

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. provisional patent applications 60/506,181 filed on September 26, 2003, entitled "c-Met Modulators and Method of Use," naming Bannen, Lynne *et. al* as inventors, 60/535,377 filed on January 9, 2004, entitled "c-Met Modulators and Method of Use," naming Bannen, Lynne *et. al* as inventors and 60/577,384 filed on June 4, 2004, entitled "Synthesis of Quinoline and Quinazoline Kinase Modulators," naming Bannen, Lynne *et. al* as inventors; each of which is hereby incorporated by reference in its entirety for all purposes.

FIELD OF THE INVENTION

[0002] This invention relates to compounds for modulating protein kinase enzymatic activity for modulating cellular activities such as proliferation, differentiation, programmed cell death, migration and chemoinvasion. Even more specifically, the invention relates to quinazolines and quinolines which inhibit, regulate and/or modulate kinase receptor signal transduction pathways related to the changes in cellular activities as mentioned above, compositions which contain these compounds, methods of using them to treat kinase-dependent diseases and conditions, synthesis of the compounds as well as processes for formulating the compounds for pharmaceutical purposes.

BACKGROUND OF THE INVENTION

[0003] Improvements in the specificity of agents used to treat cancer is of considerable interest because of the therapeutic benefits which would be realized if the side effects associated with the administration of these agents could be reduced. Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms.

[0004] Protein kinases are enzymes that catalyze the phosphorylation of proteins, in particular, hydroxy groups on tyrosine, serine and threonine residues of proteins. The consequences of this seemingly simple activity are staggering; cell differentiation and proliferation; i.e., virtually all aspects of cell life in one-way or another depend on protein kinase activity. Furthermore, abnormal protein kinase activity has been related to a host of

disorders, ranging from relatively non-life threatening diseases such as psoriasis to extremely virulent diseases such as glioblastoma (brain cancer).

[0005] Protein kinases can be categorized as receptor type or non-receptor type. Receptor-type tyrosine kinases have an extracellular, a transmembrane, and an intracellular portion, while non-receptor type tyrosine kinases are wholly intracellular.

[0006] Receptor-type tyrosine kinases are comprised of a large number of transmembrane receptors with diverse biological activity. In fact, about 20 different subfamilies of receptor-type tyrosine kinases have been identified. One tyrosine kinase subfamily, designated the HER subfamily, is comprised of EGFR (HER1), HER2, HER3, and HER4. Ligands of this subfamily of receptors identified so far include epithelial growth factor, TGF-alpha, amphiregulin, HB-EGF, betacellulin and heregulin. Another subfamily of these receptor-type tyrosine kinases is the insulin subfamily, which includes INS-R, IGF-IR, and IR-R. The PDGF subfamily includes the PDGF-alpha and beta receptors, CSFIR, c-Kit and FLK-II. Then there is the FLK family, which is comprised of the kinase insert domain receptor (KDR), fetal liver kinase-1 (FLK-1), fetal liver kinase-4 (FLK-4) and the fms-like tyrosine kinase-1 (flt-1). The PDGF and FLK families are usually considered together due to the similarities of the two groups. For a detailed discussion of the receptor-type tyrosine kinases, see Plowman et al., DN&P 7(6): 334-339, 1994, which is hereby incorporated by reference.

[0007] The non-receptor type of tyrosine kinases is also comprised of numerous subfamilies, including Src, Frk, Btk, Csk, Abl, Zap70, Fes/Fps, Fak, Jak, Ack, and LIMK. Each of these subfamilies is further sub-divided into varying receptors. For example, the Src subfamily is one of the largest and includes Src, Yes, Fyn, Lyn, Lck, Blk, Hck, Fgr, and Yrk. The Src subfamily of enzymes has been linked to oncogenesis. For a more detailed discussion of the non-receptor type of tyrosine kinases, see Bolen, Oncogene, 8:2025-2031 (1993), which is hereby incorporated by reference.

[0008] Since protein kinases and their ligands play critical roles in various cellular activities, deregulation of protein kinase enzymatic activity can lead to altered cellular properties, such as uncontrolled cell growth associated with cancer. In addition to oncological indications, altered kinase signaling is implicated in numerous other pathological diseases. These include, but are not limited to: immunological disorders, cardiovascular diseases, inflammatory diseases, and degenerative diseases. Therefore,

both receptor and non-receptor protein kinases are attractive targets for small molecule drug discovery.

[0009] One particularly attractive goal for therapeutic use of kinase modulation relates to oncological indications. For example, modulation of protein kinase activity for the treatment of cancer has been demonstrated successfully with the FDA approval of Gleevec® (imatinib mesylate, produced by Novartis Pharmaceutical Corporation of East Hanover, NJ) for the treatment of Chronic Myeloid Leukemia (CML) and gastrointestinal stroma cancers (GIST). Gleevec is a c-Kit and Abl kinase inhibitor.

[0010] Modulation (particularly inhibition) of cell proliferation and angiogenesis, two key cellular processes needed for tumor growth and survival (Matter A. Drug Disc Technol 2001 6, 1005-1024), is an attractive goal for development of small-molecule drugs. Anti-angiogenic therapy represents a potentially important approach for the treatment of solid tumors and other diseases associated with dysregulated vascularization, including ischemic coronary artery disease, diabetic retinopathy, psoriasis and rheumatoid arthritis. As well, cell antiproliferative agents are desirable to slow or stop the growth of tumors.

[0011] One particularly attractive target for small-molecule modulation, with respect to antiangiogenic and antiproliferative activity is c-Met. The kinase, c-Met, is the prototypic member of a subfamily of heterodimeric receptor tyrosine kinases (RTKs) which include Met, Ron and Sea. Expression of c-Met occurs in a wide variety of cell types including epithelial, endothelial and mesenchymal cells where activation of the receptor induces cell migration, invasion, proliferation and other biological activities associated with "invasive cell growth." As such, signal transduction through c-Met receptor activation is responsible for many of the characteristics of tumor cells.

[0012] The endogenous ligand for c-Met is the hepatocyte growth factor (HGF), a potent inducer of angiogenesis, also known as "scatter factor" (SF). Binding of HGF to c-Met induces activation of the receptor via autophosphorylation resulting in an increase of receptor dependent signaling, which promotes cell growth and invasion. Anti-HGF antibodies or HGF antagonists have been shown to inhibit tumor metastasis *in vivo* (See: Maulik et al Cytokine & Growth Factor Reviews 2002 13, 41-59).

[0013] Tumor growth progression requires the recruitment of new blood vessels into the tumor from preexisting vessels as well as invasion, adhesion and proliferation of malignant cells. Accordingly, c-Met overexpression has been demonstrated on a wide

variety of tumor types including breast, colon, renal, lung, squamous cell myeloid leukemia, hemangiomas, melanomas, astrocytomas, and glioblastomas. Additionally activating mutations in the kinase domain of c-Met have been identified in hereditary and sporadic renal papilloma and squamous cell carcinoma. (See: Maulik et al Cytokine & growth Factor reviews 2002 13, 41-59; Longati et al Curr Drug Targets 2001, 2, 41-55; Funakoshi et al Clinica Chimica Acta 2003 1-23). Thus modulation of c-Met is desirable as a means to treat cancer and cancer-related disease.

[0014] The Eph receptors comprise the largest family of receptor tyrosine kinases and are divided into two groups, EphA and EphB, based on their sequence homology. The ligands for the Eph receptors are ephrin, which are membrane anchored. Ephrin A ligands bind preferentially to EphA receptors whilst ephrin B ligands bind to EphB receptors. Binding of ephrin to Eph receptors causes receptor autophosphorylation and typically requires a cell-cell interaction since both receptor and ligand are membrane bound.

[0015] Overexpression of Eph receptors has been linked to increased cell proliferation in a variety of tumors (Zhou R 1998 Pharmacol Ther. 77, 151-181; Kiyokawa E, Takai S, Tanaka M et al 1994 Cancer Res 54, 3645-3650; Takai N Miyazaki T, Fujisawa K, Nasu K and Miyakawa. 2001 Oncology reports 8, 567-573). The family of Eph receptor tyrosine kinases and their ephrin ligands play important roles in a variety of processes during embryonic development and also in pathological angiogenesis and potentially metastasis. Therefore modulation of Eph receptor kinase activity should provide means to treat or prevent disease states associated with abnormal cell proliferation such as those described above.

[0016] Inhibition of EGF, VEGF and ephrin signal transduction will prevent cell proliferation and angiogenesis, two key cellular processes needed for tumor growth and survival (Matter A. Drug Disc. Technol. 2001 6, 1005-1024). EGF and VEGF receptors are previously described targets for small molecule inhibition. KDR and flt-4 are both VEGF receptors

[0017] One particularly attractive target for small-molecule modulation is c-Kit. The proto-oncogene *c-Kit* was first identified as the oncogenic component of the acutely transforming Hardy-Zuckerman 4-feline sarcoma virus (Besmer et al Nature 1986 320:415-421). *c-Kit* (also called stem cell factor receptor or steel factor receptor) is a type 3 receptor tyrosine kinase (RTK) belonging to the platelet-derived growth factor receptor subfamily. *c-Kit* binds the ligand stem cell factor (SCF), and triggers its multiple signal

transduction pathways including Src family kinases, phosphatidyl-inositol 3 kinase, the Ras-Raf-Map kinase cascade, and phospholipase C (Broudy et al Blood 1999 94: 1979-1986; Lennartsson et al Oncogene 1999 18: 5546-5553 ; Timokhina et al EMBO J 1998 17:6250-6262; Chian et al Blood 2001 98(5)1365-1373; Blume-Jensen et al Curr Biol 1998 8:779-782; Kissel et al EMBO J 2000 19:1312-1326; Lennartsson et al. Oncogene 1999 18: 5546-5553; Sue et al Blood, 1998 92:1242-1149; Lev et al EMBO J 1991 10:647-654). c-Kit is required for normal hematopoiesis, melanonogenesis, and gametogenesis. c-Kit is expressed in mast cells, immature myeloid cells, melanocytes, epithelial breast cells and the interstitial cells of Cajal (ICC). In mast cells, it is required not only for the differentiation, maturation, chemotaxis, and haptotaxis but also for the promotion of survival and proliferation.

[0018] Mutations in c-Kit have been implicated in human disease. Mutations in the juxtamembrane domain are found in many human gastrointestinal stromal tumors, and mutations in the kinase domain are found in mastocytosis, germ cell tumors, acute myeloid leukemia (AML), NK lymphoma, and other hematologic disorders (Hirota et al Science 1998 279:577-580; Singer et al J Clin Oncol 2002 20:3898-3905; Longley et al Proc Natl Aca Sci USA 1999: 1609-1614; Tian et al Am J Pathol 1999 154: 1643-1647; Beghini et al Blood 2000 95:726-727; Hongyo et al Cancer Res 2000 60:2345-2347). These mutations result in ligand-independent tyrosine kinase activity, autophosphorylation of c-Kit, uncontrolled cell proliferation, and stimulation of downstream signaling pathways. Overexpression of c-Kit and c-Kit ligand have also been described in other tumors including small-cell lung cancer, neuroblastomas, gynecological tumors, and colon carcinoma, which might result in autocrine or paracrine c-Kit activation.

[0019] The overexpression of c-Kit has also been implicated in the development of neoplasia associated with neurofibromatosis type 1 (NF1). Mutations in the tumor suppressor gene *NF1* lead to a deficiency in neurofibromin, a GTPase-activating protein for Ras. This deficiency results in abnormal proliferation of Schwann cells in the peripheral nervous system, and predisposes affected individuals to peripheral nerve sheath tumors (neurofibromas), astrocytomas (optic pathway gliomas), learning disabilities, seizures, strokes, macrocephaly, vascular abnormalities, and juvenile myelomonocytic leukemia (Lynch & Gutmann Neurol Clin 2002 20:841-865). Genetic experiments in mice demonstrate that haploinsufficiency at *NF1* partially rescues some of the phenotypes associated with mutations in the gene for c-Kit, indicating that these genes function along

a common developmental pathway (Ingram, et al. J. Exp Med 2000 191:181-187). Also, c-Kit is expressed in schwannoma cells from NF1 patients, but not in normal schwann cells (Ryan et al. J Neurosci Res 1994 37:415-432). These data indicate that elevated c-Kit expression and sensitivity to stem cell factor may play important roles in the development of proliferative disorders associated with NF-1. Therefore, c-Kit inhibitors may be effective chemotherapeutic agents for treating patients with NF-1.

[0020] GISTs are the most common mesenchymal tumors of the gastrointestinal tract, and they are generally resistant to chemotherapy and radiation therapy. However, recent results with the c-Kit/BCR-Abl inhibitor STI571 indicate that targeting c-Kit may be an effective therapeutic strategy for this disease (Eisenberg & Mehren Expert Opin Pharmacother 2003 4:869-874). Malignant mast cell disease often suggests an extremely poor prognosis, and no reliable effective chemotherapeutic agents have been identified (Marone et al Leuk Res 2001 25:583-594). Systemic mast cell disorders have been treated with interferon-alpha, although the effectiveness of this therapy has been variable (Lehmann & Lammle Ann Hematol 1999 78:483-484; Butterfield Br J Dermatol 1998 138: 489-495). Therefore, activated c-Kit might serve as a therapeutic target in GISTs and mast cell disease, as well as other disorders associated with activated c-Kit.

[0021] Flt-3 is normally expressed on hematopoietic progenitor cells and a subset of mature myeloid and lymphoid cells, where it modulates cell survival and proliferation. Flt-3 is constitutively activated via mutation, either in the juxtamembrane region or in the activation loop of the kinase domain, in a large proportion of patients with AML (Reilly Leuk Lymphoma 2003 44: 1-7). Also, mutations in flt-3 are significantly correlated with poor prognosis in AML patients (Sawyers Cancer Cell 2002 1: 413-415).

[0022] Accordingly, the identification of small-molecule compounds that specifically inhibit, regulate and/or modulate the signal transduction of kinases, particularly including c-Met, KDR, c-Kit, flt-3, and flt-4, is desirable as a means to treat or prevent disease states associated with abnormal cell proliferation and angiogenesis, and is an object of this invention.

[0023] Quinolines and quinazolines bearing substitution, for example at the two, four, six and seven positions of their fused ring system have been shown to be particularly attractive targets for kinase inhibition by a number of groups. Conventional quinoline and quinazoline kinase inhibitors typically have fairly simple substitution about the quinoline or quinazoline fused ten-membered ring system, but recently more complex molecules are

being disclosed. For example, we have previously disclosed, in U.S. provisional patent applications 60/506,181 and 60/535,377 which are both incorporated by reference herein in their entirety for all purposes, that certain quinolines and quinazolines are particularly well suited as kinase modulators, more particularly inhibitors of for example c-Met, KDR, c-Kit, flt-3, and flt-4. These molecules in some cases are particularly complex and although they can be made via conventional methods, more efficient routes are desirable, especially in a pharmaceutical setting.

[0024] Conventional methods of making quinolines and quinazolines with the aforementioned substitution patterns usually involve linear construction of a quinoline or quinazoline template upon which relatively simple substitutions are appended. With the advent of more complex substitution about such quinolines and quinazolines (*vide supra*), for example side chains containing cyclic and bicyclic systems with multiple functional groups, conventional methods of synthesis become problematic due to the linear or serial reactions used. Indeed, as such molecules become more complex and the utility of such complex groups is realized, the quinoline and quinazoline ring system becomes more of a sub-structure than a main structure of such inhibitors. Thus it is desirable to find more efficient methods of synthesis, particularly convergent syntheses which are an object of this invention.

SUMMARY OF THE INVENTION

[0025] In one aspect, the present invention provides compounds for modulating kinase activity and methods of treating diseases mediated by kinase activity utilizing the compounds and pharmaceutical compositions thereof. Diseases mediated by kinase activity include, but are not limited to, diseases characterized in part by migration, invasion, proliferation and other biological activities associated with invasive cell growth. In particular to this invention is modulation, even more particularly inhibition, of c-Met, KDR, c-Kit, flt-3, and flt-4.

[0026] In another aspect, the invention provides methods of screening for modulators of c-Met, KDR, c-Kit, flt-3, and flt-4 activity. The methods comprise combining a composition of the invention, a kinase, e.g. c-Met, KDR, c-Kit, flt-3, or flt-4, and at least one candidate agent and determining the effect of the candidate agent on the c-Met, KDR, c-Kit, flt-3, or flt-4, activity.

[0027] In yet another aspect, the invention also provides pharmaceutical kits comprising one or more containers filled with one or more of the ingredients of pharmaceutical

compounds and/or compositions of the present invention, including, one or more kinase, e.g. c-Met, KDR, c-Kit, flt-3, or flt-4, enzyme activity modulators as described herein. Such kits can also include, for example, other compounds and/or compositions (e.g., diluents, permeation enhancers, lubricants, and the like), a device(s) for administering the compounds and/or compositions, and written instructions in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which instructions can also reflect approval by the agency of manufacture, use or sale for human administration.

[0028] In another aspect, the invention also provides a diagnostic agent comprising a compound of the invention and, optionally, pharmaceutically acceptable adjuvants and excipients.

[0029] In still yet another aspect, the present invention provides processes for making compounds, and pharmaceutical compositions thereof, for modulating kinase activity and treating diseases mediated by kinase activity. In particular to this invention are methods for making quinolines and quinazolines used for modulation of kinase activity, even more particularly inhibition of kinase activity, and yet even more particularly inhibition of c-Met, KDR, c-Kit, flt-3, and flt-4.

[0030] These and other features and advantages of the present invention will be described in more detail below with reference to the associated drawings.


DETAILED DESCRIPTION OF THE INVENTION

[0031] The compositions of the invention are used to treat diseases associated with abnormal and or unregulated cellular activities. Disease states which can be treated by the methods and compositions provided herein include, but are not limited to, cancer (further discussed below), immunological disorders such as rheumatoid arthritis, graft-host diseases, multiple sclerosis, psoriasis; cardiovascular diseases such as arteriosclerosis, myocardioinfarction, ischemia, stroke and restenosis; other inflammatory and degenerative diseases such as interbowel diseases, osteoarthritis, macular degeneration, diabetic retinopathy.

[0032] It is appreciated that in some cases the cells may not be in a hyper- or hypo-proliferative and/or migratory state (abnormal state) and still require treatment. For example, during wound healing, the cells may be proliferating "normally", but

proliferation and migration enhancement may be desired. Alternatively, reduction in "normal" cell proliferation and/or migration rate may be desired.

[0033] Thus, in one aspect the present invention comprises a compound for modulating kinase activity according to Formula I,





or a pharmaceutically acceptable salt, hydrate, or prodrug thereof, wherein,

R^1 is selected from -H, halogen, $-OR^3$, $-NO_2$, $-NH_2$, $-NR^3R^4$, and optionally substituted lower alkyl;

A^1 is selected from $=N-$, $=C(H)-$, and $=C(CN)-$;

Z is selected from $-S(O)_{0-2}-$, $-O-$, and $-NR^5-$;

Ar is either a group of formula II, or of formula III,



wherein,

R^2 is selected from -H, halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted lower alkyl;

q is 0 to 4;

G is a group -B-L-T, wherein

B is selected from absent, $-N(R^{13})-$, $-N(SO_2R^{13})-$, $-O-$, $-S(O)_{0-2}-$, and $-C(=O)-$;

L is selected from absent, $-C(=S)N(R^{13})-$, $-C(=NR^{14})N(R^{13})-$, $-SO_2N(R^{13})-$, $-SO_2-$, $-C(=O)N(R^{13})-$, $-N(R^{13})-$, $-C(=O)C_{1-2}alkylN(R^{13})-$, $-N(R^{13})C_{1-2}alkylC(=O)-$, $-C(=O)C_{0-1}alkylC(=O)N(R^{13})-$, $-C_{0-4}alkylene-$, $-C(=O)C_{0-1}alkylC(=O)OR^3-$,

-C(=NR¹⁴)C₀₋₁alkylC(=O)-, -C(=O)-, -C(=O)C₀₋₁alkylC(=O)-, and an optionally substituted four to six-membered heterocyclyl containing between one and three annular heteroatoms including at least one nitrogen; and

T is selected from -H, -R¹³, -C₀₋₄alkyl, -C₀₋₄alkylQ, -OC₀₋₄alkylQ, -C₀₋₄alkylOQ, -N(R¹³)C₀₋₄alkylQ, -SO₂C₀₋₄alkylQ, -C(=O)C₀₋₄alkylQ, -C₀₋₄alkylN(R¹³)Q, and -C(=O)N(R¹³)C₀₋₄alkylQ, wherein each of the aforementioned C₀₋₄alkyl is optionally substituted;

J is selected from -S(O)₀₋₂-, -O-, and -NR¹⁵-;

R³ is -H or R⁴;

R⁴ is selected from optionally substituted lower alkyl, optionally substituted aryl, optionally substituted lower arylalkyl, optionally substituted heterocyclyl, and optionally substituted lower heterocyclylalkyl; or


R³ and R⁴, when taken together with a common nitrogen to which they are attached, form an optionally substituted five- to seven-membered heterocyclyl, said optionally substituted five- to seven-membered heterocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

A² and A³ are each independently selected from =N-, =C(R²)-;

R⁵ is -H or optionally substituted lower alkyl;

D is selected from -O-, -S(O)₀₋₂-, and -NR¹⁵-;

R⁵⁰ is either R³, or according to formula IV;



IV

wherein X¹, X², and optionally X³, represent the atoms of a saturated bridged ring system, said saturated bridged ring system comprising up to four annular heteroatoms represented by any of X¹, X², and X³; wherein,

each X¹ is independently selected from -C(R⁶)R⁷-, -O-, -S(O)₀₋₂-, and -NR⁸-;

each X^2 is independently an optionally substituted bridgehead methine or a bridgehead nitrogen;

each X^3 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

Y is either:

an optionally substituted lower alkylene linker, between D and either 1) any annular atom of the saturated bridged ring system, except X^2 when X^2 is a bridgehead nitrogen, or 2) any heteroatom, represented by any of R^6 or R^7 ; provided there are at least two carbon atoms between D and any annular heteroatom of the saturated bridged ring system or any heteroatom represented by any of R^6 or R^7 ;

or Y is absent, when Y is absent, said saturated bridged ring system, is directly attached to D via an annular carbon of said saturated bridged ring system, unless D is $-SO_2-$, in which case said saturated bridged ring system, is directly attached to D via an any annular atom of said saturated bridged ring system;

m and p are each independently 1-4;

n is 0-2, when n = 0, then there is a single bond between the two bridgehead X^2 's;

R^6 and R^7 are each independently selected from -H, halogen, trihalomethyl, -CN, $-NH_2$, $-NO_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^4$, $-SO_2NR^3R^4$, $-CO_2R^3$, $-C(O)NR^3R^4$, $-N(R^3)SO_2R^4$, $-N(R^3)C(O)R^3$, $-NCO_2R^3$, $-C(O)R^3$, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted lower arylalkyl, optionally substituted heterocyclyl, optionally substituted lower heterocyclylalkyl, and a bond to either Y or D; or

R^6 and R^7 , when taken together are oxo; or

R^6 and R^7 , when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

R^8 is selected from $-R^3$, Y, $-SO_2NR^3R^4$, $-CO_2R^4$, $-C(O)NR^3R^3$, $-SO_2R^4$, and $-C(O)R^3$;

R^{13} is selected from -H, $-C(=O)R^3$, $-C(=O)OR^3$, $-C(=O)SR^3$, $-SO_2R^4$, $-C(=O)N(R^3)R^3$, and optionally substituted lower alkyl,

two R^{13} , together with the atom or atoms to which they are attached, can combine to form a heteroalicyclic optionally substituted with between one and four of R^{60} , said heteroalicyclic can have up to four annular heteroatoms, and said heteroalicyclic can have an aryl or heteroaryl fused thereto, in which case said aryl or heteroaryl is optionally substituted with an additional one to four of R^{60} ;

R^{14} is selected from -H, -NO₂, -NH₂, -N(R³)R⁴, -CN, -OR³, optionally substituted lower alkyl, optionally substituted heteroalicycylalkyl, optionally substituted aryl, optionally substituted arylalkyl and optionally substituted heteroalicyclic;

R^{15} is a group -M¹-M², wherein M¹ is selected from absent, -C(=S)N(R¹³)-, -C(=NR¹⁴)N(R¹³)-, -SO₂N(R¹³)-, -SO₂-, -C(=O)N(R¹³)-, -C(=O)C(=O)N(R¹³)-, -C₀₋₄alkylene-, -C(=O)-, and an optionally substituted four to six-membered heterocyclannular containing between one and three heteroatoms including at least one nitrogen; and M² is selected from -H, -C₀₋₆alkyl, alkoxy, -C(=O)C₀₋₄alkylQ, -C₀₋₄alkylQ, -OC₀₋₄alkylQ-, -N(R¹³)C₀₋₄alkylQ-, and -C(=O)N(R¹³)C₀₋₄alkylQ; and

Q is a five- to ten-membered ring system, optionally substituted with between zero and four of R^{20} ;

R^{20} is selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl;

R^{60} is selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted heteroarylalkyl, and optionally substituted arylalkyl;

two of R^{60} , when attached to a non-aromatic carbon, can be oxo;


with the proviso, only when Ar is according to formula II, if Y is a C₁₋₆ alkylene; Z is -NH- or -N(CH₃)-; R¹ is a C₁₋₆alkyl optionally substituted in the 2-position by -OH or a C₁₋₄alkoxy group; R² is -H or halogen; n = 0; and the atoms, X¹, of one bridge of the saturated bridged ring system, when combined with both bridgehead atoms, X², of the saturated bridged ring system, represent:

- 1) either a pyrrolidine or a piperidine, and any atom, X¹ or X², of either of said pyrrolidine or said piperidine is attached to Y, then the other bridge of said saturated bridged ring system cannot be any one of -OC(O)CH₂-,

-CH₂OC(O)-, -OC(O)CH₂CH₂-, -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, -OC(O)CH₂NH-, -OC(O)CH₂N(C₁₋₄alkyl)-, and -OC(O)CH₂O-; or

- 2) either a piperazine or a 4-(C₁₋₄alkyl)-piperazine, and any atom, X¹ or X², of either of said piperazine or said 4-(C₁₋₄alkyl)-piperazine is attached to Y, then the other bridge of said saturated bridged ring system, only when attached via the 2- and the 3-position of either of said piperazine or said 4-(C₁₋₄alkyl)-piperazine, cannot be one of -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, and either of the two aforementioned bridges optionally substituted by one or two C₁₋₂alkyl groups; or
- 3) a piperazine, and any atom, X¹ or X², of said piperazine is attached to Y, then the other bridge of said saturated bridged ring system, only when attached via the 3- and the 4-position of said piperazine, cannot be one of -C(O)OCH₂CH₂-, -CH₂OC(O)CH₂-, and either of the two aforementioned bridges optionally substituted by one or two C₁₋₂alkyl groups, and only when either of the two aforementioned bridges are attached to the 3-position of said piperazine via their left-hand end as depicted above; or
- 4) a 2-oxomorpholine, said 2-oxomorpholine attached to Y via its 4-position, then the other bridge of said saturated bridged ring system, only when attached via the 5- and the 6-position of said 2-oxomorpholine, cannot be one of -(CH₂)_g-, -CH₂WCH₂-, -CH₂WCH₂CH₂-, and -CH₂CH₂WCH₂-, wherein W is -O-, -S(O)₀₋₂-, -NH-, or -N(C₁₋₄alkyl)- wherein g is 2, 3, or 4;


and with the proviso that when Z is -O-, Ar is according to formula II, and the portion of G directly attached to Ar is selected from:



then R^{50} must be of formula IV;


and with the proviso that when Ar is phenylene or substituted phenylene, Z is $-S(O)_{0-2}-$ or


$-O-$, then the portion of G directly attached to Ar cannot contain
when R^{70} is selected from $-H$, C_{1-4} alkyl, and C_{1-4} alkoxyl.



[0034] In one example, the compound is according to paragraph [0033], wherein Z is either $-O-$ or $-NR^5-$.




[0035] In another example, the compound is according to paragraph [0034], wherein G is selected from the following:





wherein wherein Q, R²⁰, and R¹³ are as defined above; each E is selected from -O-, -N(R¹³)-, -CH₂-, and -S(O)₀₋₂-; M is selected from -O-, -N(R¹³)-, -CH₂-, and -C(=O)N(R¹³)-; each V is independently either =N- or =C(H)-; each methylene in any of the above formulae is independently optionally substituted with R²⁵; and R²⁵ is selected from halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic, two of R²⁵ on a single carbon can be oxo.

[0036] In another example, the compound is according to paragraph [0035], wherein Ar is according to one of formula **IIa**, **IIb**, and **IIIa**.

**IIa****IIb****IIIa**


[0037] In another example, the compound is according to paragraph [0036], wherein D is -O- and R¹ is -OR³.

[0038] In another example, the compound is according to paragraph [0037], wherein -O-R⁵⁰ and R¹ are interchangeably located at the 6-position and 7-position of the quinazoline or quinoline according to formula I.

[0039] In another example, the compound is according to paragraph [0038], wherein R¹ is -OH or -OC₁₋₆alkyl.


[0040] In another example, the compound is according to paragraph [0039], wherein A^1 is $=N-$ or $=C(H)-$.

[0041] In another example, the compound is according to paragraph [0040], wherein G is selected from:




wherein Q, R^{20} , R^{13} , E, and R^{60} are as defined above; each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R^{25} ; and R^{25} is selected from halogen, trihalomethyl, oxo, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic.

[0042] In another example, the compound is according to paragraph [0041], wherein Q is selected from:




wherein R^{20} is defined as above, and P is a five- to seven-membered ring, including the two shared carbons of the aromatic ring to which P is fused, P optionally containing between one and three heteroatoms.


[0043] In another example, the compound is according to paragraph [0042], wherein Ar is according to formula IIa, and G is selected from:



wherein Q, R^{20} , R^{13} , E, and R^{60} are as defined above, and each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R^{25} ; and R^{25} is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic.

[0044] In another example, the compound is according to paragraph [0042], wherein Ar is according to formula IIb, and G is selected from:





wherein Q, R^{20} , R^{13} , E, and R^{60} are as defined above, and each methylene in any of the above formulae, other than those depicted in a ring, is independently optionally substituted with R^{25} ; and R^{25} is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic.

[0045] In another example, the compound is according to paragraph [0044], wherein the methylene between the two carbonyls of the depicted formulae is di-substituted with either optionally substituted lower alkyl, or an optionally substituted spirocycle.

[0046] In another example, the compound is according to either [0043] or paragraph [0044], wherein R^{50} is a heteroalicyclic or a C₁₋₆alkyl-heteroalicyclic.

[0047] In another example, the compound is according to paragraph [0046], wherein at least one of R^2 is halogen.

[0048] In another example, the compound is according to paragraph [0046], wherein R^{50} is according to formula IV.

[0049] In another example, the compound is according to paragraph [0048], wherein the saturated bridged ring system according to formula IV has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], [3.1.0], [3.3.3], [3.3.2], [3.3.1], [3.2.2], [3.2.1], [2.2.2], and [2.2.1].


[0050] In another example, the compound is according to paragraph [0049], wherein Y is selected from -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂-, -CH₂-, and absent.

[0051] In another example, the compound is according to paragraph [0050], wherein n is 0 and the saturated bridged ring system according to formula IV has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], and [3.1.0].

[0052] In another example, the compound is according to paragraph [0051], wherein said saturated bridged ring system contains at least one annular nitrogen or at least one annular oxygen.

[0053] In another example, the compound is according to paragraph [0052], wherein said saturated bridged ring system contains $\text{-NR}^8\text{-}$, wherein R^8 is selected from -H , optionally substituted lower alkyl, $\text{-CO}_2\text{R}^3$, $\text{-C(O)NR}^3\text{R}^3$, $\text{-SO}_2\text{R}^3$, and -C(O)R^3 .

[0054] In another example, the compound is according to paragraph [0052], wherein said saturated bridged ring system is of formula V,



V

wherein U^1 is selected from -O- , $\text{-S(O)}_{0-2}\text{-}$, $\text{-NR}^8\text{-}$, $\text{-CR}^6\text{R}^7\text{-}$, and absent; and e is 0 or 1.

[0055] In another example, the compound is according to paragraph [0054], wherein Y is $\text{-CH}_2\text{-}$.


[0056] In another example, the compound is according to paragraph [0055], wherein U^1 is $\text{-NR}^8\text{-}$, wherein R^8 is selected from -H , optionally substituted lower alkyl, $\text{-CO}_2\text{R}^3$, $\text{-C(O)NR}^3\text{R}^3$, $\text{-SO}_2\text{R}^3$, and -C(O)R^3 .

[0057] In another example, the compound is according to paragraph [0055], wherein U^1 is -O- .

[0058] In another example, the compound is according to paragraph [0055], wherein U^1 is absent.

[0059] In another example, the compound is according to paragraph [0052], wherein Y is selected from $\text{-CH}_2\text{CH}_2\text{-}$, $\text{-CH}_2\text{-}$, and absent.

[0060] In another example, the compound is according to paragraph [0059], wherein said saturated bridged ring system is of formula VI,



VI

wherein R^9 , R^{10} , and R^{11} are each independently selected from -H, and $-OR^{12}$; or

R^9 is selected from -H, and $-OR^{12}$, and R^{10} and R^{11} , when taken together, are either an optionally substituted alkylidene or an oxo;

R^{12} is selected from -H, $-C(O)R^3$, optionally substituted lower alkylidyne, optionally substituted lower arylalkylidyne, optionally substituted lower heterocyclalkylidyne, optionally substituted lower alkylidene, optionally substituted lower alkylidenearyl, optionally substituted lower alkylideneheterocycl, optionally substituted lower alkyl, optionally substituted lower alkylaryl, optionally substituted aryl, optionally substituted lower heterocyclalkyl, and optionally substituted heterocycl;


or two R^{12} 's, when taken together, form 1) a corresponding spirocyclic ketal when said two R^{12} 's stem from R^{10} and R^{11} , or 2) a corresponding cyclic ketal when said two R^{12} 's stem from R^9 and one of R^{10} and R^{11} .

[0061] In another example, the compound is according to paragraph [0060], wherein one of R^{10} and R^{11} is $-OR^{12}$, wherein R^{12} is selected from -H, $-C(O)R^3$, and optionally substituted lower alkyl; and R^9 and the other of R^{10} and R^{11} are both -H.

[0062] In another example, the compound is according to paragraph [0061], wherein Y is either $-CH_2-$ or absent.

[0063] In another example, the compound is according to paragraph [0062], wherein R^9 is an alkyl group containing at least one fluorine substitution thereon.

[0064] In another example, the compound is according to paragraph [0053], wherein said saturated bridged ring system is of formula VII.



VII

[0065] In another example, the compound is according to paragraph [0064], wherein Y is either $-CH_2-$ or absent.

[0066] In another example, the compound is according to paragraph [0065], wherein R^8 is methyl or ethyl.

[0067] In another example, the compound is according to paragraph [0053], wherein said saturated bridged ring system is of formula VIII.




VIII

[0068] In another example, the compound is according to paragraph [0067], wherein Y is -CH₂-.

[0069] In another example, the compound is according to paragraph [0068], wherein R⁸ is methyl or ethyl.

[0070] In another example, the compound is according to paragraph [0052], wherein said saturated bridged ring system is of formula IX



IX

wherein U² is selected from -O-, -S(O)₀₋₂-, -NR⁸-, -CR⁶R⁷-, and absent.

[0071] In another example, the compound is according to paragraph [0070], wherein R³ of formula IX is selected from -H and optionally substituted alkyl.

[0072] In another example, the compound is according to paragraph [0071], wherein U² is either -CR⁶R⁷- or absent.

[0073] In another example, the compound is according to paragraph [0072], wherein U² is either -CH₂- or absent.

[0074] In another example, the compound is according to paragraph [0073], wherein Y is -CH₂-.

[0075] In another example, the compound is according to paragraph [0053], wherein said saturated bridged ring system is according to formula X.



X

[0076] In another example, the compound is according to paragraph [0075], wherein R⁸ is methyl or ethyl.

[0077] In another example, the compound is according to paragraph [0033], selected from Table 1.

Table 1

Entry	Name	Structure
1	N-[(3-fluoro-4-[(6-(methyloxy)-7-[(3aR,6aS)-octahydrocyclopenta[c]pyrrol-5-ylmethyl]oxy)quinazolin-4-yl]oxy]phenyl]amino)carbonothioyl]-2-phenylacetamide	
2	N-{[(3-fluoro-4-{[7-((3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl]oxy)-6-(methyloxy)quinazolin-4-yl]oxy}phenyl]amino}carbonothioyl]-2-phenylacetamide	
3	N-[(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)(methyl)amino]carbonothioyl]-2-phenylacetamide	
4	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)imidazolidin-2-one	
5	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-(phenylmethyl)imidazolidin-2-one	

Table 1

Entry	Name	Structure
6	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-(phenylacetyl)imidazolidin-2-one	
7	ethyl [(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)amino](oxo)acetate	
8	N-{[(4-{[6,7-bis(methyloxy)quinazolin-4-yl]amino}-3-fluorophenyl)amino]carbonothioyl}-2-phenylacetamide	
9	N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N-methyl-N-(2-phenylethyl)sulfamide	
10	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-(phenylmethyl)-1,2,4-oxadiazol-5-amine	
11	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)piperidin-2-one	

Table 1

Entry	Name	Structure
12	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(phenylmethyl)ethanediamide	
13	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-4-phenyl-1,3-thiazol-2-amine	
14	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-phenylethyl)ethanediamide	
15	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-1-phenylmethanesulfonamide	
16	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-2-phenylethanesulfonamide	
17	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(phenylmethyl)benzenesulfonamide	

Table 1

Entry	Name	Structure
18	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(phenylmethyl)benzenesulfonamide}	
19	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(2-phenylethyl)benzenesulfonamide}	
20	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(2-phenylethyl)benzenesulfonamide}	
21	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(3-phenylpropyl)benzenesulfonamide}	
22	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}pyrrolidin-2-one)	
23	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl (phenylmethyl)carbamate}	

Table 1

Entry	Name	Structure
24	4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl (2-phenylethyl)carbamate	
25	4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(3-phenylpropyl)benzenesulfonamide	
26	N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-phenylethanediarnide	
27	N-[[[(3-fluoro-4-[[7-[[[(2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}phenyl)amino]carbonylthioyl]-2-phenylacetamide	
28	N-[(Z)-[(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)amino](imino)methyl]-2-phenylacetamide	
29	4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-[2-(phenyloxy)ethyl]benzenesulfonamide	

Table 1

Entry	Name	Structure
30	N,N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-bis-(3-phenylpropane-1-sulfonamide)	
31	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-phenylpropane-1-sulfonamide	
32	N2-[(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)sulfonyl]-N1-phenylglycinamide	
33	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-2-phenylacetamide	
34	N-[[6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl]amino]carbonothioyl]-2-phenylacetamide	
35	6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-1,3-benzothiazol-2-amine	

Table 1

Entry	Name	Structure
36	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-amine}	
37	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl})-2-phenylacetamide	
38	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl})-N'-(2-morpholin-4-ylethyl)ethanediamide	
39	benzyl-{{[4-(6,7-dimethoxyquinolin-4-yloxy)-3-fluorophenylcarbamoyl]-methyl}-carbamic acid tert-butyl ester}	
40	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl})-N2-(phenylmethyl)glycinamide	
41	N2-acetyl-N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl})-N2-(phenylmethyl)glycinamide	

Table 1







Entry	Name	Structure
42	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-1,3-benzothiazol-2-yl)-2-phenylacetamide	
43	benzyl-{{6-(6,7-dimethoxyquinolin-4-yloxy)-pyridin-3-ylcarbamoyl}-methyl}-carbamic acid tert-butyl ester	
44	N1-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-N2-(phenylmethyl)glycinamide	
45	N2-acetyl-N1-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-N2-(phenylmethyl)glycinamide	
46	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-3-phenylpropanamide	
47	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-4-phenylbutanamide	

Table 1

Entry	Name	Structure
48	N1-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}pyridin-3-yl)-N2-methyl-N2-(phenylmethyl)glycinamide	
49	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-{{2-[4-(methyloxy)phenyl]ethyl}ethanediamide	
50	N1-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N2-methyl-N2-(phenylmethyl)glycinamide	
51	4-[(2-amino-1,3-benzothiazol-6-yl)oxy]-6,7-bis(methyloxy)-1-(2-oxo-2-phenylethyl)quinolinium	
52	N-{{[(4-{{6,7-bis(methyloxy)quinolin-4-yl}amino}phenyl)amino]carbothioyl}-2-phenylacetamide	
53	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-1,3-benzothiazol-2-yl)-3-phenylpropanamide	

Table 1

Entry	Name	Structure
54	N-[[6-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-5-chloropyridin-3-yl]amino]carbonothioyl}-2-phenylacetamide	
55	N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-3-fluorophenyl)-N'-(2,3-dihydro-1H-inden-1-yl)ethanediamide	
56	N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-3-fluorophenyl)-N'-(2,3-dihydro-1H-inden-2-yl)ethanediamide	
57	N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-3-fluorophenyl)-N'-(1,2,3,4-tetrahydronaphthalen-1-yl)ethanediamide	
58	N'-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-3-fluorophenyl)-N-(2-phenylethyl)-N-(phenylmethyl)sulfamide	
59	N1-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-3-fluorophenyl)-N2-(trifluoroacetyl)glycinamide	

Table 1

Entry	Name	Structure
60	N-{{4-(6,7-dimethoxyquinolin-4-yloxy)-3-fluorophenylcarbamoyl}-methyl}-benzamide	
61	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl}-N'-(4-fluorophenyl)propanediamide	
62	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-[(2S)-1,2,3,4-tetrahydronaphthalen-2-yl]ethanediamide	
63	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-[2-(4-methylphenyl)ethyl]ethanediamide	
64	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-(2-phenylpropyl)ethanediamide	
65	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-[2-(4-chlorophenyl)ethyl]ethanediamide	

Table 1

Entry	Name	Structure
66	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N,N'-bis(phenylmethyl)sulfamide	
67	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N,N'-bis(2-phenylethyl)sulfamide	
68	ethyl [(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)amino](oxo)acetate	
69	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-phenylethyl)ethanediamide	
70	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)propanediamide	
71	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(1,2,3,4-tetrahydronaphthalen-2-yl)ethanediamide	

Table 1

Entry	Name	Structure
72	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(1-methylpyrrolidin-2-yl)ethyl]ethanediamide	
73	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(phenyloxy)ethyl]ethanediamide	
74	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-hydroxy-1-(phenylmethyl)ethyl]urea	
75	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-[(4-methylphenyl)sulfonyl]-4-(phenylmethyl)imidazolidin-2-one	
76	N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N-methyl-N-(2-phenylethyl)ethanediamide	
77	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[3-(trifluoromethyl)phenyl]methylethanediamide	

Table 1

Entry	Name	Structure
78	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-{2-[3-(trifluoromethyl)phenyl]ethyl}ethanediamide	
79	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloropyridin-3-yl)-3-oxo-4-phenylbutanamide	
80	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloropyridin-3-yl)-2-[3-(trifluoromethyl)phenyl]acetamide	
81	6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-N-[2-(phenyloxy)ethyl]-1,3-benzothiazol-2-amine	
82	6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-N-(2-piperidin-1-ylethyl)-1,3-benzothiazol-2-amine	
83	6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-N-methyl-N-(2-phenylethyl)-1,3-benzothiazol-2-amine	

Table 1

Entry	Name	Structure
84	6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-N-(2-pyrrolidin-1-ylethyl)-1,3-benzothiazol-2-amine	
85	6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-N-{{3-(trifluoromethyl)phenyl}methyl}-1,3-benzothiazol-2-amine	
86	6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-N-{2-[3-(trifluoromethyl)phenyl]ethyl}-1,3-benzothiazol-2-amine	
87	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloropyridin-3-yl)-N'-[3-(trifluoromethyl)phenyl]propanediamide	
88	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-fluoro-1,3-benzothiazol-2-yl)-2-[3-(trifluoromethyl)phenyl]acetamide	
89	N1-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N2-{{3-(trifluoromethyl)phenyl}methyl}glycinamide	

Table 1

Entry	Name	Structure
90	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-(2-phenylethyl)glycinamide	
91	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-{2-[3-(trifluoromethyl)phenyl]ethyl}glycinamide	
92	benzyl-{{5-chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl}carbonyl}-methyl}-carbamic acid tert-butyl ester	
93	N1-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N2-(phenylmethyl)glycinamide	
94	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-2-[3,5-bis(trifluoromethyl)phenyl]acetamide	
95	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-2-[2-chloro-5-(trifluoromethyl)phenyl]acetamide	

Table 1

Entry	Name	Structure
96	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl]oxyphenyl}-N'-(2-phenylethyl)ethanediamide	
97	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(1,2,3,4-tetrahydroisoquinolin-1-ylmethyl)ethanediamide	
98	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[(2-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)methyl]ethanediamide	
99	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-methyl-N2-{[3-(trifluoromethyl)phenyl]methyl}glycinamide	
100	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-methyl-N2-{2-[3-(trifluoromethyl)phenyl]ethyl}glycinamide	
101	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-methyl-N2-(2-phenylethyl)glycinamide	

Table 1

Entry	Name	Structure
102	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-4-(phenylmethyl)imidazolidin-2-one	
103	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridazin-3-yl)-N'-(4-fluorophenyl)propanediamide	
104	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-chlorophenyl)propanediamide	
105	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(3-chlorophenyl)propanediamide	
106	N1-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N2-methyl-N2-(phenylmethyl)glycinamide	
107	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-chlorophenyl)propanediamide	

Table 1

Entry	Name	Structure
108	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[(methyloxy)imino]propanamide	
109	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[(ethyloxy)imino]propanamide	
110	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[[phenylmethyl]oxy]imino]propanamide	
111	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-1-(phenylmethyl)prolinamide	
112	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-3-[(4-methylphenyl)sulfonyl]-4-(phenylmethyl)imidazolidin-2-one	
113	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-4-(phenylmethyl)imidazolidin-2-one	

Table 1

Entry	Name	Structure
114	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-4-(phenylmethyl)-4,5-dihydro-1,3-oxazol-2-amine	
115	6,7-bis(methyloxy)-4-({4-[4-(phenylmethyl)piperazin-1-yl]phenyl}oxy)quinoline	
116	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-4-(phenylmethyl)piperazin-2-one	
117	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N2-(phenylmethyl)alaninamide	
118	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N2-methyl-N2-(phenylmethyl)alaninamide	
119	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N2-(phenylmethyl)leucinamide	

Table 1

Entry	Name	Structure
120	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N2-methyl-N2-(phenylmethyl)leucinamide	
121	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N2-(phenylmethyl)valinamide	
122	4-(6,7-dimethoxy-quinolin-4-ylamino)-N-(3-phenylpropyl)-benzamide	
123	4-benzyl-1-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenyl]-tetrahydro-pyrimidin-2-one	
124	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	

Table 1

Entry	Name	Structure
125	2-(Benzyl-methyl-amino)-N-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenyl]-3-methyl-butylamide (note: Alphabetic order of prefixes ignored while selecting parent chain)	
126	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-phenyl]-2-phenoxyimino-propionamide	
127	2-Benzyloxyimino-N-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenyl]-2-phenyl-acetamide	
128	4-[4-(4-Benzyl-piperidin-1-yl)-phenoxy]-6,7-dimethoxy-quinoline	
129	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-N'-(2-isopropyl-1,2,3,4-tetrahydro-isoquinolin-1-ylmethyl)-oxalamide	

Table 1

Entry	Name	Structure
130	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-N'-(2-ethyl-1,2,3,4-tetrahydro-isoquinolin-1-ylmethyl)-oxalamide	
131	4-(4-{3-Chloro-5-[2-(4-fluoro-phenylcarbamoyl)-acetylamino]-pyridin-2-yloxy}-6-methoxy-quinolin-7-yloxymethyl)-piperidine-1-carboxylic acid tert-butyl ester	
132	N-{5-Chloro-6-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	
133	N-{5-Chloro-6-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	

Table 1

Entry	Name	Structure
134	N-{4-[7-(3-Diethylamino-propoxy)-6-methoxy-quinolin-4-yloxy]-3-fluoro-phenyl}-N'-phenethyl-oxalamide	
135	N-{3-Fluoro-4-[6-methoxy-7-(3-morpholin-4-yl-propoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
136	N-{3-Fluoro-4-[6-methoxy-7-(3-piperidin-1-yl-propoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
137	N-{4-[7-(2-Diethylamino-ethoxy)-6-methoxy-quinolin-4-yloxy]-3-fluoro-phenyl}-N'-phenethyl-oxalamide	

Table 1





Entry	Name	Structure
138	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-methyl-N'-phenethyl-oxalamide	
139	N-{3-Fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
140	N-{3-Fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
141	2-(3,4-Dihydro-1H-isoquinolin-2-yl)-N-{3-fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-acetamide	

Table 1





Entry	Name	Structure
142	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-2-(3-phenyl-pyrrolidin-1-yl)-acetamide	
143	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-2-(2-phenyl-morpholin-4-yl)-acetamide	
144	N-(2-Dimethylamino-2-phenyl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
145	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-oxo-2-phenyl-ethyl)-oxalamide	

Table 1

Entry	Name	Structure
146	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-2,2-difluoro-N'-(4-fluoro-phenyl)-malonamide	
147	N-Benzyl-N'-{3-fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
148	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(2-fluoro-phenyl)-ethyl]-oxalamide	
149	N-[2-(3-Chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
150	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(2-methoxy-phenyl)-ethyl]-oxalamide	
151	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-pyridin-3-yl-ethyl)-oxalamide	
152	N-Benzyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
153	N-[2-(2,5-Dimethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
154	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(2-trifluoromethyl-phenyl)-ethyl]-oxalamide	
155	N-[2-(2-Ethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
156	N-[2-(2,4-Dimethyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
157	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[(1S-phenyl-2-p-tolyl-ethyl)-oxalamide	

Table 1

Entry	Name	Structure
158	N-[2-(4-Chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
159	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamic acid	
160	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(3-fluoro-phenyl)-ethyl]-oxalamide	
161	N-[2-(2-Chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
162	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(3-methoxy-phenyl)-ethyl]-oxalamide	
163	N-(1,2-Diphenyl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
164	N-[2-(2,4-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
165	N-[2-(3,4-Dimethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
166	N-[2-(4-Ethyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
167	N-[2-(4-Ethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
168	N-[2-(4-Ethoxy-3-methoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
169	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-phenoxy-phenyl)-ethyl]-oxalamide	
170	N-[2-(3-Ethoxy-4-methoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
171	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-pyridin-2-yl-ethyl)-oxalamide	
172	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-pyridin-4-yl-ethyl)-oxalamide	

Table 1





Entry	Name	Structure
173	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-fluoro-phenyl)-ethyl]-oxalamide	
174	N-[2-(2-Bromo-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
175	N-[2-(2-Chloro-6-fluoro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
176	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2 <i>R</i> -phenyl-propyl)-oxalamide	

Table 1

Entry	Name	Structure
177	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-indan-1-yl-oxalamide	
178	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-isobutyl-oxalamide	
179	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methyl-butyl)-oxalamide	
180	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2R-phenyl-propyl)-oxalamide	

Table 1

Entry	Name	Structure
181	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-phenyl-propyl)-oxalamide	
182	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-indan-2-yl-oxalamide	
183	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R-phenyl-ethyl)-oxalamide	
184	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1S-phenyl-ethyl)-oxalamide	

Table 1

Entry	Name	Structure
185	N-[2-(3-Bromo-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
186	N-[2-(2,6-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
187	N-[2-(2,4-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
188	N-(2-Benzo[1,3]dioxol-5-yl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
189	N-[2-(3-Bromo-4-methoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
190	N-[2-(3,5-Dimethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
191	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-o-tolyl-ethyl)-oxalamide	

Table 1





Entry	Name	Structure
192	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-m-tolyl-ethyl)-oxalamide	
193	N-[2-(3-Ethoxy-phenyl)-ethyl]-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	
194	N-[2-(3,4-Dimethyl-phenyl)-ethyl]-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	
195	N-[2-(2,5-Dimethyl-phenyl)-ethyl]-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	

Table 1

Entry	Name	Structure
196	N-[2-(3-Chloro-4-propoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
197	N-[2-(4-Butoxy-3-chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
198	N-[2-(4-tert-Butyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1




Entry	Name	Structure
199	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-sulfamoyl-phenyl)-ethyl]-oxalamide	
200	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-hydroxy-3-methoxy-phenyl)-ethyl]-oxalamide	
201	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(3-hydroxy-4-methoxy-phenyl)-ethyl]-oxalamide	

Table 1




Entry	Name	Structure
202	N-(2,4-Dichloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
203	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-fluoro-2-trifluoromethyl-benzyl)-oxalamide	
204	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1-p-tolyl-ethyl)-oxalamide	

Table 1

Entry	Name	Structure
205	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-fluoro-4-trifluoromethyl-benzyl)-oxalamide	
206	N-(3-Chloro-4-fluoro-benzyl)-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	
207	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[1-(3-methoxy-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
208	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1-naphthalen-2-yl-ethyl)-oxalamide	
209	N-(4-Chloro-3-trifluoromethyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
210	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1-p-tolyl-ethyl)-oxalamide	

Table 1




Entry	Name	Structure
211	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(6-trifluoromethyl-pyridin-3-ylmethyl)-oxalamide	
212	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-methyl-benzyl)-oxalamide	
213	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methyl-benzyl)-oxalamide	

Table 1

Entry	Name	Structure
214	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-fluoro-3-trifluoromethyl-benzyl)-oxalamide	
215	N-(3,5-Dichloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
216	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R,2,3,4-tetrahydronaphthalen-1-yl)-oxalamide	
217	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1S,2,3,4-tetrahydronaphthalen-1-yl)-oxalamide	

Table 1

Entry	Name	Structure
218	N-Cyclopentyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
219	N-[1-(4-Bromo-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
220	N-(2-Fluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
221	N-[2-(3,4-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
222	N-(4-Fluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
223	N-(2,3-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
224	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-phenoxy-ethyl)-oxalamide	
225	N-(2,2-Diphenyl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
226	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-methoxy-phenyl)-ethyl]-oxalamide	
227	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-phenyl-propyl)-oxalamide	
228	N-[2-(4-Bromo-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
229	N-{4-[7-(1-Ethyl-piperidin-4-ylmethoxy)-6-methoxy-quinolin-4-yloxy]-3-fluoro-phenyl}-2-oxo-2-(2-phenyl-morpholin-4-yl)-acetamide	

Table 1

Entry	Name	Structure
230	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-fluoro-5-trifluoromethyl-benzyl)-oxalamide	
231	N-(3,5-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
232	N-(2-Chloro-5-trifluoromethyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
233	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-3-fluorophenyl]-N'-(2-dimethylamino-2-phenylethyl)-oxalamide	

Table 1

Entry	Name	Structure
234	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-methoxy-benzyl)-oxalamide	
235	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-trifluoromethyl-benzyl)-oxalamide	
236	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methoxy-benzyl)-oxalamide	

Table 1





Entry	Name	Structure
237	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-trifluoromethyl-benzyl)-oxalamide	
238	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-trifluoromethoxy-benzyl)-oxalamide	
239	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-methoxy-benzyl)-oxalamide	
240	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-trifluoromethyl-benzyl)-oxalamide	

Table 1

Entry	Name	Structure
241	N-(3-Chloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
242	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-trifluoromethoxy-benzyl)-oxalamide	
243	N-(2-Chloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
244	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-trifluoromethoxy-benzyl)-oxalamide	
245	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-methoxy-benzyl)-oxalamide	
246	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-trifluoromethyl-benzyl)-oxalamide	

Table 1

Entry	Name	Structure
247	N-{4-[7-(Azetidin-3-ylmethoxy)-6-methoxy-quinolin-4-yloxy]-3-fluorophenyl}-N'-phenethyl-oxalamide	
248	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-azetidin-3-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
249	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-hydroxy-2-phenyl-ethyl)-oxalamide	
250	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(2,4-difluorophenyl)-malonamide	

Table 1




Entry	Name	Structure
251	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(4-fluorophenyl)-N'-methyl-malonamide	
252	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R-phenyl-propyl)-oxalamide	
253	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R-phenyl-propyl)-oxalamide	

Table 1





Entry	Name	Structure
254	N-(3,4-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
255	N-(2,6-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
256	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-fluoro-phenyl)-ethyl]-oxalamide	
257	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenyl-oxalamide	

Table 1

Entry	Name	Structure
258	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-fluoro-phenyl)-oxalamide	
259	N-(4-Chloro-3-fluorophenyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
260	N-(3,4-Dimethoxy-phenyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
261	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methyl-butyl)-oxalamide	

Table 1

Entry	Name	Structure
262	N-(3,3-Dimethyl-butyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
263	N-{5-Chloro-6-[6-methoxy-7-(3-piperidin-1-yl-propoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	
264	N-{5-Chloro-6-[6-methoxy-7-(3-morpholin-4-yl-propoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	

Table 1

Entry	Name	Structure
265	N-{5-Chloro-6-[7-(3-diethylamino-propoxy)-6-methoxy-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluorophenyl)-malonamide	
266	N-(4-Chloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
267	N-(3,5-Dimethoxy-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1




Entry	Name	Structure
268	N-(4-Butyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
269	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-p-tolyl-ethyl)-oxalamide	
270	N-(3,5-Bis-trifluoromethyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
271	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-pyrazin-2-ylmethyl-oxalamide	
272	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-pyridin-2-ylmethyl-oxalamide	
273	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
274	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	

Table 1

Entry	Name	Structure
275	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-fluoro-3-trifluoromethyl-benzyl)-oxalamide	
276	N-[2-(2-Bromo-6-methoxy-phenyl)-ethyl]-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	
277	N-[2-(3,4-Dimethoxy-phenyl)-ethyl]-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-N-methyl-oxalamide	
278	N-[2-(5-Bromo-2-methoxy-phenyl)-ethyl]-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	

Table 1

Entry	Name	Structure
279	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-fluoro-5-trifluoromethylbenzyl)-oxalamide	
280	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[1-(4-fluorophenyl)-ethyl]-oxalamide	
281	N-(1S-Benzyl-2-oxo-2-pyrrolidin-1-yl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
282	N-{3-Fluoro-4-[6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	

Table 1




Entry	Name	Structure
283	N-[2-(4-Amino-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
284	2-(4-Benzyl-piperidin-1-yl)-N-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-acetamide	
285	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-phenyl]-N'-(4-fluoro-phenyl)-malonamide	

Table 1





Entry	Name	Structure
286	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(3-fluorophenyl)-malonamide	
287	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-phenyl-malonamide	
288	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(4-fluorophenyl)-2,2-dimethyl-malonamide	
289	N-Ethyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1


Entry	Name	Structure
290	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-isopropyl-oxalamide	
291	N-Butyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
292	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-methoxy-ethyl)-oxalamide	
293	N-Cyclopropylmethyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
294	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-morpholin-4-yl-ethyl)-oxalamide	

Table 1


Entry	Name	Structure
295	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-2-pyrrolidin-1-yl-acetamide	
296	N-Ethyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N-methyl-oxalamide	

[0078] In another aspect, the invention comprises a compound for modulating kinase activity of formula A-B-C, or a pharmaceutically acceptable salt, hydrate, or prodrug thereof, wherein, A is selected from:


$-R^3$		




B is selected from:



and, C is selected from:





wherein R^2 is selected from -H, halogen, trihalomethyl, -CN, -NH₂, -NO₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl;

q is 0 to 2;

each R^3 is independently selected from -H, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted arylalkyl, and optionally substituted heteroarylalkyl;

two R^3 , together with the nitrogen to which they are attached, form a four- to seven-membered heteroalicyclic, said four- to seven-membered heteroalicyclic optionally containing one additional heteroatom; when one said additional heteroatom is a nitrogen, then said nitrogen is optionally substituted with a group selected from -H, trihalomethyl, -SO₂R⁵, -SO₂NR⁵R⁵, -CO₂R⁵, -C(O)NR⁵R⁵, -C(O)R⁵, and optionally substituted lower alkyl;

each R^{35} is independently selected from -H, -C(=O)R³, -C(=O)OR³, -C(=O)SR³, -SO₂R³, -C(=O)N(R³)R³, and optionally substituted lower alkyl;

two R^{35} , together with the nitrogen to which they are attached, can combine to form a heteroalicyclic optionally substituted with between one and four of R^{60} , said heteroalicyclic may have an additional annular heteroatom, and said heteroalicyclic may have an aryl fused thereto, said aryl optionally substituted with an additional one to four of R^{60} ;

A^1 is selected from $=N-$, $=C(H)-$, and $=C(CN)-$;

A^2 is either $=N-$ or $=C(H)-$;

R^5 is $-H$ or optionally substituted lower alkyl;

R^8 is selected from R^3 , $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-SO_2R^3$, and $-C(O)R^3$;

R^9 , R^{10} , and R^{11} are each independently selected from $-H$, and $-OR^{12}$; or

R^9 is selected from $-H$, and $-OR^{12}$, and R^{10} and R^{11} , when taken together, are either an optionally substituted alkylidene or an oxo; and

R^{12} is selected from $-H$, $-C(O)R^3$, optionally substituted lower alkylidyne, optionally substituted lower arylalkylidyne, optionally substituted lower heterocyclylalkylidyne, optionally substituted lower alkylidene, optionally substituted lower alkylidenearyl, optionally substituted lower alkylideneheterocyclyl, optionally substituted lower alkyl, optionally substituted lower alkylaryl, optionally substituted aryl, optionally substituted lower heterocyclylalkyl, and optionally substituted heterocyclyl;

or two R^{12} 's, when taken together, form 1) a corresponding spirocyclic ketal when said two R^{12} 's stem from R^{10} and R^{11} , or 2) a corresponding cyclic ketal when said two R^{12} 's stem from R^9 and one of R^{10} and R^{11} ;

E^1 is selected from $-O-$, $-CH_2-$, $-N(R^5)-$, and $-S(O)_{0-2}-$;

Q is a five- to ten-membered ring system, optionally substituted with between zero and four of R^{20} ;

R^{20} is selected from $-H$, halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted lower alkyl;

R^{60} is selected from $-H$, halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$,



$-N(R^3)CO_2R^3$, $-C(O)R^3$, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted heteroarylalkyl, and optionally substituted arylalkyl;

two of R^{60} , when attached to a non-aromatic carbon, can be oxo;


each methylene in any of the above formulae is independently optionally substituted with R^{25} ;


each R^{25} is independently selected from halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic, two of R^{25} on a single carbon can be oxo;

with the proviso that when B is selected from:




and C contains $(R^2)_q$, and the remaining portion of C contains one of:






directly attached to $(R^2)_q$, then A must be one of:




and with the proviso that when C contains $(R^2)_q$, and B is selected from:

then the portion of C directly attached to





cannot contain



, when R^{70} is selected from -H, C_{1-4} alkyl, and C_{1-4} alkoxyl.


[0079] In another example the compound is according to paragraph [0078], wherein Q is selected from phenyl, naphthyl, 1,2,3,4-tetrahydronaphthyl, indanyl, benzodioxanyl, benzofuranyl, phenazinyl, phenothiazinyl, phenoxazinyl, tetrahydroisoquinolyl, pyrrolyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazoliny, imidazolidinyl, tetrahydropyridinyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazoliny, oxazolidinyl, triazolyl, isoxazolyl, isoxazolidinyl, thiazolyl, thiazoliny, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, isoindolyl, indoliny, isoindoliny, octahydroindolyl, octahydroisoindolyl, quinolyl, isoquinolyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, thienyl, benzothieliyl, and oxadiazolyl; each optionally substituted with between one and four of R^{20} ; wherein each R^{20} is independently selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl.

[0080] In another example the compound is according to paragraph [0079], wherein B is either of the following:





wherein A^1 is either =N- or =C(H)-.

[0081] In another example the compound is according to paragraph [0080], wherein B is



[0082] In another example the compound is according to paragraph [0081], wherein C is selected from:






wherein R^2 , R^3 , R^5 , R^{20} , R^{25} and R^{60} are as defined above.

[0083] In another example the compound is according to paragraph [0082], R² is selected from halogen, trihalomethyl, -CN, -NO₂, -OR³, -NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl

[0084] In another example the compound is according to paragraph [0083], wherein R² is halogen.

[0085] In another example the compound is according to paragraph [0084], wherein R² is either fluorine or chlorine.

[0086] In another aspect, the invention comprises a compound for modulating kinase activity according to Formula XI,



XII

or a pharmaceutically acceptable salt, hydrate, or prodrug thereof, wherein,

each R^1 is independently selected from halogen, $-OR^3$, $-NO_2$, $-NH_2$, $-NR^3R^4$, $-D-R^{50}$ and optionally substituted C_{1-6} alkyl;

R^{70} is selected from $-H$, halogen, $-OR^3$, $-S(O)_{0-2}R^3$, $-NO_2$, $-NH_2$, $-NR^3R^4$, and optionally substituted C_{1-6} alkyl;

Q is selected from $=N-$, $=C(H)-$, and $=C(CN)-$;

Z is selected from $-S(O)_{0-2}-$, $-O-$, and $-NR^5-$;

Ar is either a five- or six-membered arylene or a five- or six-membered heteroarylene containing between one and three heteroatoms;

G is either an optionally substituted cycloalkyl or an optionally substituted heteroalicyclic;

each R^2 is independently selected from halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted C_{1-6} alkyl;

each R^3 is independently $-H$ or R^4 ;


each R^4 is independently selected from optionally substituted C_{1-6} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C_{1-6} alkyl; or

R^3 and R^4 , when taken together with a common nitrogen to which they are attached, form an optionally substituted five- to seven-membered heterocyclyl, said optionally substituted five- to seven-membered heterocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

R^5 is $-H$ or optionally substituted C_{1-6} alkyl;

each D is independently selected from $-O-$, $-S(O)_{0-2}-$, and $-NR^5-$;

each R^{50} is independently either R^3 , or according to formula **XII**;



XII

wherein X^1 , X^2 , and optionally X^3 , represent the atoms of a saturated bridged ring system, said saturated bridged ring system comprising up to four annular heteroatoms represented by any of X^1 , X^2 , and X^3 ; wherein,

each X^1 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

each X^2 is independently an optionally substituted bridgehead methine or a bridgehead nitrogen;

each X^3 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

Y is either:

an optionally substituted lower alkylene linker, between D and either 1) any annular atom of the saturated bridged ring system, except X^2 when X^2 is a bridgehead nitrogen, or 2) any heteroatom, represented by any of R^6 or R^7 ; provided there are at least two carbon atoms between D and any annular heteroatom of the saturated bridged ring system or any heteroatom represented by any of R^6 or R^7 ;

or Y is absent, when Y is absent, said saturated bridged ring system, is directly attached to D via an annular carbon of said saturated bridged ring system, unless D is $-SO_2-$, in which case said saturated bridged ring system, is directly attached to D via an any annular atom of said saturated bridged ring system;

m and p are each independently one to four;

n is zero to two, when n equals zero, then there is a single bond between the two bridgehead X^2 's;

R^6 and R^7 are each independently selected from $-H$, halogen, trihalomethyl, $-CN$, $-NH_2$, $-NO_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^4$, $-SO_2NR^3R^4$, $-CO_2R^3$, $-C(O)NR^3R^4$, $-N(R^3)SO_2R^4$, $-N(R^3)C(O)R^3$, $-NCO_2R^3$, $-C(O)R^3$, optionally substituted C_{1-6} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, optionally substituted heterocyclyl, optionally substituted heterocyclyl C_{1-6} alkyl, and a bond to either Y or D; or

R^6 and R^7 , when taken together are oxo; or

R^6 and R^7 , when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

R^8 is selected from $-R^3$, Y, $-SO_2NR^3R^4$, $-CO_2R^4$, $-C(O)NR^3R^3$, $-SO_2R^4$, and $-C(O)R^3$; and


each R^{30} is independently selected from halogen, trihalomethyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OR}^3$, $-\text{NR}^3\text{R}^4$, $-\text{S}(\text{O})_{0-2}\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{N}(\text{R}^3)\text{SO}_2\text{R}^3$, $-\text{N}(\text{R}^3)\text{C}(\text{O})\text{R}^3$, $-\text{N}(\text{R}^3)\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{R}^3$, and optionally substituted C_{1-6} alkyl.

[0087] In one example, the compound is according to paragraph [0086], wherein Z is either $-\text{O}-$ or $-\text{NR}^5-$.

[0088] In another example, the compound is according to paragraph [0087], wherein at least one of R^1 is $-\text{D}-\text{R}^{50}$.

[0089] In another example, the compound is according to paragraph [0088], wherein D is $-\text{O}-$ and at least one other R^1 is $-\text{OR}^3$.

[0090] In another example, the compound is according to paragraph [0089], of formula **XIIIa** or **XIIIb**:



wherein Q^1 is either $=\text{N}-$ or $=\text{C}(\text{H})-$.

[0091] In another example, the compound is according to paragraph [0090], wherein R^{50} is selected from C_{1-6} alkyl optionally substituted with at least one of optionally substituted amino, optionally substituted C_{1-6} alkyl amino, optionally substituted C_{1-6} dialkyl amino, optionally substituted heteroalicyclic, and a group of formula **XII**.

[0092] In another example, the compound is according to paragraph [0091], wherein R^{3a} is C_{1-6} alkyl.


[0093] In another example, the compound is according to paragraph [0092], wherein Z is $-\text{O}-$.

[0094] In another example, the compound is according to paragraph [0093], wherein G is selected from cyclopropyl, aziradine, cyclobutyl, and azetidine, each optionally substituted with between zero and four of R^{30} .

[0095] In another example, the compound is according to paragraph [0094], wherein Q is either =N- or =C(H)-.

[0096] In another example, the compound is according to paragraph [0095], wherein R^2 is selected from -H, halogen, C_{1-6} alkyl and perfluoro C_{1-6} alkyl.

[0097] In another example, the compound is according to paragraph [0096], wherein $-N(R^{3b})R^4$ is selected from the following:



wherein J, is a five- to ten-membered ring, optionally substituted with between zero and five of R^{20} ;

each R^{20} is independently selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted C_{1-6} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C_{1-6} alkyl;

two of R^{20} , together with the atom or atoms to which they are attached, combine to form an optionally substituted three- to seven-membered heteroalicyclic, said optionally substituted three- to seven-membered heteroalicyclic either spiro- to J or fused to J;

E is selected from -O-, -N(R⁵)-, -CH₂-, and -S(O)₀₋₂-;

each R^{60} is independently selected from halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted C_{1-6} alkyl, optionally substituted aryl, optionally substituted heteroaryl C_{1-6} alkyl, and optionally substituted aryl C_{1-6} alkyl;

each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R^{25} ; and


R^{25} is selected from halogen, trihalomethyl, oxo, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OR}^3$, $-\text{NR}^3\text{R}^4$, $-\text{S}(\text{O})_{0-2}\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{N}(\text{R}^3)\text{SO}_2\text{R}^3$, $-\text{N}(\text{R}^3)\text{C}(\text{O})\text{R}^3$, $-\text{N}(\text{R}^3)\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{R}^3$, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, heteroaryl C_{1-6} alkyl, and optionally substituted C_{1-6} alkyl; or

two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic;

R^{3b} is equivalent to R^3 as defined above; and

R^4 and R^5 are as defined above.

[0098] In another example, the compound is according to paragraph [0097], of formula **XIVa** or **XIVb**:



[0099] In another example, the compound is according to paragraph [0098], wherein R^{50} is C_{1-6} alkyl optionally substituted with a group selected from optionally substituted amino, an optionally substituted alkylamino, optionally substituted dialkylamino, and optionally substituted heteroalicyclic.

[0100] In another example, the compound is according to paragraph [0099], wherein the heteroalicyclic portion of said optionally substituted heteroalicyclic of R^{50} is selected from the group consisting of piperidine, piperazine, morpholine, thiomorpholine, thiomorpholine 1-oxide, thiomorpholine 1,1-dioxide, 2-oxo-morpholine, pyrrolidine, and azepine.

[0101] In another example, the compound is according to paragraph [0099], wherein R^{50} is according to formula **XII**.

[0102] In another example, the compound is according to paragraph [0101], wherein the saturated bridged ring system according to formula **XII** has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], [3.1.0], [3.3.3], [3.3.2], [3.3.1], [3.2.2], [3.2.1], [2.2.2], and [2.2.1].


[0103] In another example, the compound is according to paragraph [0102], wherein Y is selected from $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2-$, and absent.

[0104] In another example, the compound is according to paragraph [0103], wherein n is 0 and the saturated bridged ring system according to formula **XII** has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], and [3.1.0].

[0105] In another example, the compound is according to paragraph [0104], wherein said saturated bridged ring system contains at least one annular nitrogen or at least one annular oxygen.

[0106] In another example, the compound is according to paragraph [0105], wherein said saturated bridged ring system contains $-\text{NR}^8-$, wherein R^8 is selected from $-\text{H}$, optionally substituted $\text{C}_{1-6}\text{alkyl}$, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{SO}_2\text{R}^3$, and $-\text{C}(\text{O})\text{R}^3$.

[0107] In another example, the compound is according to paragraph [0105], wherein said saturated bridged ring system is of formula **XV**,



XV

wherein U^1 is selected from $-\text{O}-$, $-\text{S}(\text{O})_{0-2}-$, $-\text{NR}^8-$, $-\text{CR}^6\text{R}^7-$, and absent; and e is 0 or 1.

[0108] In another example, the compound is according to paragraph [0107], wherein Y is $-\text{CH}_2-$.


[0109] In another example, the compound is according to paragraph [0108], wherein U^1 is $-\text{NR}^8-$, wherein R^8 is selected from $-\text{H}$, optionally substituted lower alkyl, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{SO}_2\text{R}^3$, and $-\text{C}(\text{O})\text{R}^3$.

[0110] In another example, the compound is according to paragraph [0108], wherein U^1 is $-\text{O}-$.

[0111] In another example, the compound is according to paragraph [0108], wherein U^1 is absent.

[0112] In another example, the compound is according to paragraph [0103], wherein Y is selected from $-CH_2CH_2-$, $-CH_2-$, and absent.

[0113] In another example, the compound is according to paragraph [0112], wherein said saturated bridged ring system is of formula XVI,



XVI

wherein R^9 , R^{10} , and R^{11} are each independently selected from $-H$, and $-OR^{12}$; or

R^9 is selected from $-H$, and $-OR^{12}$, and R^{10} and R^{11} , when taken together, are either an optionally substituted alkylidene or an oxo;

R^{12} is selected from $-H$, $-C(O)R^3$, optionally substituted lower alkylidene, optionally substituted lower arylalkylidene, optionally substituted lower heterocyclylalkylidene, optionally substituted lower alkylidene, optionally substituted lower alkylidenearyl, optionally substituted lower alkylideneheterocyclyl, optionally substituted lower alkyl, optionally substituted lower alkylaryl, optionally substituted aryl, optionally substituted lower heterocyclylalkyl, and optionally substituted heterocyclyl;


or two R^{12} 's, when taken together, form 1) a corresponding spirocyclic ketal when said two R^{12} 's stem from R^{10} and R^{11} , or 2) a corresponding cyclic ketal when said two R^{12} 's stem from R^9 and one of R^{10} and R^{11} .

[0114] In another example, the compound is according to paragraph [0113], wherein one of R^{10} and R^{11} is $-OR^{12}$, wherein R^{12} is selected from $-H$, $-C(O)R^3$, and optionally substituted lower alkyl; and R^9 and the other of R^{10} and R^{11} are both $-H$.

[0115] In another example, the compound is according to paragraph [0114], wherein Y is either $-CH_2-$ or absent.

[0116] In another example, the compound is according to paragraph [0113], wherein R^9 is an alkyl group containing at least one fluorine substitution thereon.

[0117] In another example, the compound is according to paragraph [0106], wherein said saturated bridged ring system is of formula **XVII**.

**XVII**

[0118] In another example, the compound is according to paragraph [0117], wherein Y is either -CH₂- or absent.

[0119] In another example, the compound is according to paragraph [0118], wherein R⁸ is methyl or ethyl.

[0120] In another example, the compound is according to paragraph [0119], wherein at least one of R² is halogen.


[0121] In another example, the compound is according to paragraph [0106], wherein said saturated bridged ring system is of formula **XVIII**.

**XVIII**

[0122] In another example, the compound is according to paragraph [0121], wherein Y is -CH₂-.

[0123] In another example, the compound is according to paragraph [0122], wherein R⁸ is methyl or ethyl.

[0124] In another example, the compound is according to paragraph [0105], wherein said saturated bridged ring system is of formula **XIX**

**XIX**

wherein U² is selected from -O-, -S(O)₀₋₂-, -NR⁸-, -CR⁶R⁷-, and absent.

[0125] In another example, the compound is according to paragraph [0124], wherein R³ of formula **XIX** is selected from -H and optionally substituted alkyl.

[0126] In another example, the compound is according to paragraph [0125], wherein U^2 is either $-CR^6R^7-$ or absent.

[0127] In another example, the compound is according to paragraph [0126], wherein U^2 is either $-CH_2-$ or absent.

[0128] In another example, the compound is according to paragraph [0127], wherein Y is $-CH_2-$.

[0129] In another example, the compound is according to paragraph [0106], wherein said saturated bridged ring system is according to formula **XX**.



XX

[0130] In another example, the compound is according to paragraph [0129], wherein R^8 is methyl or ethyl.

[0131] In another example, the compound is according to any of paragraphs [0099] – [0130], wherein R^2 is selected from C_{1-6} alkyl, perfluoro C_{1-6} alkyl, and halogen.

[0132] In another example, the compound is according to paragraph [0131], wherein R^2 is selected from perfluoro C_{1-3} alkyl and halogen.

[0133] In another example, the compound is according to any of paragraphs [0099] – [0130], wherein R^{20} is selected from halogen, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C_{1-6} alkyl, and (two of R^{20}) together with the atom or atoms to which they are attached, an optionally substituted three- to six-membered heteroalicyclic, said optionally substituted three- to six-membered heteroalicyclic fused to the phenyl as in **XIVa** or **XIVb**.

[0134] In another example, the compound is according to paragraph [0133], wherein R^{20} is selected from halogen, $-NR^3R^4$, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C_{1-6} alkyl, and (two of R^{20}) together with the atom or atoms to which they are attached, an optionally substituted five- to six-membered heteroalicyclic, said optionally substituted five- to six-membered heteroalicyclic fused to the phenyl as in **XIVa** or **XIVb**.

[0135] In another example, the compound is according to paragraph [0134], wherein R² is selected from C₁₋₆ alkyl, perfluoro C₁₋₆ alkyl, and halogen.

[0136] In another example, the compound is according to paragraph [0135], wherein R² is selected from perfluoro C₁₋₃ alkyl and halogen.

[0137] In another example, the compound is according to paragraph [0086], selected from Table 2.

Table 2

Entry	Name	Structure
1	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
2	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
3	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(phenylmethyl)cyclopropane-1,1-dicarboxamide	
4	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-phenylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
5	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
6	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperidin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
7	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperidin-1-ylbutyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
8	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-phenylethyl)cyclopropane-1,1-dicarboxamide	
9	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-methylpyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
10	N-{4-[(7-chloroquinolin-4-yl)oxy]-3-fluorophenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
11	N-{4-[(7-chloroquinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
12	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
13	N-(4-{[6,7-bis(methyloxy)quinazolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
14	N-(4-{[6,7-bis(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
15	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
16	N-{5-chloro-6-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy}pyridin-3-yl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
17	N-[5-chloro-6-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
18	N-[5-chloro-6-({6-(methyloxy)-7-[(phenylmethyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
19	N-(4-{[7-{{2-(diethylamino)ethyl}oxy}-6-(methyloxy)quinolin-4-yl}oxy]-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
20	N-(4-{[7-{{2-(diethylamino)ethyl}oxy}-6-(methyloxy)quinolin-4-yl}oxy]-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
21	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy]quinazolin-4-yl}oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
22	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
23	N-(4-fluorophenyl)-N'-[2-methyl-6-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]cyclopropane-1,1-dicarboxamide	
24	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
25	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloro-2-methylpyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
26	N-[3-fluoro-4-({7-(methyloxy)-6-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
27	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3,5-difluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
28	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-2,5-difluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
29	N-[3-fluoro-4-({7-(methyloxy)-6-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
30	N-[3-fluoro-4-[(6-(methyloxy)-7-(2-methyloctahydrocyclopenta[c]pyrrol-5-ylmethoxy)quinazolin-4-yl]oxy]phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
31	N-[3-fluoro-4-[(7-(methyloxy)-6-[(1-methylpiperidin-4-yl)methyl]oxy]quinazolin-4-yl]oxy]phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
32	N-[5-fluoro-2-methyl-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
33	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2,3,5-trifluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
34	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-2-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
35	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-chloro-5-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
36	N-(3-fluoro-4-{[6-hydroxy-7-(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
37	N-(4-fluorophenyl)-N'-[2-methyl-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]cyclopropane-1,1-dicarboxamide	
38	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
39	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
40	N-{3-fluoro-4-[(6-(methyloxy)-7-{[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
41	N-(4-fluorophenyl)-N'-[4-[(6-(methyloxy)-7-{[3-(morpholin-4-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl]cyclopropane-1,1-dicarboxamide	
42	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
43	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-2-chloro-5-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
44	N-(4-{{6,7-bis(methyloxy)-2-(methylthio)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
45	N-(4-fluorophenyl)-N'-(4-{{2-methyl-6,7-bis(methyloxy)quinazolin-4-yl}oxy}phenyl)cyclopropane-1,1-dicarboxamide	
46	N-(4-{{2-amino-6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
47	N-(3-fluoro-4-{{2-(methylamino)-6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
48	(1S,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
49	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
50	N-(4-{[6-{[3-(diethylamino)propyl]oxy}-7-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
51	N-(4-{[6-{[2-(diethylamino)ethyl]oxy}-7-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
52	1,1-dimethylethyl 4-(3- {[4-[(2-fluoro-4- {[(1- {[(4- fluorophenyl)amino]carbo nyl}cyclopropyl)carbonyl]amino}phenyl)oxy]-6- (methyloxy)quinolin-7- yl]oxy}propyl)piperazine- 1-carboxylate	
53	(1R,2R)-N-[3-fluoro-4- (6-(methyloxy)-7-[(3- morpholin-4- ylpropyl)oxy]quinazolin- 4-yl]oxy)phenyl]-N'-(4- fluorophenyl)-2- methylcyclopropane-1,1- dicarboxamide	
54	(1R,2R)-N-(4- {[7- {[2- (diethylamino)ethyl]oxy}- 6-(methyloxy)quinazolin- 4-yl]oxy}-3- fluorophenyl)-N'-(4- fluorophenyl)-2- methylcyclopropane-1,1- dicarboxamide	
55	N-(4- {[7- {[3- (diethylamino)propyl]oxy }-6- (methyloxy)quinazolin-4- yl]oxy}-3-fluorophenyl)- N'-(4- fluorophenyl)cyclopropan e-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
56	N-(4-{[7-{[3-(4-acetylpiperazin-1-yl)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
57	1,1-dimethylethyl 4-(3-{[4-[(2-fluoro-4-{[[(1R,2R)-1-{[(4-fluorophenyl)amino]carbonyl}-2-methylcyclopropyl)carbonyl]amino}phenyl]oxy]-6-(methyloxy)quinolin-7-yl]oxy}propyl)piperazine-1-carboxylate	
58	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)-1-(phenylmethyl)azetidine-3,3-dicarboxamide	

Table 2

Entry	Name	Structure
59	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)azetidine-3,3-dicarboxamide	
60	(1R,2S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
61	(1R,2R)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
62	(1R,2R)-N-[3-fluoro-4-((6-(methyloxy)-7-[(3-piperazin-1-yl)propyl]oxy)quinolin-4-yl)oxy]phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
63	N-(3-fluoro-4-{{7-({3-[4-(1-methylethyl)piperazin-1-yl]propyl}oxy)-6-(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
64	N-(4-{{7-{{3-(diethylamino)propyl}oxy}-6-(methyloxy)quinazolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
65	(1R,2R)-N-(4-{{7-{{3-(diethylamino)propyl}oxy}-6-(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
66	(1R,2R)-N-(4-{{7-{{2-(diethylamino)ethyl}oxy}-6-(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
67	(1R,2S)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
68	(1R,2S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
69	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
70	(1R,2S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl]oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
71	(1R,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
72	(1R,2R,3S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
73	(1R,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
74	(1R,2R,3S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	

Table 2





Entry	Name	Structure
75	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
76	(2R,3R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
77	(2R,3R)-N-(4-{[7-([3-(diethylamino)propyl)oxy]-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
78	N-(4-{[7-([3-(diethylamino)propyl)oxy]-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
79	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
80	(1R,2R,3S)-N-(4-{[7-{{3-(diethylamino)propyl}oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
81	N-(4-{[7-{{2-(diethylamino)ethyl}oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
82	(1R,2R,3S)-N-(4-{[7-{{2-(diethylamino)ethyl}oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
83	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
84	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
85	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
86	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
87	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl]oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
88	N-[3-fluoro-4-[(6-(methyloxy)-7-{[3-piperazin-1-yl]propyl]oxy}quinazolin-4-yl]oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
89	(2R,3R)-N-[3-fluoro-4-((6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl)oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
90	N-(4-{[7-[[3-(diethylamino)propyl]oxy]-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
91	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
92	(1R,2R)-N-(4-{[7-[[3-(diethylamino)propyl]oxy]-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
93	(1R,2R)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
94	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
95	(2R,3R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
96	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
97	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	

Table 2


Entry	Name	Structure
98	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[(4-fluorophenyl)methyl]cyclopropane-1,1-dicarboxamide	
99	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-morpholin-4-ylethyl]cyclopropane-1,1-dicarboxamide	
100	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	
101	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	
102	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	

Table 2



Entry	Name	Structure
103	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-[2-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	
104	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-phenylcyclopropane-1,1-dicarboxamide	
105	N-[3-(aminomethyl)phenyl]-N'-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)cyclopropane-1,1-dicarboxamide	
106	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-[3-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	
107	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-[3-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	

- [0176] Another aspect of the invention is a pharmaceutical composition comprising a compound according to any one of paragraphs [0033]-[0120] and a pharmaceutically acceptable carrier.
- [0177] Another aspect of the invention is a metabolite of the compound or the pharmaceutical composition according to any one of paragraphs [0022]-[0122].
- [0178] Another aspect of the invention is a method of modulating the *in vivo* activity of a kinase, the method comprising administering to a subject an effective amount of the compound or the pharmaceutical composition according to any of paragraphs [0033]-[0121].
- [0179] Another aspect of the invention is the method according to paragraph [0123], wherein modulating the *in vivo* activity of the kinase comprises inhibition of said kinase.
- [0180] Another aspect of the invention is the method according to paragraph [0124], wherein the kinase is at least one of c-Met, KDR, c-Kit, flt-3, and flt-4.
- [0181] Another aspect of the invention is the method according to paragraph [0125], wherein the kinase is c-Met.
- [0182] Another aspect of the invention is a method of treating diseases or disorders associated with uncontrolled, abnormal, and/or unwanted cellular activities, the method comprising administering, to a mammal in need thereof, a therapeutically effective amount of the compound or the pharmaceutical composition as described in any one of paragraphs [0033]-[0121].
- [0183] Another aspect of the invention is a method of screening for a modulator of a kinase, said kinase selected from c-Met, KDR, c-Kit, flt-3, and flt-4, the method comprising combining a compound according to any one of paragraphs [0033]-[0120], and at least one candidate agent and determining the effect of the candidate agent on the activity of said kinase.
- [0184] Another aspect of the invention is a method of inhibiting proliferative activity in a cell, the method comprising administering an effective amount of a composition comprising a compound according any one of paragraphs [0033]-[0120] to a cell or a plurality of cells.
- [0185] As mentioned, although improved quinolines and quinazolines of the invention can be made via conventional serial methods, due to their complex structure, more efficient

routes are desirable, particularly convergent syntheses. Thus, the present invention also comprises a process for preparing a compound of Formula **XXI**,



comprising reaction of a compound of Formula **XXII**, with a compound of Formula **XXIII**



wherein,

each R^1 is independently selected from halogen, $-OR^3$, $-NO_2$, $-NH_2$, $-NR^3R^3$, $-D-R^{50}$ and optionally substituted C_{1-6} alkyl;

R^{70} is selected from $-H$, halogen, $-OR^3$, $-S(O)_{0-2}R^3$, $-NO_2$, $-NH_2$, $-NR^3R^3$, and optionally substituted C_{1-6} alkyl;

J is selected from $=N-$, $=C(H)-$, $=C(halogen)-$, and $=C(CN)-$;

Z is selected from $-S(O)_{0-2}-$, $-O-$, and $-NR^5-$;

each R^5 is independently selected from $-H$, optionally substituted C_{1-6} alkyl, optionally substituted aryl, and optionally substituted aryl C_{1-6} alkyl;

Ar is either a five- to ten-membered arylene or a five- to ten-membered heteroarylene containing between one and three heteroatoms;

R^2 is selected from $-H$, halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted C_{1-6} alkyl;

each R^3 is independently selected from $-H$, $-Si(R^5)(R^5)R^5$, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted arylalkyl, and optionally substituted heteroarylalkyl;

two R^3 , together with the nitrogen to which they are attached, form a four- to seven-membered heteroalicyclic, said four- to seven-membered heteroalicyclic optionally containing one additional heteroatom; when one said additional heteroatom is a nitrogen, then said nitrogen is optionally substituted with a group selected from -H, trihalomethyl, $-SO_2R^5$, $-SO_2NR^5R^5$, $-CO_2R^5$, $-C(O)NR^5R^5$, $-C(O)R^5$, and optionally substituted lower alkyl;

B is selected from absent, $-N(R^{13})-$, $-N(SO_2R^{13})-$, $-O-$, $-S(O)_{0.2}-$, and $-C(=O)-$;

L is selected from absent, $-C(=S)N(R^{13})-$, $-C(=NR^{14})N(R^{13})-$, $-SO_2N(R^{13})-$, $-SO_2-$, $-C(=O)N(R^{13})-$, $-N(R^{13})-$, $-C(=O)C_{1.2}alkylN(R^{13})-$, $-N(R^{13})C_{1.2}alkylC(=O)-$, $-C(=O)C_{0.1}alkylC(=O)N(R^{13})-$, $-C(=O)-$, $-C_{0.4}alkylene-$, $-C(=O)C_{0.1}alkylC(=O)OR^3-$, $-C(=NR^{14})C_{0.1}alkylC(=O)-$, $-C(=O)C_{0.1}alkylC(=O)-$, and an optionally substituted four- to six-membered heterocyclyl containing between one and three annular heteroatoms and comprising at least one nitrogen;

T is selected from -H, $-R^{13}$, $-C_{0.4}alkyl$, $-C_{0.4}alkylQ$, $-OC_{0.4}alkylQ$, $-C_{0.4}alkylOQ$, $-N(R^{13})C_{0.4}alkylQ$, $-SO_2C_{0.4}alkylQ$, $-C(=O)C_{0.4}alkylQ$, $-C_{0.4}alkylN(R^{13})Q$, and $-C(=O)N(R^{13})C_{0.4}alkylQ$, wherein each of the aforementioned $C_{0.4}alkyl$ is optionally substituted;


Q is a five- to ten-membered ring system, optionally substituted with between zero and four of R^{20} ;

each R^{20} is independently selected from -H, halogen, trihalomethyl, -CN, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0.2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, optionally substituted $C_{1.6}alkyl$, optionally substituted aryl, optionally substituted aryl $C_{1.6}alkyl$, optionally substituted heterocyclyl, and optionally substituted heterocyclyl $C_{1.6}alkyl$;

two of R^{20} , together with the atom or atoms to which they are attached, combine to form an optionally substituted three- to seven-membered heteroalicyclic, said optionally substituted three- to seven-membered heteroalicyclic either spiro- to Q or fused to Q;

D is selected from $-O-$, $-S(O)_{0.2}-$, and $-NR^{15}-$;

R^{50} is either R^3 , or according to formula XXIV;



XXIV

wherein X^1 , X^2 , and optionally X^3 , represent the atoms of a saturated bridged ring system, said saturated bridged ring system comprising up to four annular heteroatoms represented by any of X^1 , X^2 , and X^3 ; wherein,

each X^1 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

each X^2 is independently an optionally substituted bridgehead methine or a bridgehead nitrogen;

each X^3 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

Y is either:

an optionally substituted C_{1-6} alkylene linker, between D and either 1) any annular atom of the saturated bridged ring system, except X^2 when X^2 is a bridgehead nitrogen, or 2) any heteroatom, represented by any of R^6 or R^7 ; provided there are at least two carbon atoms between D and any annular heteroatom of the saturated bridged ring system or any heteroatom represented by any of R^6 or R^7 ;

or Y is absent, when Y is absent, said saturated bridged ring system, is directly attached to D via an annular carbon of said saturated bridged ring system, unless D is $-SO_2-$, in which case said saturated bridged ring system, is directly attached to D via an any annular atom of said saturated bridged ring system;

m and p are each independently one to four;

n is zero to two, when n is zero, then there is a single bond between the two bridgehead X^2 's;

R^6 and R^7 are each independently selected from $-H$, halogen, trihalomethyl, $-CN$, $-NH_2$, $-NO_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-NCO_2R^3$, $-C(O)R^3$, optionally substituted C_{1-6} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, optionally substituted heterocyclyl, optionally substituted heterocyclyl a C_{1-6} lky; and a bond to either Y or D; or

R^6 and R^7 , when taken together are oxo; or

R^6 and R^7 , when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

R^8 is selected from $-R^3$, Y, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{SO}_2\text{R}^3$, and $-\text{C}(\text{O})\text{R}^3$;

R^{13} is selected from $-\text{H}$, $-\text{C}(=\text{O})\text{R}^3$, $-\text{C}(=\text{O})\text{OR}^3$, $-\text{C}(=\text{O})\text{SR}^3$, $-\text{SO}_2\text{R}^3$, $-\text{C}(=\text{O})\text{N}(\text{R}^3)\text{R}^3$, and optionally substituted $\text{C}_{1-6}\text{alkyl}$;

two R^{13} , together with the atom or atoms to which they are attached, can combine to form a heteroalicyclic optionally substituted with between one and four of R^{60} , said heteroalicyclic comprising up to four annular heteroatoms, and said heteroalicyclic optionally comprising an aryl or heteroaryl fused thereto, in which case said aryl or heteroaryl is optionally substituted with an additional one to four of R^{60} ;

R^{14} is selected from $-\text{H}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{N}(\text{R}^3)\text{R}^3$, $-\text{CN}$, $-\text{OR}^3$, optionally substituted $\text{C}_{1-6}\text{alkyl}$, optionally substituted heteroalicyclyl $\text{C}_{1-6}\text{alkyl}$, optionally substituted aryl, optionally substituted aryl $\text{C}_{1-6}\text{alkyl}$ and optionally substituted heteroalicyclic;

R^{15} is a group $-\text{M}^1-\text{M}^2$, wherein M^1 is selected from absent, $-\text{C}(=\text{S})\text{N}(\text{R}^{13})-$, $-\text{C}(=\text{NR}^{14})\text{N}(\text{R}^{13})-$, $-\text{SO}_2\text{N}(\text{R}^{13})-$, $-\text{SO}_2-$, $-\text{C}(=\text{O})\text{N}(\text{R}^{13})-$, $-\text{C}(=\text{O})\text{C}(=\text{O})\text{N}(\text{R}^{13})-$, $-\text{C}_{0-4}\text{alkylene}-$, $-\text{C}(=\text{O})-$, and an optionally substituted four to six-membered heterocyclyl containing between one and three heteroatoms but comprising at least one nitrogen; and M^2 is selected from $-\text{H}$, $-\text{C}_{0-6}\text{alkyl}$, alkoxy, $-\text{C}(=\text{O})\text{C}_{0-4}\text{alkylQ}$, $-\text{C}_{0-4}\text{alkylQ}$, $-\text{OC}_{0-4}\text{alkylQ}-$, $-\text{N}(\text{R}^{13})\text{C}_{0-4}\text{alkylQ}-$, and $-\text{C}(=\text{O})\text{N}(\text{R}^{13})\text{C}_{0-4}\text{alkylQ}$;

R^{60} is selected from $-\text{H}$, halogen, trihalomethyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OR}^3$, $-\text{NR}^3\text{R}^3$, $-\text{S}(\text{O})_{0-2}\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{N}(\text{R}^3)\text{SO}_2\text{R}^3$, $-\text{N}(\text{R}^3)\text{C}(\text{O})\text{R}^3$, $-\text{N}(\text{R}^3)\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{R}^3$, optionally substituted $\text{C}_{1-6}\text{alkyl}$, optionally substituted aryl, optionally substituted heteroaryl $\text{C}_{1-6}\text{alkyl}$, and optionally substituted aryl $\text{C}_{1-6}\text{alkyl}$;

two of R^{60} , when attached to a non-aromatic carbon, can be oxo;


P^1 is a suitable leaving group; and


P^2 is selected from $-\text{H}$, a metal, and a group removed *in-situ* when combining **XXII** and **XXIII** to make **XXI**.

[0131] In one example, the process is according to paragraph [0130], wherein Ar is *para*-phenylene as defined by the substitution pattern of -Z- and -B-L-T about said phenylene.

[0132] In another example, the process is according to paragraph [0131], wherein Z is either -O- or -NR⁵-.

[0133] In another example, the process is according to paragraph [0132], wherein -B-L-T is selected from the following:





wherein Q, R²⁰, and R¹³ are as defined above; each E is selected from -O-, -N(R¹³)-, -CH₂-, and -S(O)₀₋₂-; M is selected from -O-, -N(R¹³)-, -CH₂-, and -C(=O)N(R¹³)-; each V is independently either =N- or =C(H)-; each methylene in any of the above formulae is

independently optionally substituted with R^{25} ; and R^{25} is selected from halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, heteroaryl C_{1-6} alkyl, and optionally substituted C_{1-6} alkyl; two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form an optionally substituted three- to seven-membered alicyclic or heteroalicyclic; two of R^{25} on a single carbon can be oxo.

[0134] In another example, the process is according to paragraph [0133], wherein there is one of R^1 that is $-D-R^{50}$ and another of R^1 that is $-OR^{3a}$.


[0135] In another example, the process is according to paragraph [0134], wherein D is $-O-$.


[0136] In another example, the process is according to paragraph [0135], wherein $-O-R^{50}$ and $-OR^{3a}$ are interchangeably located at the 6-position and 7-position of the quinazoline or quinoline according to Formula XXI.

[0137] In another example, the process is according to paragraph [0136], wherein $-OR^{3a}$ is selected from $-OH$, $-OSi(R^5)(R^5)R^5$, and optionally substituted $-OC_{1-6}$ alkyl.

[0138] In another example, the process is according to paragraph [0137], wherein J is $=N-$ or $=C(H)-$.


[0139] In another example, the process is according to paragraph [0138], wherein $-B-L-T$ is selected from:






wherein Q, R²⁰, R¹³, E, and R⁶⁰ are as defined above; each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R²⁵; and R²⁵ is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted aryl C₁₋₆alkyl, heteroaryl C₁₋₆alkyl, and optionally substituted C₁₋₆alkyl; two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered optionally substituted alicyclic or heteroalicyclic.

[0140] In another example, the process is according to paragraph [0139], wherein Q is selected from the following three formula:




wherein R²⁰ is defined as above, and P is a five- to seven-membered ring, including the two shared carbons of the aromatic ring to which P is fused, P optionally containing between one and three heteroatoms.

[0141] In another example, the process is according to paragraph [0140], wherein -B-L-T is either of formula XXV or formula XXVI,



XXV




XXVI

wherein R²⁰ is defined as above; G is either an optionally substituted cycloalkyl or an optionally substituted heteroalicyclic; each R³⁰ is independently selected from halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally

substituted C₁₋₆alkyl; and R^{3a} and R^{3b} are each independently selected from -H and optionally substituted C₁₋₆alkyl.

[0142] In another example, the process is according to paragraph [0141], wherein a compound of formula **XXIIa** is combined with a compound of formula **XXIIIa** to make a compound of formula **XXIa**,



wherein -B-L-T, Z, J, R⁵⁰, and R² are as defined above; R⁷⁰ is selected from -H, -NO₂, -NH₂, and -NR³R³; provided when Z is -N(R⁵)- that R⁵ is selected from -H, C₁₋₃alkyl, and aryl C₁₋₃alkyl; P¹ is selected from halogen, optionally substituted alkyl-S(O)₀₋₂-, optionally substituted arylsulfonate, optionally substituted alkylsulfonate, a group containing boron, an azide, a group containing phosphorus, and a metal; and P² is selected from -H and a metal.

[0143] In another example, the process is according to paragraph [0142], wherein P² is selected from -H, lithium, sodium, potassium, cesium, copper, palladium, and titanium.

[0144] In another example, the process is according to paragraph [0143], wherein Z is -O-.

[0145] In another example, the process is according to paragraph [0144], wherein P¹ is selected from chlorine, bromine, a toluene sulfonate, and trifluoromethanesulfonate.

[0146] In another example, the process is according to paragraph [0145], wherein R⁷⁰ is -H.

[0147] In another example, the process is according to paragraph [0146], wherein J is =C(H)-.

[0148] In another example, the process is according to paragraph [0147], wherein R² is selected from C₁₋₆ alkyl, perfluoro C₁₋₆ alkyl, and halogen.

- [0149] In another example, the process is according to paragraph [0148], wherein **XXIIa** and **XXIIIa** are heated together, optionally with a base, optionally with microwave radiation, to form **XXIa**.
- [0150] In another example, the process is according to paragraph [0149], wherein the base is selected from an organic base, an inorganic base, and a combination of an organic base and an inorganic base.
- [0151] In another example, the process is according to paragraph [0150], wherein the base is selected from 2,6-lutidine, 4-N,N-dimethylaminopyridine, and a metal carbonate.
- [0152] In another example, the process is according to paragraph [0151], wherein **XXIIa** and **XXIIIa** are heated together in a solvent with said base, at between about 40°C and 200°C for between about one hour and twenty-four hours to form **XXIa**.
- [0153] In another example, the process is according to paragraph [0152], wherein the solvent is an organic solvent.
- [0154] In another example, the process is according to paragraph [0153], wherein one molar equivalent of **XXIIa** is combined with between about one quarter and four molar equivalents of **XXIIIa**.
- [0155] In another example, the process is according to paragraph [0154], wherein one molar equivalent of **XXIIa** is combined with more than one but less than two molar equivalents of **XXIIIa**.
- [0156] In another example, the process is according to paragraph [0155], wherein **XXIIa** is combined with **XXIIIa** and said base in an aromatic solvent to form a mixture, and said mixture is heated to between about 100°C and 200°C for between about one and ten hours to form **Ia**.
- [0157] In another example, the process is according to paragraph [0156], wherein the aromatic solvent is an optionally substituted benzene.
- [0158] In another example, the process is according to paragraph [0157], wherein the aromatic solvent is bromobenzene.
- [0159] In another example, the process is according to paragraph [0158], wherein the base is 4-N,N-dimethylaminopyridine.

- [0160] In another example, the process is according to paragraph [0159], wherein said mixture is heated to reflux for between about three and seven hours.
- [0161] In another example, the process is according to paragraph [0160], wherein said mixture is heated to reflux for between about four and six hours.
- [0162] In another example, the process is according to paragraph [0155], wherein **XXIIa** is combined with **XXIIIa** and said base in a non-aromatic solvent to form a mixture, and said mixture is heated to between about 40°C and 100°C for between about one and twenty hours to form **XXIa**.
- [0163] In another example, the process is according to paragraph [0162], wherein the non-aromatic solvent comprises a functional group selected from an amide, and ether, a nitrile, a halide, an ester, an amine, and a ketone.
- [0164] In another example, the process is according to paragraph [0163], wherein the non-aromatic solvent is N,N-dimethylacetamide.
- [0165] In another example, the process is according to paragraph [0164], wherein the base is potassium carbonate.
- [0166] In another example, the process is according to paragraph [0165], wherein said mixture is heated to about 50°C between about ten and twenty hours.
- [0167] In another example, the process is according to paragraph [0166], wherein the aromatic solvent is an optionally substituted pyridine.
- [0168] In another example, the process is according to paragraph [0167], wherein the aromatic solvent is 2,6-lutidine.
- [0169] In another example, the process is according to paragraph [0168], wherein the base is 2,6-lutidine.
- [0170] In another example, the process is according to paragraph [0169], wherein said mixture is heated to reflux for between about three and seven hours.
- [0171] In another example, the process is according to paragraph [0170], wherein said mixture is heated to reflux for between about four and six hours.
- [0172] In another example, the process is according to paragraph [0154], wherein one molar equivalent of **XXIIIa** is combined with more than one but less than two molar equivalents of **XXIIa**.

[0173] In another example, the process is according to paragraph [0172], wherein **XXIIa** is combined with **XXIIIa** and said base in an aromatic solvent to form a mixture, and said mixture is heated to between about 100°C and 200°C for between about ten and twenty hours to form **XXIa**.




[0174] In another example, the process is according to paragraph [0173], wherein the aromatic solvent is an optionally substituted pyridine.



[0175] In another example, the process is according to paragraph [0174], wherein the aromatic solvent is 2,6-lutidine.

[0176] In another example, the process is according to paragraph [0175], wherein the base is 2,6-lutidine.

[0177] In another example, the process is according to paragraph [0176], wherein said mixture is heated to between about 150°C and 200°C for between about fifteen and twenty hours.

[0178] In another example, the process is according to any of paragraphs [0149] – [0177], wherein a compound of formula **XXIIb** is substituted for the compound of formula **XXIIa**, and either a compound of formula **XXIIIb** or a compound of formula **XXIIIc** is substituted for the compound of formula **XXIIIa**, in order to make a compound of formula **XXIb** or a compound of formula **XXIc**, respectively,





wherein J, R⁵⁰, R²⁰ and R² are as defined above.

[0179] In another example, the process is according to paragraph [0178], wherein R², if present, is halogen.

[0180] In another example, the process is according to paragraph [0179], wherein R², if present, is fluorine.

[0181] In another example, the process is according to paragraph [0180], wherein R², if present, is up to two fluorines *ortho* to the oxygen of the phenylene to which R² is attached.



[0182] In another example, the process is according to paragraph [0130], used to make a compound listed in either Table 1 or Table 2.

[0183] In another example the process is according to any of paragraphs [0130] - [0182], further comprising converting said compound to a pharmaceutically acceptable salt, hydrate, or prodrug thereof.


Definitions

[0184] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in


which they are used indicates otherwise or they are expressly defined to mean something different.

[0185] The symbol “-” means a single bond, “=” means a double bond, “≡” means a triple bond. The symbol “” refers to a group on a double-bond as occupying either position on the terminus of a double bond to which the symbol is attached; that is, the geometry, *E*- or *Z*-, of the double bond is ambiguous. When a group is depicted removed from its parent formula, the “” symbol will be used at the end of the bond which was theoretically cleaved in order to separate the group from its parent structural formula.


[0186] When chemical structures are depicted or described, unless explicitly stated otherwise, all carbons are assumed to have hydrogen substitution to conform to a valence of four. For example, in the structure on the left-hand side of the schematic below there are nine hydrogens implied. The nine hydrogens are depicted in the right-hand structure. Sometimes a particular atom in a structure is described in textual formula as having a hydrogen or hydrogens as substitution (expressly defined hydrogen), for example, -CH₂CH₂-. It is understood by one of ordinary skill in the art that the aforementioned descriptive techniques are common in the chemical arts to provide brevity and simplicity to description of otherwise complex structures.



[0187] In this application, some ring structures are depicted generically and will be described textually. For example, in the schematic below, if in the structure on the left, ring A is used to describe a “spirocyclyl,” then if ring A is cyclopropyl, there are at most four hydrogens on ring A (when “R” can also be -H). In another example, as depicted on the right side of the schematic below, if ring B is used to describe a “phenylene” then there can be at most four hydrogens on ring B (assuming depicted cleaved bonds are not C-H bonds).




[0188] If a group “R” is depicted as “floating” on a ring system, as for example in the formula:




then, unless otherwise defined, a substituent "R" may reside on any atom of the ring system, assuming replacement of a depicted, implied, or expressly defined hydrogen from one of the ring atoms, so long as a stable structure is formed.

[0189] If a group "R" is depicted as floating on a fused ring system, as for example in the formulae:




then, unless otherwise defined, a substituent "R" may reside on any atom of the fused ring system, assuming replacement of a depicted (for example the -NH- in the formula above), implied (for example as in the formula above, where the hydrogens are not shown but understood to be present), or expressly defined hydrogen (for example where in the formula above, "X" equals =CH-) from one of the ring atoms, so long as a stable structure is formed. In the example depicted, the "R" group may reside on either the 5-membered or the 6-membered ring of the fused ring system. In the formula depicted above, when y is 2 for example, then the two "R's" may reside on any two atoms of the ring system, again assuming each replaces a depicted, implied, or expressly defined hydrogen on the ring.

[0190] When there are more than one such depicted "floating" groups, as for example in the formulae:




where there are two groups, namely, the "R" and the bond indicating attachment to a parent structure; then, unless otherwise defined, the "floating" groups may reside on any atoms of the ring system, again assuming each replaces a depicted, implied, or expressly defined hydrogen on the ring.

[0191] When a group "R" is depicted as existing on a ring system containing saturated carbons, as for example in the formula:



where, in this example, “y” can be more than one, assuming each replaces a currently depicted, implied, or expressly defined hydrogen on the ring; then, unless otherwise defined, where the resulting structure is stable, two “R’s” may reside on the same carbon. A simple example is when R is a methyl group; there can exist a geminal dimethyl on a carbon of the depicted ring (an “annular” carbon). In another example, two R’s on the same carbon, including that carbon, may form a ring, thus creating a spirocyclic ring (a “spirocyclyl” group) structure with the depicted ring as for example in the formula:



[0192] “Alkyl” is intended to include linear, branched, or cyclic hydrocarbon structures and combinations thereof, inclusively. For example, “C₈ alkyl” may refer to an *n*-octyl, *iso*-octyl, cyclohexylethyl, and the like. Lower alkyl refers to alkyl groups of from one to six carbon atoms. Examples of lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *s*-butyl, *t*-butyl, isobutyl, pentyl, hexyl and the like. Higher alkyl refers to alkyl groups containing more than eight carbon atoms. Exemplary alkyl groups are those of C₂₀ or below. Cycloalkyl is a subset of alkyl and includes cyclic hydrocarbon groups of from three to thirteen carbon atoms. Examples of cycloalkyl groups include *c*-propyl, *c*-butyl, *c*-pentyl, norbornyl, adamantyl and the like. In this application, alkyl refers to alkanyl, alkenyl, and alkynyl residues (and combinations thereof); it is intended to include cyclohexylmethyl, vinyl, allyl, isoprenyl, and the like. Thus when an alkyl residue having a specific number of carbons is named, all geometric isomers having that number of carbons are intended to be encompassed; thus, for example, either “butyl” or “C₄ alkyl” is meant to include *n*-butyl, *sec*-butyl, isobutyl, *t*-butyl, isobutenyl and but-2-yne radicals; and for example, “propyl” or “C₃ alkyl” each include *n*-propyl, propenyl, and isopropyl.

[0193] “Alkylene” refers to straight or branched chain divalent radical consisting solely of carbon and hydrogen atoms, containing no unsaturation and having from one to ten carbon atoms, for example, methylene, ethylene, propylene, *n*-butylene and the like. Alkylene is a subset of alkyl, referring to the same residues as alkyl, but having two points of attachment and, specifically, fully saturated. Examples of alkylene include ethylene

(-CH₂CH₂-), propylene (-CH₂CH₂CH₂-), dimethylpropylene (-CH₂C(CH₃)₂CH₂-), and cyclohexylpropylene (-CH₂CH₂CH(C₆H₁₃)).

[0194] "Alkylidene" refers to a straight or branched chain unsaturated divalent radical consisting solely of carbon and hydrogen atoms, having from two to ten carbon atoms, for example, ethylidene, propylidene, *n*-butylidene, and the like. Alkylidene is a subset of alkyl, referring to the same residues as alkyl, but having two points of attachment and, specifically, double bond unsaturation. The unsaturation present includes at least one double bond.

[0195] "Alkylidyne" refers to a straight or branched chain unsaturated divalent radical consisting solely of carbon and hydrogen atoms having from two to ten carbon atoms, for example, propylid-2-ynyl, *n*-butylid-1-ynyl, and the like. Alkylidyne is a subset of alkyl, referring to the same residues as alkyl, but having two points of attachment and, specifically, triple bond unsaturation. The unsaturation present includes at least one triple bond.

[0196] Any of the above radicals, "alkylene," "alkylidene" and "alkylidyne," when optionally substituted, may contain alkyl substitution which itself contains unsaturation. For example, 2-(2-phenylethynyl-but-3-enyl)-naphthalene (IUPAC name) contains an *n*-butylid-3-ynyl radical with a vinyl substituent at the 2-position of said radical.

[0197] "Alkoxy" or "alkoxyl" refers to the group -O-alkyl, for example including from one to eight carbon atoms of a straight, branched, cyclic configuration, unsaturated chains, and combinations thereof attached to the parent structure through an oxygen atom. Examples include methoxy, ethoxy, propoxy, isopropoxy, cyclopropyloxy, cyclohexyloxy and the like. Lower-alkoxy refers to groups containing one to six carbons.

[0198] "Substituted alkoxy" refers to the group -O-(substituted alkyl), the substitution on the alkyl group generally containing more than only carbon (as defined by alkoxy). One exemplary substituted alkoxy group is "polyalkoxy" or -O-optionally substituted alkylene-optionally substituted alkoxy, and includes groups such as -OCH₂CH₂OCH₃, and glycol ethers such as polyethyleneglycol and -O(CH₂CH₂O)_xCH₃, where *x* is an integer of between about two and about twenty, in another example, between about two and about ten, and in a further example between about two and about five. Another exemplary substituted alkoxy group is hydroxyalkoxy or -OCH₂(CH₂)_yOH, where *y* is for example an

integer of between about one and about ten, in another example y is an integer of between about one and about four.

[0199] "Acyl" refers to groups of from one to ten carbon atoms of a straight, branched, cyclic configuration, saturated, unsaturated and aromatic and combinations thereof, attached to the parent structure through a carbonyl functionality. One or more carbons in the acyl residue may be replaced by nitrogen, oxygen or sulfur as long as the point of attachment to the parent remains at the carbonyl. Examples include acetyl, benzoyl, propionyl, isobutyryl, *t*-butoxycarbonyl, benzyloxycarbonyl and the like. Lower-acyl refers to groups containing one to six carbons.

[0200] " α -Amino Acids" refer to naturally occurring and commercially available amino acids and optical isomers thereof. Typical natural and commercially available α -amino acids are glycine, alanine, serine, homoserine, threonine, valine, norvaline, leucine, isoleucine, norleucine, aspartic acid, glutamic acid, lysine, ornithine, histidine, arginine, cysteine, homocysteine, methionine, phenylalanine, homophenylalanine, phenylglycine, ortho-tyrosine, meta-tyrosine, para-tyrosine, tryptophan, glutamine, asparagine, proline and hydroxyproline. A "side chain of an α -amino acid" refers to the radical found on the α -carbon of an α -amino acid as defined above, for example, hydrogen (for glycine), methyl (for alanine), benzyl (for phenylalanine), and the like.


[0201] "Amino" refers to the group -NH_2 . "Substituted amino," refers to the group -N(H)R or -N(R)R where each R is independently selected from the group: optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted heterocyclyl, acyl, carboxy, alkoxycarbonyl, sulfanyl, sulfinyl and sulfonyl, for example, diethylamino, methylsulfonylamino, furanyl-oxy-sulfonamino.

[0202] "Aryl" refers to aromatic six- to fourteen-membered carbocyclic ring, for example, benzene, naphthalene, indane, tetralin, fluorene and the like, univalent radicals. As univalent radicals, the aforementioned ring examples are named, phenyl, naphthyl, indanyl, tetralinyl, and fluorenyl.

[0203] "Arylene" generically refers to any aryl that has at least two groups attached thereto. For a more specific example, "phenylene" refers to a divalent phenyl ring radical. A phenylene, thus may have more than two groups attached, but is defined by a minimum of two non-hydrogen groups attached thereto.

[0204] "Arylalkyl" refers to a residue in which an aryl moiety is attached to a parent structure via one of an alkylene, alkylidene, or alkylidyne radical. Examples include benzyl, phenethyl, phenylvinyl, phenylallyl and the like. Both the aryl, and the corresponding alkylene, alkylidene, or alkylidyne radical portion of an arylalkyl group may be optionally substituted. "Lower arylalkyl" refers to an arylalkyl where the "alkyl" portion of the group has one to six carbons; this can also be referred to as C₁₋₆ arylalkyl.

[0205] "Exo-alkenyl" refers to a double bond that emanates from an annular carbon, and is not within the ring system, for example the double bond depicted in the formula below.



[0206] In some examples, as appreciated by one of ordinary skill in the art, two adjacent groups on an aromatic system may be fused together to form a ring structure. The fused ring structure may contain heteroatoms and may be optionally substituted with one or more groups. It should additionally be noted that saturated carbons of such fused groups (i.e. saturated ring structures) can contain two substitution groups.

[0207] "Fused-polycyclic" or "fused ring system" refers to a polycyclic ring system that contains bridged or fused rings; that is, where two rings have more than one shared atom in their ring structures. In this application, fused-polycyclics and fused ring systems are not necessarily all aromatic ring systems. Typically, but not necessarily, fused-polycyclics share a vicinal set of atoms, for example naphthalene or 1,2,3,4-tetrahydro-naphthalene. A spiro ring system is not a fused-polycyclic by this definition, but fused polycyclic ring systems of the invention may themselves have spiro rings attached thereto via a single ring atom of the fused-polycyclic.

[0208] "Halogen" or "halo" refers to fluorine, chlorine, bromine or iodine. "Haloalkyl" and "haloaryl" refer generically to alkyl and aryl radicals that are substituted with one or more halogens, respectively. Thus, "dihaloaryl," "dihaloalkyl," "trihaloaryl" etc. refer to aryl and alkyl substituted with a plurality of halogens, but not necessarily a plurality of the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl.

[0209] "Heteroarylene" generically refers to any heteroaryl that has at least two groups attached thereto. For a more specific example, "pyridylene" refers to a divalent pyridyl

ring radical. A pyridylene, thus may have more than two groups attached, but is defined by a minimum of two non-hydrogen groups attached thereto.

[0210] "Heteroatom" refers to O, S, N, or P.

[0211] "Heterocyclyl" refers to a stable three- to fifteen-membered ring radical that consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, phosphorus, oxygen and sulfur. For purposes of this invention, the heterocyclyl radical may be a monocyclic, bicyclic or tricyclic ring system, which may include fused or bridged ring systems as well as spirocyclic systems; and the nitrogen, phosphorus, carbon or sulfur atoms in the heterocyclyl radical may be optionally oxidized to various oxidation states. In a specific example, the group $-S(O)_{0-2}-$, refers to $-S-$ (sulfide), $-S(O)-$ (sulfoxide), and $-SO_2-$ (sulfone). For convenience, nitrogens, particularly but not exclusively, those defined as annular aromatic nitrogens, are meant to include their corresponding *N*-oxide form, although not explicitly defined as such in a particular example. Thus, for a compound of the invention having, for example, a pyridyl ring; the corresponding pyridyl-*N*-oxide is meant to be included as another compound of the invention. In addition, annular nitrogen atoms may be optionally quaternized; and the ring radical may be partially or fully saturated or aromatic. Examples of heterocyclyl radicals include, but are not limited to, azetidiny, acridiny, benzodioxoly, benzodioxany, benzofurany, carbazoy, cinnolinyl, dioxolany, indoliziny, naphthyridiny, perhydroazepiny, phenaziny, phenothiaziny, phenoxaziny, phthalaziny, pteridiny, puriny, quinazolinyl, quinoxaliny, quinolinyl, isoquinolinyl, tetrazoyl, tetrahydroisoquinoly, piperidiny, piperaziny, 2-oxopiperaziny, 2-oxopiperidiny, 2-oxopyrrolidiny, 2-oxoazepiny, azepiny, pyrroly, 4-piperidonyl, pyrrolidiny, pyrazoly, pyrazolidiny, imidazoly, imidazoliny, imidazolidiny, dihydropyridiny, tetrahydropyridiny, pyridiny, pyraziny, pyrimidiny, pyridaziny, oxazoly, oxazoliny, oxazolidiny, triazoly, isoxazoly, isoxazolidiny, morpholinyl, thiazoly, thiazoliny, thiazolidiny, isothiazoly, quinuclidiny, isothiazolidiny, indoly, isoindoly, indoliny, isoindoliny, octahydroindoly, octahydroisoindoly, quinoly, isoquinoly, decahydroisoquinoly, benzimidazoly, thiadiazoly, benzopyrany, benzothiazoly, benzoxazoly, furyl, tetrahydrofuryl, tetrahydropyrany, thienyl, benzothieliyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, dioxaphospholany, and oxadiazoly.

[0212] “Heteroalicyclic” refers specifically to a non-aromatic heterocyclyl radical. A heteroalicyclic may contain unsaturation, but is not aromatic.


[0213] “Heteroaryl” refers specifically to an aromatic heterocyclyl radical.

[0214] “Heterocyclylalkyl” refers to a residue in which a heterocyclyl is attached to a parent structure via one of an alkylene, alkylidene, or alkylidyne radical. Examples include (4-methylpiperazin-1-yl) methyl, (morpholin-4-yl) methyl, (pyridine-4-yl) methyl, 2-(oxazolin-2-yl) ethyl, 4-(4-methylpiperazin-1-yl)-2-butenyl, and the like. Both the heterocyclyl, and the corresponding alkylene, alkylidene, or alkylidyne radical portion of a heterocyclylalkyl group may be optionally substituted. “Lower heterocyclylalkyl” refers to a heterocyclylalkyl where the “alkyl” portion of the group has one to six carbons. “Heteroalicycylalkyl” refers specifically to a heterocyclylalkyl where the heterocyclyl portion of the group is non-aromatic; and “heteroarylalkyl” refers specifically to a heterocyclylalkyl where the heterocyclyl portion of the group is aromatic. Such terms may be described in more than one way, for example, “lower heterocyclylalkyl” and “heterocyclyl C₁₋₆alkyl” are equivalent terms.

[0215] “Optional” or “optionally” means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. One of ordinary skill in the art would understand that, with respect to any molecule described as containing one or more optional substituents, that only sterically practical and/or synthetically feasible compounds are meant to be included. “Optionally substituted” refers to all subsequent modifiers in a term, for example in the term “optionally substituted arylC₁₋₈ alkyl,” optional substitution may occur on both the “C₁₋₈ alkyl” portion and the “aryl” portion of the molecule; and for example, optionally substituted alkyl includes optionally substituted cycloalkyl groups, which in turn are defined as including optionally substituted alkyl groups, potentially *ad infinitum*. A list of exemplary optional substitution are listed below in the definition of “substituted.”

[0216] “Saturated bridged ring system” refers to a bicyclic or polycyclic ring system that is not aromatic. Such a system may contain isolated or conjugated unsaturation, but not aromatic or heteroaromatic rings in its core structure (but may have aromatic substitution thereon). For example, hexahydro-furo[3,2-b]furan, 2,3,3a,4,7,7a-hexahydro-1H-indene, 7-aza-bicyclo[2.2.1]heptane, and 1,2,3,4,4a,5,8,8a-octahydro-naphthalene are all included in the class “saturated bridged ring system.”

[0217] “Spirocyclyl” or “spirocyclic ring” refers to a ring originating from a particular annular carbon of another ring. For example, as depicted below, a ring atom of a saturated bridged ring system (rings B and B’), but not a bridgehead atom, can be a shared atom between the saturated bridged ring system and a spirocyclyl (ring A) attached thereto. A spirocyclyl can be carbocyclic or heteroalicyclic.



[0218] “Substituted” alkyl, aryl, and heterocyclyl, refer respectively to alkyl, aryl, and heterocyclyl, wherein one or more (for example up to about five, in another example, up to about three) hydrogen atoms are replaced by a substituent independently selected from: optionally substituted alkyl (for example, fluoromethyl), optionally substituted aryl (for example, 4-hydroxyphenyl), optionally substituted arylalkyl (for example, 1-phenyl-ethyl), optionally substituted heterocyclylalkyl (for example, 1-pyridin-3-yl-ethyl), optionally substituted heterocyclyl (for example, 5-chloro-pyridin-3-yl or 1-methyl-piperidin-4-yl), optionally substituted alkoxy, alkylenedioxy (for example methylenedioxy), optionally substituted amino (for example, alkylamino and dialkylamino), optionally substituted amidino, optionally substituted aryloxy (for example, phenoxy), optionally substituted arylalkyloxy (for example, benzyloxy), carboxy ($-\text{CO}_2\text{H}$), carboalkoxy (that is, acyloxy or $-\text{OC}(=\text{O})\text{R}$), carboxyalkyl (that is, esters or $-\text{CO}_2\text{R}$), carboxamido, benzyloxycarbonylamino (CBZ-amino), cyano, acyl, halogen, hydroxy, nitro, sulfanyl, sulfinyl, sulfonyl, thiol, halogen, hydroxy, oxo, carbamyl, acylamino, and sulfonamido.

[0219] “Suitable leaving group” is defined as the term would be understood by one of ordinary skill in the art; that is, a carbon with such a group attached, upon reaction wherein a new bond is to be formed, loses such a group upon formation of the new bond. The invention pertains particularly with respect convergent synthesis, to reactions where such a leaving group is bonded to a reaction partner that is aromatic, undergoes a bond-forming reaction and remains aromatic. A typical example of such a reaction is a nucleophilic aromatic substitution reaction, as would be understood by one of ordinary skill in the art. However, the invention is not limited to such mechanistic restrictions; for example, reactions where there is, for example, an insertion reaction (for example by a transition metal) into the bond between the aromatic reaction partner and its leaving group

followed by reductive coupling can also be used within the scope of the invention. Examples of suitable leaving groups include halogens, optionally substituted aryl or alkyl sulfonates, phosphonates, azides, $RS(O)_{0-2}$ where R is, for example optionally substituted alkyl, optionally substituted aryl, or optionally substituted heteroaryl.

[0220] "Sulfanyl" refers to the groups: -S-(optionally substituted alkyl), -S-(optionally substituted aryl), and -S-(optionally substituted heterocyclyl).

[0221] "Sulfinyl" refers to the groups: -S(O)-H, -S(O)-(optionally substituted alkyl), -S(O)-optionally substituted aryl, and -S(O)-(optionally substituted heterocyclyl).

[0222] "Sulfonyl" refers to the groups: -S(O₂)-H, -S(O₂)-(optionally substituted alkyl), -S(O₂)-optionally substituted aryl, -S(O₂)-(optionally substituted heterocyclyl), -S(O₂)-(optionally substituted alkoxy), -S(O₂)-optionally substituted aryloxy, and -S(O₂)-(optionally substituted heterocycliloxy).

[0223] "Yield" for each of the reactions described herein is expressed as a percentage of the theoretical yield.

[0224] Some of the compounds of the invention may have imino, amino, oxo or hydroxy substituents off aromatic heterocyclyl systems. For purposes of this disclosure, it is understood that such imino, amino, oxo or hydroxy substituents may exist in their corresponding tautomeric form, i.e., amino, imino, hydroxy or oxo, respectively.

[0225] Compounds of the invention are named according to systematic application of the nomenclature rules agreed upon by the International Union of Pure and Applied Chemistry (IUPAC), International Union of Biochemistry and Molecular Biology (IUBMB), and the Chemical Abstracts Service (CAS).

[0226] The compounds of the invention, or their pharmaceutically acceptable salts, may have asymmetric carbon atoms, oxidized sulfur atoms or quaternized nitrogen atoms in their structure.

[0227] The compounds of the invention and their pharmaceutically acceptable salts may exist as single stereoisomers, racemates, and as mixtures of enantiomers and diastereomers. The compounds may also exist as geometric isomers. All such single stereoisomers, racemates and mixtures thereof, and geometric isomers are intended to be within the scope of this invention.

[0228] It is assumed that when considering generic descriptions of compounds of the invention for the purpose of constructing a compound, such construction results in the creation of a stable structure. That is, one of ordinary skill in the art would recognize that there can theoretically be some constructs which would not normally be considered as stable compounds (that is, sterically practical and/or synthetically feasible, *supra*).

[0229] When a particular group with its bonding structure is denoted as being bonded to two partners; that is, a divalent radical, for example, $\text{-OCH}_2\text{-}$, then it is understood that either of the two partners may be bound to the particular group at one end, and the other partner is necessarily bound to the other end of the particular group, unless stated explicitly otherwise. Stated another way, divalent radicals are not to be construed as limited to the depicted orientation, for example " $\text{-OCH}_2\text{-}$ " is meant to mean not only " $\text{-OCH}_2\text{-}$ " as drawn, but also " $\text{-CH}_2\text{O-}$."

[0230] Methods for the preparation and/or separation and isolation of single stereoisomers from racemic mixtures or non-racemic mixtures of stereoisomers are well known in the art. For example, optically active (R)- and (S)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. Enantiomers (R- and S-isomers) may be resolved by methods known to one of ordinary skill in the art, for example by: formation of diastereoisomeric salts or complexes which may be separated, for example, by crystallization; via formation of diastereoisomeric derivatives which may be separated, for example, by crystallization, selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where a desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be required to liberate the desired enantiomeric form. Alternatively, specific enantiomer may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation. For a mixture of enantiomers, enriched in a particular enantiomer, the major component enantiomer may be further enriched (with concomitant loss in yield) by recrystallization.

[0231] "Patient" for the purposes of the present invention includes humans and other animals, particularly mammals, and other organisms. Thus the methods are applicable to

both human therapy and veterinary applications. In a preferred embodiment the patient is a mammal, and in a most preferred embodiment the patient is human.

[0232] “Kinase-dependent diseases or conditions” refer to pathologic conditions that depend on the activity of one or more protein kinases. Kinases either directly or indirectly participate in the signal transduction pathways of a variety of cellular activities including proliferation, adhesion, migration, differentiation and invasion. Diseases associated with kinase activities include tumor growth, the pathologic neovascularization that supports solid tumor growth, and associated with other diseases where excessive local vascularization is involved such as ocular diseases (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

[0233] While not wishing to be bound to theory, phosphatases can also play a role in “kinase-dependent diseases or conditions” as cognates of kinases; that is, kinases phosphorylate and phosphatases dephosphorylate, for example protein substrates. Therefore compounds of the invention, while modulating kinase activity as described herein, may also modulate, either directly or indirectly, phosphatase activity. This additional modulation, if present, may be synergistic (or not) to activity of compounds of the invention toward a related or otherwise interdependent kinase or kinase family. In any case, as stated previously, the compounds of the invention are useful for treating diseases characterized in part by abnormal levels of cell proliferation (i.e. tumor growth), programmed cell death (apoptosis), cell migration and invasion and angiogenesis associated with tumor growth.

[0234] “Therapeutically effective amount” is an amount of a compound of the invention, that when administered to a patient, ameliorates a symptom of the disease. The amount of a compound of the invention which constitutes a “therapeutically effective amount” will vary depending on the compound, the disease state and its severity, the age of the patient to be treated, and the like. The therapeutically effective amount can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

[0235] “Cancer” refers to cellular-proliferative disease states, including but not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma),

alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiforme, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma [embryonal rhabdomyosarcoma], fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma,

dermatofibroma, keloids, psoriasis; and Adrenal lands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions.

[0236] "Pharmaceutically acceptable acid addition salt" refers to those salts that retain the biological effectiveness of the free bases and that are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like, as well as organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

[0237] "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Exemplary salts are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins, and the like. Exemplary organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline, and caffeine. (See, for example, S. M. Berge, et al., "Pharmaceutical Salts," J. Pharm. Sci., 1977;66:1-19 which is incorporated herein by reference.)

[0238] "Prodrug" refers to compounds that are transformed (typically rapidly) *in vivo* to yield the parent compound of the above formulae, for example, by hydrolysis in blood. Common examples include, but are not limited to, ester and amide forms of a compound having an active form bearing a carboxylic acid moiety. Examples of pharmaceutically acceptable esters of the compounds of this invention include, but are not limited to, alkyl esters (for example with between about one and about six carbons) wherein the alkyl group is a straight or branched chain. Acceptable esters also include cycloalkyl esters and

arylalkyl esters such as, but not limited to benzyl. Examples of pharmaceutically acceptable amides of the compounds of this invention include, but are not limited to, primary amides, and secondary and tertiary alkyl amides (for example with between about one and about six carbons). Amides and esters of the compounds of the present invention may be prepared according to conventional methods. A thorough discussion of prodrugs is provided in T. Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference for all purposes.

[0239] "Metabolite" refers to the break-down or end product of a compound or its salt produced by metabolism or biotransformation in the animal or human body; for example, biotransformation to a more polar molecule such as by oxidation, reduction, or hydrolysis, or to a conjugate (see Goodman and Gilman, "The Pharmacological Basis of Therapeutics" 8^{sup}.th Ed., Pergamon Press, Gilman et al. (eds), 1990 for a discussion of biotransformation). As used herein, the metabolite of a compound of the invention or its salt may be the biologically active form of the compound in the body. In one example, a prodrug may be used such that the biologically active form, a metabolite, is released *in vivo*. In another example, a biologically active metabolite is discovered serendipitously, that is, no prodrug design *per se* was undertaken. An assay for activity of a metabolite of a compound of the present invention is known to one of skill in the art in light of the present disclosure.

[0240] In addition, the compounds of the present invention can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.

[0241] In addition, it is intended that the present invention cover compounds made either using standard organic synthetic techniques, including combinatorial chemistry or by biological methods, such as bacterial digestion, metabolism, enzymatic conversion, and the like.

[0242] "Treating" or "treatment" as used herein covers the treatment of a disease-state in a human, which disease-state is characterized by abnormal cellular proliferation, and invasion and includes at least one of: (i) preventing the disease-state from occurring in a human, in particular, when such human is predisposed to the disease-state but has not yet

been diagnosed as having it; (ii) inhibiting the disease-state, i.e., arresting its development; and (iii) relieving the disease-state, i.e., causing regression of the disease-state. As is known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by one of ordinary skill in the art.

[0243] One of ordinary skill in the art would understand that certain crystallized, protein-ligand complexes, in particular c-Met, c-Kit, KDR, flt-3, or flt-4-ligand complexes, and their corresponding x-ray structure coordinates can be used to reveal new structural information useful for understanding the biological activity of kinases as described herein. As well, the key structural features of the aforementioned proteins, particularly, the shape of the ligand binding site, are useful in methods for designing or identifying selective modulators of kinases and in solving the structures of other proteins with similar features. Such protein-ligand complexes, having compounds of the invention as their ligand component, are an aspect of the invention.

[0244] As well, one of ordinary skill in the art would appreciate that such suitable x-ray quality crystals can be used as part of a method of identifying a candidate agent capable of binding to and modulating the activity of kinases. Such methods may be characterized by the following aspects: a) introducing into a suitable computer program, information defining a ligand binding domain of a kinase in a conformation (e.g. as defined by x-ray structure coordinates obtained from suitable x-ray quality crystals as described above) wherein the computer program creates a model of the three dimensional structures of the ligand binding domain, b) introducing a model of the three dimensional structure of a candidate agent in the computer program, c) superimposing the model of the candidate agent on the model of the ligand binding domain, and d) assessing whether the candidate agent model fits spatially into the ligand binding domain. Aspects a-d are not necessarily carried out in the aforementioned order. Such methods may further entail: performing rational drug design with the model of the three-dimensional structure, and selecting a potential candidate agent in conjunction with computer modeling.

[0245] Additionally, one skilled in the art would appreciate that such methods may further entail: employing a candidate agent, so-determined to fit spatially into the ligand binding domain, in a biological activity assay for kinase modulation, and determining whether said candidate agent modulates kinase activity in the assay. Such methods may also include

administering the candidate agent, determined to modulate kinase activity, to a mammal suffering from a condition treatable by kinase modulation, such as those described above.

[0246] Also, one skilled in the art would appreciate that compounds of the invention can be used in a method of evaluating the ability of a test agent to associate with a molecule or molecular complex comprising a ligand binding domain of a kinase. Such a method may be characterized by the following aspects: a) creating a computer model of a kinase binding pocket using structure coordinates obtained from suitable x-ray quality crystals of the kinase, b) employing computational algorithms to perform a fitting operation between the test agent and the computer model of the binding pocket, and c) analyzing the results of the fitting operation to quantify the association between the test agent and the computer model of the binding pocket.

General Administration

[0247] Administration of the compounds of the invention, or their pharmaceutically acceptable salts, in pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration or agents for serving similar utilities. Thus, administration can be, for example, orally, nasally, parenterally (intravenous, intramuscular, or subcutaneous), topically, transdermally, intravaginally, intravesically, intracisternally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages.

[0248] The compositions will include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, etc. Compositions of the invention may be used in combination with anticancer or other agents that are generally administered to a patient being treated for cancer. Adjuvants include preserving, wetting, suspending, sweetening, flavoring, perfuming, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form

can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0249] If desired, a pharmaceutical composition of the invention may also contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, antioxidants, and the like, such as, for example, citric acid, sorbitan monolaurate, triethanolamine oleate, butylated hydroxytoluene, etc.

[0250] Compositions suitable for parenteral injection may comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

[0251] One preferable route of administration is oral, using a convenient daily dosage regimen that can be adjusted according to the degree of severity of the disease-state to be treated.

[0252] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders, as for example, cellulose derivatives, starch, alginates, gelatin, polyvinylpyrrolidone, sucrose, and gum acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, croscarmellose sodium, complex silicates, and sodium carbonate, (e) solution retarders, as for example paraffin, (f) absorption accelerators, as for example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol, and glycerol monostearate, magnesium stearate and the like (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents.

[0253] Solid dosage forms as described above can be prepared with coatings and shells, such as enteric coatings and others well known in the art. They may contain pacifying agents, and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedded compositions that can be used are polymeric substances and waxes. The active compounds can also be in microencapsulated form, if appropriate, with one or more of the above-mentioned excipients.

[0254] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. Such dosage forms are prepared, for example, by dissolving, dispersing, etc., a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol and the like; solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, dimethylformamide; oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethyleneglycols and fatty acid esters of sorbitan; or mixtures of these substances, and the like, to thereby form a solution or suspension.

[0255] Suspensions, in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, or mixtures of these substances, and the like.

[0256] Compositions for rectal administrations are, for example, suppositories that can be prepared by mixing the compounds of the present invention with for example suitable non-irritating excipients or carriers such as cocoa butter, polyethyleneglycol or a suppository wax, which are solid at ordinary temperatures but liquid at body temperature and therefore, melt while in a suitable body cavity and release the active component therein.

[0257] Dosage forms for topical administration of a compound of this invention include ointments, powders, sprays, and inhalants. The active component is admixed under sterile conditions with a physiologically acceptable carrier and any preservatives, buffers, or propellants as may be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

[0258] Generally, depending on the intended mode of administration, the pharmaceutically acceptable compositions will contain about 1% to about 99% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and 99% to 1% by weight of a suitable pharmaceutical excipient. In one example, the composition will be between about 5% and about 75% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, with the rest being suitable pharmaceutical excipients.

[0259] Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, 18th Ed., (Mack Publishing Company, Easton, Pa., 1990). The composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state in accordance with the teachings of this invention.

[0260] The compounds of the invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount which will vary depending upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of the compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular disease-states, and the host undergoing therapy. The compounds of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 1,000 mg per day. For a normal human adult having a body weight of about 70 kilograms, a dosage in the range of about 0.01 to about 100 mg per kilogram of body weight per day is an example. The specific dosage used, however, can vary. For example, the dosage can depend on a number of factors including the requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the compound being used. The determination of optimum dosages for a particular patient is well known to one of ordinary skill in the art.

Utility of compounds of the invention as screening agents

[0261] To employ the compounds of the invention in a method of screening for candidate agents that bind to, for example c-Met, KDR, c-Kit, flt-3, or flt-4, the protein is bound to a support, and a compound of the invention is added to the assay. Alternatively, the

compound of the invention is bound to the support and the protein is added. Classes of candidate agents among which novel binding agents may be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

[0262] The determination of the binding of the candidate agent to, for example, c-Met, KDR, c-Kit, flt-3, or flt-4 protein may be done in a number of ways. In one example, the candidate agent (the compound of the invention) is labeled, for example, with a fluorescent or radioactive moiety and binding determined directly. For example, this may be done by attaching all or a portion of the c-Met, KDR, c-Kit, flt-3, or flt-4 protein to a solid support, adding a labeled agent (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

[0263] By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g., radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

[0264] In some embodiments, only one of the components is labeled. For example, c-Met, KDR, c-Kit, flt-3, or flt-4 protein may be labeled at tyrosine positions using ^{125}I , or with fluorophores. Alternatively, more than one component may be labeled with different labels; using ^{125}I for the proteins, for example, and a fluorophor for the candidate agents.

[0265] The compounds of the invention may also be used as competitors to screen for additional drug candidates. "Candidate bioactive agent" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity.

They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. In the case where protein binding or activity is screened, some embodiments exclude molecules already known to bind to that particular protein. Exemplary embodiments of assays described herein include candidate agents, which do not bind the target protein in its endogenous native state, termed herein as "exogenous" agents. In one example, exogenous agents further exclude antibodies to c-Met, KDR, c-Kit, flt-3, or flt-4.

[0266] Candidate agents can encompass numerous chemical classes, though typically they are organic molecules having a molecular weight of more than about 100 daltons and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, for example at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclyl structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs, or combinations thereof.

[0267] Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

[0268] In one example, the binding of the candidate agent is determined through the use of competitive binding assays. In this example, the competitor is a binding moiety known to bind to c-Met, KDR, c-Kit, flt-3, or flt-4, such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the

candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

[0269] In some embodiments, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to for example c-Met, KDR, c-Kit, flt-3, or flt-4 for a time sufficient to allow binding, if present. Incubations may be performed at any temperature that facilitates optimal activity, typically between 4°C and 40°C

[0270] Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

[0271] In one example, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is binding to c-Met, KDR, c-Kit, flt-3, or flt-4 and thus is capable of binding to, and potentially modulating, the activity of the c-Met, KDR, c-Kit, flt-3, or flt-4. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

[0272] In an alternative embodiment, the candidate agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate the candidate agent is bound to c-Met, KDR, c-Kit, flt-3, or flt-4 with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable of binding to c-Met, KDR, c-Kit, flt-3, or flt-4.

[0273] It may be of value to identify the binding site of c-Met, KDR, c-Kit, flt-3, or flt-4. This can be done in a variety of ways. In one embodiment, once c-Met, KDR, c-Kit, flt-3, or flt-4 has been identified as binding to the candidate agent, the c-Met, KDR, c-Kit, flt-3, or flt-4 is fragmented or modified and the assays repeated to identify the necessary components for binding.

[0274] Modulation is tested by screening for candidate agents capable of modulating the activity of c-Met, KDR, c-Kit, flt-3, or flt-4 comprising the steps of combining a candidate agent with c-Met, KDR, c-Kit, flt-3, or flt-4, as above, and determining an alteration in the

biological activity of the c-Met, KDR, c-Kit, flt-3, or flt-4. Thus, in this embodiment, the candidate agent should both bind to (although this may not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both *in vitro* screening methods and *in vivo* screening of cells for alterations in cell viability, morphology, and the like.

[0275] Alternatively, differential screening may be used to identify drug candidates that bind to native c-Met, KDR, c-Kit, flt-3, or flt-4, but cannot bind to modified c-Met, KDR, c-Kit, flt-3, or flt-4.

[0276] Positive controls and negative controls may be used in the assays. For example, all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of samples is for a time sufficient for the binding of the agent to the protein. Following incubation, samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

[0277] A variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

Abbreviations and their Definitions

[0278] The following abbreviations and terms have the indicated meanings throughout.

Abbreviation	Meaning
Ac	acetyl
ATP	adenosine triphosphate
BNB	4-bromomethyl-3-nitrobenzoic acid
Boc	t-butyloxy carbonyl

Abbreviation	Meaning
br	broad
Bu	butyl
°C	degrees Celsius
c-	cyclo
CBZ	CarboBenZoxy = benzyloxycarbonyl
d	doublet
dd	doublet of doublet
dt	doublet of triplet
DBU	Diazabicyclo[5.4.0]undec-7-ene
DCM	dichloromethane = methylene chloride = CH ₂ Cl ₂
DCE	dichloroethylene
DEAD	diethyl azodicarboxylate
DIC	diisopropylcarbodiimide
DIEA	N,N-diisopropylethyl amine
DMAP	4-N,N-dimethylaminopyridine
DMF	N,N-dimethylformamide
DMSO	dimethyl sulfoxide
DVB	1,4-divinylbenzene
EEDQ	2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline
EI	Electron Impact ionization
Et	ethyl
Fmoc	9-fluorenylmethoxycarbonyl
g	gram(s)
GC	gas chromatography
h or hr	hour(s)
HATU	O-(7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HMDS	hexamethyldisilazane
HOAc	acetic acid
HOBt	hydroxybenzotriazole
HPLC	high pressure liquid chromatography


Abbreviation	Meaning
L	liter(s)
M	molar or molarity
m	multiplet
Me	methyl
mesyl	methanesulfonyl
mg	milligram(s)
MHz	megahertz (frequency)
Min	minute(s)
mL	milliliter(s)
mM	millimolar
mmol	millimole(s)
mol	mole(s)
MS	mass spectral analysis
MTBE	methyl t-butyl ether
N	normal or normality
NBS	N-bromosuccinimide
NCS	N-chlorosuccinimide
nM	nanomolar
NMO	N-methylmorpholine oxide
NMR	nuclear magnetic resonance spectroscopy
PEG	polyethylene glycol
pEY	poly-glutamine, tyrosine
Ph	phenyl
PhOH	phenol
PfP	pentafluorophenol
PfPy	pentafluoropyridine
PPTS	Pyridinium p-toluenesulfonate
Py	pyridine
PyBroP	bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
q	quartet

Abbreviation	Meaning
RT	Room temperature
Sat'd	saturated
s	singlet
s-	secondary
t-	tertiary
t or tr	triplet
TBDMS	t-butyldimethylsilyl
TES	triethylsilane
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TMOF	trimethyl orthoformate
TMS	trimethylsilyl
tosyl	p-toluenesulfonyl
Trt	triphenylmethyl
uL	microliter(s)
uM	Micromole(s) or micromolar

Synthesis of Compounds

[0279] Schemes 1 and 2 depict general synthetic routes for compounds of the invention and are not intended to be limiting. More specifically, Scheme 1 depicts synthesis of quinazoline compounds, and Scheme 2 depicts synthesis of quinoline compounds. Specific examples are described subsequently to these general synthetic descriptions so as to allow one skilled in the art to make and use either quinazolines or quinolines of the invention.


Scheme 1



[0280] Referring to Scheme 1, a benzoic ester 1, where R is typically but not necessarily a methyl radical and P is typically but not necessarily an alkyl group, is O-alkylated at the oxygen *para* to the carboxylate group with an electrophile to afford a substituted derivative 2. P is typically a lower alkyl group, but may be a protecting group that is removed later in a synthesis. When P is a lower alkyl group it may possess functionality initially, or be derivitized to contain such functionality at various stages of the synthesis. The group, E¹, may represent either a protecting group, e.g. benzyl, or a group that either has moieties present in compounds of the invention or possesses functionality that serve as precursors to such groups. Aromatic ring nitration and reduction of the corresponding nitro group are carried out in a regio- and chemoselective manner by methods well known in the art to give anthranilate derivative 3. Formation of quinazolin-4-one 4 is carried out by methods well known in the art, for example by heating 3 in formamide solution in the presence of ammonium formate or for example by heating directly with formamidine hydrochloride. Introduction of 4-position functionality groups is carried out by methods known in the art. For example, quinazolin-4-one 4 is converted to an intermediate quinazoline 5, where "L" represents a leaving group, e.g. chlorine. Quinazoline 5 is then converted to 6 by reaction with a range of nucleophiles, e.g. amines, alcohols, and thiols. After formation of 6, group "Z" is either left "as is" or converted at some subsequent stage to a derivative thereof. For example when Z is -NH-, then the hydrogen on the nitrogen

may optionally be replaced with an alkyl group, or when Z is sulfur, then that sulfur atom may be oxidized to, for example, a sulfone. Structure 6 may represent compounds of the invention or, for example when E¹ serves as a protecting group, E¹ may be removed to provide phenol 7. Introduction of a group E² is carried out by methods well established in the art; for example alkylation with an appropriately derivatized alkyl halide (or mesylate or the like) to give 8 which also represents compounds of the invention.

Scheme 2



[0281] Scheme 2 shows a general route used to make exemplary quinolines of the invention. For example, compound 9 contains an alkyl group, R¹, a protecting group, P. The arrangement of the protected and alkylated phenolic oxygens may vary from the pattern depicted in compound 9. Compound 9 is nitrated to provide compound 10. The nitro group of compound 10 is reduced to give aniline 11. Compound 11 is treated, for example, with ethyl formate under basic conditions followed by acidification and isolation to form 4-hydroxy quinoline 12. Quinoline 12 may be converted to compounds of the invention in a number of ways. For example, the 4-oxygen is used as a nucleophile in a nucleophilic aromatic substitution reaction to form quinoline-aryl-ether 13. In another example, compound 13 is further derivatized, via removal of protecting group P, to afford compound 14. The 7-hydroxy of compound 14 is alkylated, for example with electrophile E, to provide a compound of the invention. As discussed in relation to Scheme 1,

variations on any of the above steps are possible, and intermediates in these schemes, for example compounds **12**, **13**, and **14** may also be compounds of the invention according to formula **I**. Also, for example, the 4-hydroxy quinoline compound **12** are converted to a corresponding 4-nitrogen or 4-sulfur quinoline using chemistry known in the art to make compounds of the invention, or alternatively the corresponding 4-nitrogen or 4-sulfur quinolines are made via routes analogous to that depicted in Schemes 1 and 2.


[0282] Schemes 1 and 2 are illustrative of quinolines and quinazolines having oxygen substitution at their respective 6- and 7-positions; the invention is not so limited, but rather is intended to encompass quinolines and quinazolines not necessarily having substitution, oxygen or otherwise, at their respective 6- or 7-positions.

[0283] Schemes 3 and 4 depict generalized synthetic routes to show the process of the invention to make compounds of formula **XXI** and is not intended to be limiting. More specifically, Schemes 3 and 4 depict convergent syntheses of quinoline and quinazoline compounds as described herein. Specific examples are described subsequently to this general synthetic description so as to allow one of ordinary skill in the art to practice the invention.

[0284] Referring to Scheme 3, a benzoic ester **16** for example, where R is typically but not necessarily a methyl radical and R¹ is typically but not necessarily one or more alkoxy or hydroxy groups. In a typical synthesis, at least one of R¹ within Scheme 3 is a hydroxyl which is converted (or protected) via one or more steps to a group important to the activity of the compounds as described as kinase modulators (in the case that -OH itself is desired in the final compound, then deprotection affords the -OH, *vide supra*). Preferably, but not necessarily, this group is complete once the synthesis of **XXII** is complete. By building desired complexity into **XXII** prior to combination with **XXIII**, convergent syntheses' advantages over serial syntheses are realized more fully. Regioselective aromatic ring nitration, and reduction of the corresponding nitro group, are carried out in a regio- and chemoselective manner by methods well known in the art to give anthranilate derivative **17**. Formation of quinazoline or quinoline 4-one **18** is carried out by methods well known in the art. For example by heating **17** in formamide solution in the presence of ammonium formate, or by heating **17** with formamidine hydrochloride, the quinazoline-4-one analog is made. In another example **17** is treated, for example, with ethyl formate under basic conditions followed by acidification and isolation to form the 4-hydroxy quinoline analog (a tautomer of the 4-one). In this scheme J' represents either carbon or nitrogen atom with


the appropriate number of hydrogens to fill their respective normal valence bonding schemes; J' is a precursor to J. Radicals J and R⁷⁰ are in accord with formula **XXI**. Introduction of 4-position functionality is carried out by methods known in the art. For example, 4-one **18** is converted to **XXII**, where "P¹" represents a suitable leaving group (in accord with formula **XXI**), e.g. chlorine (via dehydration/chlorination of **18** to give **XXII**). In another example, a 4-hydroxy analog is converted to a sulfonyl ester, e.g. the trifluoromethane sulfonate.

Scheme 3




[0285] Scheme 4 shows a general route used to make compounds of formula **XXIII**. For example, aromatic compound **19**, where "X" is a leaving group, such as fluorine and "E" is an electron withdrawing group such as nitro, is converted to **20** by reaction with a range of nucleophiles, e.g. amines, alcohols, and thiols (where "Z" is oxygen, nitrogen (substituted or not), or sulfur). In this case, "R" represents a removable group, for example benzyl. In a typical synthesis, after formation of **20**, group "E" is either left "as is" or converted at some subsequent stage to a derivative thereof. In the example depicted, E is converted to B', a precursor to B in accord with formula **XXI**, to make **21**. For example if E is a nitro, then B' could be an amino group, made via reduction of the nitro group. Structure **21** may be further derivitized by synthesis of -B-L-T in accord with formula **XXI**. In scheme 4, this is depicted as a serial process whereby L', a precursor to L, is introduced to give **22**, followed by introduction of T' (a precursor to T) to give **23**. In some cases, -L-T is preformed and appended to B. One of ordinary skill in the art would appreciate that variations on any of the above steps are possible. Compound **23** is converted to **XXIII** via conversion of T' to T and introduction of P² (for example, when R is benzyl, removal of the benzyl after completion of -B-L-T).

Scheme 4



[0286] As discussed above, one aspect of the invention encompasses combination of **XXII** and **XXIII** to make compounds of formula **XXI**. Because of the diversity and complexity of compounds described for kinase modulation (*vide supra*), methods of the invention provide advantages to serial synthesis.




Examples

[0287] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. All references cited herein are incorporated by reference in their entirety. Generally, but not necessarily, each example set out below describes a multi-step synthesis as outlined above.

Quinoline and Quinazoline Syntheses

Example 1




[0288] Synthesis of 1-(4-Benzyloxy-5-methoxy-2-nitro-phenyl)-ethanone. 1-(4-Benzyloxy-3-methoxy-phenyl)-ethanone (200 mmol, 51.3 g) dissolved in DCM (750ml) and the mixture cooled to 0° C. Nitric acid (90%, 300 mmol, 14 ml) was added dropwise to the cooled solution over 20 minutes. Sulfuric acid (96.2%, 300 mmol, 8.75 ml) was then added dropwise over 40 minutes at 0° C.

[0289] Additional nitric acid (200 mmol, 9.4 ml) was added dropwise over 20 minutes. The reaction mixture was diluted with water (300 ml) and wash with water (3 X 200 ml), Sat. NaHCO₃ (4 X 200 ml, or until neutral). The organic layer was dried over Na₂SO₄ and concentrated.

[0290] The crude mixture was recrystallized with DMF to give 22.5 g of the nitro product. The DMF layer was concentrated and recrystallized with ethyl acetate to give additional 8.75g of the product. The ethyl acetate layer was concentrated and purified on silica column using 20% EtOAc/hexanes to gave another 4.75 g of the product. Total yield is 36 g, (~60%). ¹H NMR (CDCl₃): 7.647 (1H, s), 7.446-7.333 (5H, m), 6.745 (1H, s), 5.210 (2H, s), 3.968 (3H, s), 2.487 (3H, s).


Example 2



[0291] Synthesis of 1-(2-Amino-4-benzyloxy-5-methoxy-phenyl)-ethanone. A Mixture of iron powder (477 mmol, 27 g), ammonium acetate (500 mmol, 31.g), 1-(4-Benzyloxy-5-methoxy-2-nitro-phenyl)-ethanone (120 mmol, 36 g), toluene (500 ml) and water (500 ml) was refluxed overnight, or until completion. The mixture was filtered through celite and


washed with EtOAc. The organic layer was washed with water and Sat. NaCl, dried over Na₂SO₄, and concentrated to afford the product, 90%. ¹H NMR (CDCl₃): 7.408-7.298 (5H, m), 7.130 (1H, s), 6.155 (2H, br), 6.104 (1H, s), 5.134 (2H, s), 3.834 (3H, s), 2.507 (3H, s). LC/MS (M+1 = 272).

Example 3



[0292] Synthesis of 7-Benzyloxy-6-methoxy-quinolin-4-ol. To a solution of 1-(2-Amino-4-benzyloxy-5-methoxy-phenyl)-ethanone (108 mmol, 29.3 g) in DME (700 ml) was added sodium methoxide (432 mmol, 23.35 g). The mixture was stirred for 30 minutes. Ethyl formate (540 mmol, 44 ml) was added and the mixture was stirred overnight. (Additional sodium methoxide may be needed if reaction is not complete as monitored by LC/MS.) After the reaction was completion, the mixture was diluted with water (40 ml) and acidified to neutral with 1M HCl. The precipitate was filtered and washed with water, dried *in vacuo* to afford 22g (72%) of 7-benzyloxy-6-methoxy-quinolin-4-ol. ¹H NMR (CDCl₃): 10.7 (1H, br), 7.703 (1H, s), 7.493-7.461 (1H, t), 7.431-7.413 (2H, br d), 7.372-7.333 (2H, t), 7.296-7.283 (1H, d), 6.839 (1H, s), 6.212-6.193 (1H, d), 5.212 (2H, s), 3.965 (3H, s). LC/MS (M+1 = 282).


Example 4



[0293] 7-Benzyloxy-4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-quinoline. To a round bottom flask equipped with a magnetic stir bar was added 7-Benzyloxy-6-methoxy-1H-quinolin-4-one (12.2 g, 43.3 mmol, 1.0 eq.), acetonitrile (150ml), DMF (150ml) and


cesium carbonate (28.2 g, 86.5 mmol, 2.0 eq). The mixture was stirred at room temperature for 30 minutes at which time 1,2-difluoro-4-nitro-benzene (7.57 g, 47.6 mmol, 1.1 eq) was added over a 10 minute period. After 2 hours the reaction was complete at which time 75% of the MeCN and DMF was removed and the resulting solution was poured over into ice water. The solid was filtered and dried and further columned with a biotage system. The eluent was 1:3 ethyl acetate/hexane. Removal of the solvent afforded 7-Benzyloxy-4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-quinoline as a pale green solid (7.4 g, 41% yield). ^1H NMR (400 MHz, CDCl_3): 8.53 (d, 1H), 8.42 (dd, 1H), 8.16 (m, 1H), 7.5 (m, 8H), 6.76 (d, 1H), 5.31 (s, 2H), 3.92 (s, 3H); MS (EI) for $\text{C}_{23}\text{H}_{27}\text{FN}_2\text{O}_5$: 421 (MH^+).

Example 5




[0294] 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-ol. To a round bottom flask equipped with a magnetic stir bar was added 7-benzyloxy-4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-quinoline (2.9 g, 6.9 mmol, 1.0eq) and 33% HBr in acetic acid (30 ml). The mixture was stirred at room temperature for 3 hours and diluted with ether to give a pale white solid. The solid was filtered, washed with ether and dried to yield 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-ol as a pale white solid (2.74 g, 97.5% yield). ^1H NMR (400 MHz, CDCl_3): 11.89 (bs, 1H), 8.87 (d, 1H), 8.57 (d, 1H), 8.30 (d, 1H), 7.89 (m, 1H), 7.73 (s, 1H), 7.55 (s, 1H), 4.03 (s, 3H); MS (EI) for $\text{C}_{16}\text{H}_{11}\text{FN}_2\text{O}_5$: 421 ($\text{M}+\text{H}^+$).

Example 6



[0295] 5-[4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-yloxymethyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester. To a round bottom flask equipped with a magnetic stir bar was added 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-ol (2.74 g, 6.7 mmol, 1.0 eq.), DMA (30ml) and cesium carbonate (6.6 g, 20.2 mmol, 3.0 eq.). The mixture was stirred at room temperature for 30 minutes at which time 5-methanesulfonyloxymethyl-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester (2.6 g, 7.3 mmol, 1.1 eq) was added. The reaction was heated to 75⁰ C and allowed to stir overnight. After allowing the reaction to cool to room temperature the reaction was poured into water. The solid was filtered and was then dissolved in EtOAc and washed 2X water, 1X brine and dried over NaSO₄. The solvent was removed to yield 5-[4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-yloxymethyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester as a cream solid (3.7 g, 94% yield). ¹H NMR (400 MHz, CDCl₃): 8.55 (d, 1H), 8.15 (d, 1H), 8.09 (d, 1H), 7.32 (m, 8H), 6.52 (d, 1H), 5.11 (d, 2H), 4.13 (d, 2H), 3.95 (s, 3H), 3.57 (m, 2H), 3.43 (m, 2H), 2.93 (m, 3H), 2.16 (m, 2H), 1.39 (m, 2H); MS (EI) for C₃₂H₃₀FN₃O₇: 588 (M+H⁺).


Example 7



[0296] 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinoline. To a round bottom flask equipped with a magnetic stir bar was added 5-[4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-yloxymethyl]-hexahydrocyclopenta-[c]pyrrole-2-carboxylic acid benzyl ester (2.5 g, 4.1 mmol, 1.0eq), 33% HBr in acetic acid (5 ml) and acetic acid (5 ml). The mixture was stirred at room temperature for 1 hour and diluted with EtOAc to give a pale orange solid. The solid was filtered, washed with EtOAc and dried, giving 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinoline (2.1 g, 95% yield). ¹H NMR (400 MHz, CDCl₃): 8.83 (d, 1H), 8.32 (m, 2H), 8.02 (s, 1H), 7.76 (t, 1H), 7.65 (s, 1H), 6.89 (d,


1H), 5.3 (d, 2H), 4.11 (m, 3H), 3.26 (m, 4H), 2.95 (m, 2H), 2.68 (m, 3H), 2.36 (m, 2H), 1.68 (m, 2H); MS (EI) for $C_{24}H_{24}FN_3O_5$: 454 ($M+H^+$).

Example 8




[0297] 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinoline. To a round bottom flask equipped with a magnetic stir bar was added 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinoline (2.1 g, 3.9 mmol, 1.0 eq.) and acetonitrile/water 1:1 (5ml, 5ml). The reaction mixture was then cooled to 0°C and 37% solution of formaldehyde in water was added (0.2 g, 7.8 mmol, 2.0 eq). While keeping the temperature at 0°C $Na(OAc)_3BH$ was added (4.4g, 20.7 mmol, 3.0 eq). After 1 hour the pH was adjusted to 10 and the aqueous was extracted 2 x DCM (100 ml). Removal of the DCM resulted in a white solid. The compound was further purified with a biotage system using an eluent EtOAc and 5% MeOH, affording 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-7-(2-methyl-octahydrocyclopenta-[c]pyrrol-5-ylmethoxy)-quinoline (0.9 g, 50% yield). 1H NMR (400 MHz, $CDCl_3$): 8.57 (d, 1H), 8.14 (dd, 1H), 8.12 (dd, 1H), 7.41 (s, 2H), 7.34 (t, 1H), 6.54 (d, 1H), 4.19 (d, 2H), 4.01 (s, 3H), 2.61 (m, 4H), 2.43 (m, 1H), 2.33 (s, 3H), 2.11 (m, 4H), 1.32 (m, 2H); MS (EI) for $C_{25}H_{26}FN_3O_5$: 468 ($M+H^+$).

Example 9



[0298] 3-Fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinolin-4-yloxy]-phenylamine. To a par hydrogenation reaction vessel was added 4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinoline (0.800 g, 1.6 mmol, 1.0 eq.), DMF (50 ml), EtoAc (50ml), MeOH (50ml), TEA (5ml) and 10% Pd/C (200 mg). The vessel was placed on the par hydrogenator at 35 psi overnight. The Pd was filtered and the solvent removed to give 3-fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinolin-4-yloxy]-phenylamine as an off yellow solid (0.78 g, 99% yield). ¹H NMR (400 MHz, CDCl₃): 8.45 (d, 1H), 7.57 (s, 1H), 7.36 (s, 1H), 7.05 (t, 1H), 6.54 (m, 2H), 6.39 (d, 1H), 4.16 (d, 2H), 4.01 (s, 3H), 3.81 (m, 3H), 2.61 (m, 3H), 2.41 (m, 1H), 2.29 (s, 3H), 2.23 (m, 2H), 1.32 (m, 2H); MS (EI) for C₂₅H₂₈FN₃O₃: 438 (M+H⁺).


Example 10



[0299] 1-[3-Fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinolin-4-yloxy]-phenyl]-3-phenylacetyl-thiourea. To a round bottom flask equipped with a magnetic stir bar was added 3-fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinolin-4-yloxy]-phenylamine (0.78 mg, 1.7 mmol, 1.0 eq.), toluene (10ml), ethanol (10ml) and phenyl-acetyl isothiocyanate (1.64 g, 9.2 mmol, 4.5 eq). The reaction mixture was stirred at room temperature overnight. After removal of the solvent the product was purified with a biotage system using an eluent EtOAc and 4% TEA (2L) then EtOAc, 4% TEA, 1% MeOH (1L). The solvent was removed to give 1-[3-fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinolin-4-yloxy]-phenyl]-3-phenylacetyl-thiourea (0.5 g, 50% yield). ¹H NMR (400 MHz, DMSO): 8.48 (d, 1H), 7.92 (dd, 1H), 7.53 (s, 1H), 7.40 (m, 4H), 7.33 (d, 2H), 7.23 (m, 2H), 6.54 (d, 2H), 6.39 (d, 1H), 4.21 (d, 2H), 4.02 (s, 3H), 3.81 (m, 3H),


2.87 (d, 2H), 2.73 (m, 4H), 2.53 (m, 1H), 2.27 (m, 2H), 2.01 (s, 3H), 1.36 (m, 2H); MS (EI) for $C_{34}H_{35}FN_4O_4S$: 615 ($M+H^+$).

Example 11



[0300] 6-(6,7-Dimethoxy-quinolin-4-yloxy)-5-fluoro-benzothiazol-2-ylamine. 4-(6,7-dimethoxy-quinolin-4-yloxy)-3-fluoro-phenylamine (1.00g, 3.18mmol) was dissolved in AcOH (8.0ml), to which was added NH_4SCN (486mg, 6.38mmol) and the mixture cooled in an ice bath. Br_2 (0.33ml, 6.42mmol) in AcOH (0.33ml) was added dropwise with stirring. After addition was complete, the reaction mixture was stirred at room temperature. After one hour, more NH_4SCN (1.0g, 13.1mmol) was added, followed by more Br_2 (0.33ml, 6.42mmol) in AcOH (0.33ml), dropwise with stirring. The reaction mixture was then heated to reflux for several minutes. Upon cooling to room temperature, solids were filtered and washed with AcOH, followed by H_2O . The volume of the filtrate was reduced *in vacuo* and the pH adjusted to pH 9-10 with 1.0N NaOH. The resulting solids were filtered, washed with H_2O , and dried under high vacuum to give 6-(6,7-dimethoxy-quinolin-4-yloxy)-5-fluoro-benzothiazol-2-ylamine (568mg, 48%). 1H -NMR (400MHz, DMSO): 8.45 (d, 1H), 7.82 (d, 1H), 7.73 (br s, 2H), 7.53 (s, 1H), 7.38 (m, 2H), 6.44 (d, 1H), 3.94 (s, 6H). LC/MS Calcd for $[M+H]^+$ 372.1, found 372.2


Example 12



[0301] N-[6-(6,7-Dimethoxy-quinolin-4-yloxy)-5-fluoro-benzothiazol-2-yl]-2-phenyl-acetamide. 6-(6,7-dimethoxy-quinolin-4-yloxy)-5-fluoro-benzothiazol-2-ylamine (95mg,


0.25mmol), Et₃N (0.10ml, 0.72mmol), phenylacetyl chloride (0.044ml, 0.33mmol), and THF (1.0ml) were combined and stirred at room temperature for 1 hr. Additional phenylacetyl chloride (0.044ml, 0.33mmol) was added and the mixture heated to reflux for 1-2 hrs. After cooling to room temperature, the reaction mixture was diluted with 1:1 AcCN:H₂O (1.0ml) and the resulting solids filtered, washed with 1:1 AcCN:H₂O and dried under high vacuum to give N-[6-(6,7-dimethoxy-quinolin-4-yloxy)-5-fluoro-benzothiazol-2-yl]-2-phenyl-acetamide (72mgs, 59%). ¹H-NMR (400MHz, DMSO): 12.80 (s, 1H), 8.54 (d, 1H), 8.18 (d, 1H), 7.91 (d, 1H), 7.60 (s, 1H), 7.45 (s, 1H), 7.34 (m, 4H), 7.28 (m, 1H), 6.60 (d, 1H), 3.98 (s, 3H), 3.96 (s, 3H), 3.86 (s, 2H). LC/MS Calcd for [M+H]⁺ 490.1, found 490.0.

Example 13




[0302] 5-[4-(4-Amino-2-fluoro-phenoxy)-6-methoxy-quinazolin-7-yloxymethyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester. 4-Amino-2-fluoro-phenol (1.53g, 12.0mmol) was dissolved in dry DMF (30ml) to which was added 60% NaH (774mg, 19.3mmol). After the mixture was stirred at room temperature for several minutes, a suspension of 5-(4-chloro-6-methoxy-quinazolin-7-yloxymethyl)-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester (4.70g, 6.7mmol) in dry DMF (40ml) was added. The reaction mixture was stirred at room temperature for 1-2 hrs, then diluted with EtOAc and washed with sat'd NaHCO₃ (3x), H₂O (1x), sat'd NaCl (1x), dried (Na₂SO₄), and concentrated *in vacuo* to give crude 5-[4-(4-amino-2-fluoro-phenoxy)-6-methoxy-quinazolin-7-yloxymethyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester (5.6g, ~100%) which was used in the next reaction without further purification. ¹H-NMR (400MHz, DMSO): 8.50 (s, 1H), 7.48 (s, 1H), 7.34 (m, 5H), 7.28 (m, 1H), 7.02 (t, 1H), 6.48 (dd, 1H), 6.40 (dd, 1H), 5.40 (br s, 2H), 5.05 (s, 2H), 4.16 (d, 2H), 3.92 (s, 3H), 3.48 (m, 2H), 3.30 (m, 2H), 2.65 (m, 2H), 2.52 (m, 1H), 2.10 (m, 2H), 1.30 (m, 2H). LC/MS Calcd for [M+H]⁺ 559.2, found 559.4.

Example 14




[0303] 5-[4-[2-Fluoro-4-(3-phenylacetyl-thioureido)-phenoxy]-6-methoxy-quinazolin-7-yloxymethyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester. Phenylacetyl chloride (2.65ml, 20.0mmol) and AgSCN (4.92g, 29.6mmol) were combined in dry toluene (50ml) and heated to reflux for 2 hrs. The reaction mixture was allowed to cool to room temperature, the solids were filtered through celite and the filtrate concentrated *in vacuo*. The resulting oil was combined with 5-[4-(4-amino-2-fluorophenoxy)-6-methoxy-quinazolin-7-yloxymethyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester (5.6g, 10mmol) in 1:1 EtOH:toluene (100ml) and the mixture stirred at room temperature for 1-2 hrs. The reaction mixture was diluted with EtOAc and washed with sat'd NaHCO₃ (3x), H₂O (1x), sat'd NaCl (1x), dried (Na₂SO₄), and concentrated *in vacuo*. The resulting oil was purified by flash chromatography (3:1 EtOAc:hexanes) to give 5-[4-[2-fluoro-4-(3-phenylacetyl-thioureido)-phenoxy]-6-methoxy-quinazolin-7-yloxymethyl]-hexahydrocyclopenta[c]pyrrole-2-carboxylic acid benzyl ester (3.61g, 49%) as a dark brown foam. ¹H-NMR (400MHz, DMSO): 12.44 (s, 1H), 11.80 (s, 1H), 8.54 (s, 1H), 7.90 (m, 1H), 7.53 (s, 1H), 7.48 (m, 2H), 7.38 (s, 1H), 7.34 (m, 7H), 7.28 (m, 3H), 5.05 (s, 2H), 4.16 (d, 2H), 3.94 (s, 3H), 3.72 (s, 2H), 3.48 (m, 2H), 3.30 (m, 2H), 2.65 (m, 2H), 2.52 (m, 1H), 2.10 (m, 2H), 1.30 (m, 2H). LC/MS Calcd for [M+H]⁺ 736.2, found 736.0.

Example 15



[0304] 1-{3-Fluoro-4-[6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-3-phenylacetyl-thiourea, dihydrobromide salt. 5-{4-[2-Fluoro-4-(3-phenylacetyl-thioureido)-phenoxy]-6-methoxy-quinazolin-7-yloxymethyl}-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid benzyl ester (3.3g, 4.5mmol) was dissolved in AcOH (70ml) to which was added 33% HBr in AcOH (12ml). The reaction mixture was stirred at room temperature for 1 hr, diluted with Et₂O (1000ml) and the resulting solids filtered, washed with Et₂O, and dried under high vacuum to give the 1-{3-fluoro-4-[6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-3-phenylacetyl-thiourea, dihydrobromide salt (3.4g, 100%). ¹H-NMR (400MHz, DMSO): 12.42 (s, 1H), 11.80 (s, 1H), 8.84 (br s, 2H), 8.64 (s, 1H), 7.92 (m, 1H), 7.59 (s, 1H), 7.49 (m, 2H), 7.41 (s, 1H), 7.33 (m, 4H), 7.27 (m, 1H), 4.17 (d, 2H), 3.95 (s, 3H), 3.73 (s, 2H), 3.17 (m, 2H), 3.10 (m, 2H), 2.83 (m, 2H), 2.45 (m, 1H), 2.15 (m, 2H), 1.30 (m, 2H). LC/MS Calcd for [M+H]⁺ 602.2, found 602.1.


Example 16



[0305] 1-{3-Fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-3-phenylacetyl-thiourea. 1-{3-Fluoro-4-[6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-3-phenylacetyl-thiourea, dihydrobromide salt (3.4g, 4.5mmol) was dissolved in a combination of AcCN (100ml), H₂O (30ml), and AcOH (2.45ml). Formaldehyde (37% in H₂O, 855ml, 10.5mmol) was added and the mixture cooled in an ice bath. Na(OAc)₃BH (2.99g, 14.1mmol) was added and the reaction mixture was stirred at 0 C for 1 hr, followed by stirring at room temperature for 2 hrs. The reaction mixture was neutralized with the addition of sat'd NaHCO₃ and then concentrated *in vacuo*. The resulting aqueous mixture was extracted with CH₂Cl₂ (3x). The combined extractions were washed with sat'd NaHCO₃ (1x), sat'd NaCl (1x), dried (Na₂SO₄), and concentrated *in vacuo*. The

resulting residue was purified by flash chromatography (100% EtOAc, followed by 4% Et₃N in EtOAc) to give the free base of 1-{3-fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-3-phenylacetyl-thiourea (1.13g, 40%). The free base was converted to the HCl salt by dissolving the free base in a mixture of 1:1 AcCN:H₂O containing 2-3 equivalents of 1 N HCl and lyophilizing to give the HCl salt of 1-{3-fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-3-phenylacetyl-thiourea as a white solid. ¹H-NMR (400MHz, DMSO): 12.44 (s, 1H), 11.83 (s, 1H), 10.24 (br s, 1H), 8.59 (s, 1H), 7.93 (m, 1H), 7.59 (s, 1H), 7.50 (m, 2H), 7.42 (s, 1H), 7.36 (m, 4H), 7.30 (m, 1H), 4.20 (m, 2H), 3.95 (s, 3H), 3.73 (s, 2H), 3.39 (m, 2H), 3.06 (m, 2H), 2.95-2.77 (m, 5H), 2.35 (m, 1H), 2.15 (m, 2H), 1.45 (m, 2H). LC/MS Calcd for [M+H]⁺ 616.2, found 616.2. Alternatively, the free base was converted to the acetate salt by dissolving the free base in a mixture of MeOH and CH₂Cl₂ to which was added 3 equivalents of acetic acid. The resulting mixture was concentrated *in vacuo* and the resulting residue lyophilized from 1:1 AcCN:H₂O to give the acetate salt of 1-{3-fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-3-phenylacetyl-thiourea as a white solid. ¹H-NMR (400MHz, CDCl₃): d 12.45 (s, 1H), 8.65 (s, 1H), 7.98 (dd, 1H), 7.50 (s, 1H), 7.40 (m, 4H), 7.29 (m, 4H), 4.17 (d, 2H), 4.05 (s, 3H), 3.75 (s, 2H), 2.93 (m, 2H), 2.80 (m, 2H), 2.72 (m, 2H), 2.53 (s, 3H), 2.47 (m, 1H), 2.25 (m, 2H), 2.02 (s, 3H), 1.35 (m, 2H). LC/MS Calcd for [M+H]⁺ 616.2, found 616.2.


Example 17



[0306] (6,7-Dimethoxy-quinazolin-4-yl)-(2-fluoro-4-nitro-phenyl)-amine. A mixture of 4-chloro-6,7-dimethoxy-quinazoline (548mg, 2.4mmol), 2-fluoro-4-nitro-phenylamine (392mg, 2.5mmol), AcCN (10ml), and conc'd HCl (0.050ml) was heated to reflux for several hrs. After the reaction mixture was allowed to cool to room temperature, the resulting solids were filtered, washed with AcCN and air-dried to give (6,7-dimethoxy-quinazolin-4-yl)-(2-fluoro-4-nitro-phenyl)-amine (673mgs, 80%). ¹H-NMR (400MHz,

DMSO): 12.18 (br s, 1H), 8.91 (s, 1H), 8.45 (s, 1H), 8.36 (dd, 1H), 8.24 (dd, 1H), 7.91 (dd, 1H), 7.44 (s, 1H), 4.04 (s, 3H), 4.02 (s, 3H). LC/MS Calcd for $[M+H]^+$ 345.1, found 345.4.


Example 18



[0307] N¹-(6,7-Dimethoxy-quinazolin-4-yl)-(2-fluoro-benzene-1,4-diamine. (6,7-

Dimethoxy-quinazolin-4-yl)-(2-fluoro-4-nitro-phenyl)-amine (673mg, 1.95mmol) was dissolved in a combination of DMF (20ml) and MeOH (20ml), to which was added 10% Pd/C (227mg). The mixture was shaken under an atmosphere of H₂ on a Parr hydrogenator at 40psi for 3hrs. The reaction mixture was filtered through celite and the filtrate concentrated *in vacuo*. The resulting residue was triturated in EtOAc/Et₂O. The resulting solids were filtered, washed with Et₂O, and dried under vacuum to give N¹-(6,7-dimethoxy-quinazolin-4-yl)-(2-fluoro-benzene-1,4-diamine (398mg, 65%) which was used in the next reaction without further purification. ¹H-NMR (400MHz, DMSO): 10.80 (br s, 1H), 10.30 (br s, 1H), 8.63 (s, 1H), 8.15 (s, 1H), 7.33 (s, 1H), 7.15 (m, 1H), 6.45 (m, 1H), 3.96 (s, 6H). LC/MS Calcd for $[M+H]^+$ 315.1, found 315.4.


Example 19



[0308] 1-[4-(6,7-Dimethoxy-quinazolin-4-ylamino)-3-fluoro-phenyl]-3-phenylacetylthiourea. Phenylacetyl chloride (0.18ml, 1.4mmol) and AgSCN (338mg, 2.0mmol) were

combined in dry toluene (5ml) and heated to reflux for 2 hrs. The reaction mixture was allowed to cool to room temperature, the solids were filtered through celite and the filtrate concentrated *in vacuo*. The resulting oil was combined with N¹-(6,7-Dimethoxy-quinazolin-4-yl)-2-fluoro-benzene-1,4-diamine (398mg, 1.3mmol) in 1:1:2 EtOH:toluene:MeOH (30ml) and the mixture stirred at room temperature overnight. The resulting solids were filtered and washed with toluene, followed by hexanes. The solids were dissolved/suspended in a mixture of EtOAc/MeOH. Insoluble material was filtered and the filtrate concentrated *in vacuo*. The resulting solids were once again dissolved/suspended in a mixture of EtOAc/MeOH. Insoluble material was filtered and the filtrate concentrated *in vacuo* to give 1-[4-(6,7-dimethoxy-quinazolin-4-ylamino)-3-fluoro-phenyl]-3-phenylacetyl-thiourea (105mg, 17%). ¹H-NMR (400MHz, DMSO): 12.53 (s, 1H), 11.86 (s, 1H), 11.44 (br s, 1H), 8.81(s, 1H), 8.25 (s, 1H), 7.94 (dd, 1H), 7.54 (m, 2H), 7.16 (m, 5H), 7.10 (m, 1H), 4.02 (s, 6H), 3.84 (s, 2H). LC/MS Calcd for [M+H]⁺ 492.1, found 492.4.


Example 20



[0309] 6,7-Dimethoxy-4-(5-nitro-pyridin-2-yloxy)-quinoline. To a round bottom flask equipped with a magnetic stir bar was added 6,7-dimethoxy-1H-quinolin-4-one (1.8 g, 8.77 mmol, 1.0 eq.), anhydrous acetonitrile (90 mL) and Cs₂CO₃ (3.13 g, 9.65 mmole, 1.1 eq.). The reaction mixture was stirred at room temperature for 5 minutes. Then, 2-Cl-5-nitropyridine (1.53 g, 9.65 mmol, 1.1 eq.) was added. The reaction mixture was stirred at room temperature for 16 hours. The solids were then filtered off and the filtrate was concentrated via rotary evaporation. The resulting material was taken up in EtOAc, and again the solids were filtered off. The EtOAc filtrate was concentrated. Purification was done on Biotage with solvent system EtOAc 100%. The collected pure fractions were concentrated and dried on high vacuum overnight to give 6,7-dimethoxy-4-(5-nitro-pyridin-2-yloxy)-quinoline as a yellow foam solid (0.902 g, 31.4% yield). ¹H NMR (400


MHz, CDCl₃): 9.08 (d, 1H), 8.74 (d, 1H), 8.60 (dd, 1H), 7.49 (s, 1H), 7.26 (d, 1H), 7.16 (s, 1H), 7.07 (d, 1H), 4.06 (s, 3H), 3.95 (s, 3H); MS (EI) for C₁₆H₁₃N₃O₅: 328 (M+H⁺).

Example 21



[0310] 6-(6,7-Dimethoxy-quinolin-4-yloxy)-pyridin-3-ylamine. To a round bottom flask equipped with a magnetic stir bar was added 6,7-dimethoxy-4-(5-nitro-pyridin-2-yloxy)-quinoline (0.46 g, 1.41 mmol, 1.0 eq.), and THF (10 mL), MeOH (4 mL), DMF (2 mL), and TEA (2 mL). The 6,7-Dimethoxy-4-(5-nitro-pyridin-2-yloxy)-quinoline was dissolved completely in the above solution mixture, and was flushed with nitrogen for at least 5 minutes. The Pd/C (10% by weight) (0.090 g, 20% by weight) was then added. A balloon filled with H₂ was connected to the flask after the nitrogen was vacuumed out. The reaction mixture was stirred at room temperature for 4 hours. The palladium was filtered out through Celite, and the filtrate was collected and concentrated via rotary evaporation. The resulting oil-like product was taken up into 5 mL of water and 1 mL of acetonitrile and lyophilized to yield 6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-ylamine as a light brown solid (0.411 g, 98.1%). ¹H NMR (400 MHz, CDCl₃): 8.54 (d, 1H), 7.85 (d, 1H), 7.53 (s, 1H), 7.41 (s, 1H), 7.18 (dd, 1H), 6.96 (d, 1H), 6.61 (d, 1H), 4.05 (s, 3H), 4.03 (s, 3H), 3.73 (s, 2H); MS (EI) for C₁₆H₁₅N₃O₃: 298 (M+H⁺).


Example 22



[0311] 1-[6-(6,7-Dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-3-phenylacetylthiourea. To a round bottom flask equipped with a magnetic stir bar was added 6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-ylamine (85 mg, 0.0285 mmol, 1.0 eq.), and Phenyl-acetyl

isothiocyanate (256 mg, 1.44 mmol, 5.0 eq.) dissolved in EtOAc/MeOH 50:50 (2 mL). The reaction mixture was stirred at room temperature for 12 hours, and the solvent was evaporated via rotary evaporation. Purification was done on Biotage with solvent system 95% EtOAc, 4% TEA and 1% MeOH. The combined pure fractions were concentrated and dried under vacuum overnight to yield 1-[6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-3-phenylacetyl-thiourea as a light yellow solid (40.4 mg, 29.7%). ^1H NMR (400 MHz, CDCl_3): 8.65 (d, 1H), 8.33 (d, 1H), 8.27 (dd, 1H), 7.35 (m, 7H), 7.15 (d, 1H), 6.92 (d, 1H), 4.05 (s, 3H), 3.99 (s, 3H), 3.76 (s, 2H); MS (EI) for $\text{C}_{25}\text{H}_{22}\text{N}_4\text{O}_4\text{S}$: 475 ($\text{M}+\text{H}^+$).


Example 23



[0312] N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-N'-phenethyl-oxalamide.

To a solution of 4-(6,7-dimethoxy-quinolin-4-yloxy)-3-fluorophenylamine (263 mg, 0.83 mmol) and Et_3N (0.223 ml, 1.67 mmol) in CH_2Cl_2 (10 mL) was added dropwise a solution of ethyl oxalyl chloride in CH_2Cl_2 (1 mL). The stirring was continued for 0.5 h at rt. The reaction mixture was then washed with aqueous saturated NaHCO_3 and dried over NaSO_4 . Removal of the solvent gave the crude oxamate, which was treated with neat phenethylamine (1.0 g, 8.3 mmol) at 80°C for 3 h. Purification by flash column chromatography (hexanes:EtOAc = 1:3) gave N-[4-(6,7-dimethoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-N'-phenethyl-oxalamide (310 mg, 76%). ^1H NMR (400 MHz, CDCl_3) δ 9.35 (br s, 1 H), 8.70 (d, $J = 6.3$ Hz, 1 H), 7.83 (dd, $J = 11.9, 2.5$ Hz, 1 H), 7.60-7.54 (m, 2 H), 7.43 (s, 1 H), 7.38-7.32 (m, 3 H), 7.30-7.20 (m, 4 H), 6.41 (d, $J = 5.3$ Hz, 1 H), 4.07 (s, 3 H), 4.05 (s, 3 H), 3.67 (dt, $J = 7.0, 7.0$ Hz, 2 H), 2.92 (t, $J = 7.2$ Hz, 2 H). LC-MS: 490 [$\text{M}+\text{H}$] $^+$

Example 24




- [0313] N-(3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-N'-phenethyl-oxalamide. To a flask containing 7-benzyloxy-4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-quinoline (850 mg, 2.0 mmol) was added 20 mL of 30% HBr in AcOH. The resulted solution was stirred for 4 h at rt; at this time, a large amount of precipitate formed. The crude product was filtered, washed with Et₂O and dried in air, giving 4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-7-hydroxyquinoline (609 mg, 92% yield).
- [0314] To a solution of the 4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-7-hydroxyquinoline (609 mg, 1.8 mmol) in DMF (9 mL) was added K₂CO₃ (1.24 g, 9.0 mmol) and N-Boc-4-piperidinemethanol mesylate (732 mg, 2.5 mmol). The mixture was then stirred at 80 °C for 2.5 h. After it was cooled to rt, the mixture was loaded directly to a Biotage column, and eluted with solvents (hexanes:EtOAc = 1:3). The resulting product, 4-[4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester, was obtained as a solid (556 mg, 56%).
- [0315] To a solution of 4-[4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-quinolin-7-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester (305 mg, 0.58 mmol) in CH₂Cl₂ (1 mL) was added 0.4 mL of TFA. The reaction mixture was stirred for 1.5 h and the solvents were removed under reduced pressure. The crude product was treated with NaBH(OAc)₃ (381 mg, 1.80 mmol) and formaldehyde (0.5 mL, 37% in H₂O). The stirring was continued for 12 h. The reaction was quenched with sat. aqueous NaHCO₃. 15% NaOH was added until PH = 14. The product was extracted with EtOAc. Removal of the solvent *in vacuo* gave the crude product, 4-(2-fluoro-4-nitro-phenoxy)-6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinoline, (240 mg, 93%), which was used directly in the next reaction.
- [0316] To a solution of 4-(2-Fluoro-4-nitro-phenoxy)-6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinoline (240 mg, 0.54 mmol) in EtOH (20 mL) was added 10% Pd/C (50

mg). The mixture was then hydrogenated on a Parr hydrogenator (40 psi) for 10 h. AcOH was added to dissolve the intermediate (mostly the hydroxylamine) and the hydrogenation was continued for additional 12 h. LC-MS was used to monitor the reaction progress. The solvents were removed under reduced pressure and the resulting crude product of 3-fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenylamine (about 220 mg) was used directly in the next reaction.

[0317] To a 0 °C solution of 3-fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenylamine (66 mg, 0.13 mmol) and Et₃N (0.34 mL) in CH₂Cl₂ (6 mL) was added slowly ethyl oxalyl chloride (98 mg). The reaction mixture was stirred at rt for 30 min, then diluted with CH₂Cl₂ and washed with sat. aqueous NaHCO₃. After dried over MgSO₄ and concentrated, the crude ethyl oxamate was reacted with phenethylamine (80 mg, 0.64 mmol) at 80 °C for 2 h. Purification by HPLC gave product, N-{3-fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide (52 mg, 68% yield). ¹H NMR (400 MHz) δ 9.38 (br s, 1 H), 8.48 (d, J = 5.2 Hz, 1 H), 7.83 (dd, J = 11.7, 2.6 Hz, 1 H), 7.59 (t, J = 6.2 Hz, 1 H), 7.55 (s, 1 H), 7.40-7.20 (8 H), 6.39 (d, J = 5.3 Hz, 1 H), 4.06 (d, J = 6.6 Hz, 2 H), 4.04 (s, 3 H), 3.67 (q, J = 6.8 Hz, 2 H), 2.98 (br d, J = 11.5 Hz, 2 H), 2.92 (t, J = 7.0 Hz, 2 H), 2.34 (s, 3 H), 2.10-1.80 (m, 5 H), 1.60-1.54 (m, 2 H).


Example 25



[0318] 1-(4-Fluorophenylcarbamoyl)-cyclopropanecarboxylic acid. The title compound was prepared based on a modified procedure of Shih and Rankin [*Synthetic Communications*, 1996, 26(4), 833-836]: To a mixture of cyclopropane-1,1-dicarboxylic acid (21.2 g, 0.163 mol, 1.0 eq.) in anhydrous THF (200 mL) under nitrogen was added dropwise triethylamine (16.49 g, 0.163 mol, 1.0 eq.) with stirring for 30 minutes at 0°C, followed by the addition of thionyl chloride (19.39 g, 0.163 mol, 1.0 eq.) with stirring for another 30 minutes at 0°C. To the resulting mixture under nitrogen was added dropwise a solution of 4-fluoroaniline (19.92 g, 0.179 mol, 1.1 eq.) in anhydrous THF (100 mL) with stirring for 1.5 hours at 0°C. The reaction mixture was diluted with ethyl acetate and


washed with 1N NaOH. The layers were separated, and the ethyl acetate layer was concentrated in vacuo to give a brownish solid. The brownish solid was washed with small amount of cold ethyl acetate, filtered and dried under vacuum to yield 1-(4-fluorophenylcarbamoyl)-cyclopropanecarboxylic acid as a white solid (23.71 g, 65.18%). ^1H NMR (400 MHz, CD_3OD): 7.57-7.53 (m, 2H), 7.05-7.00 (m, 2H) 1.46-1.43 (m, 2H), 1.40-1.37 (m, 2H).

Example 26



[0319] 1-(4-Fluoro-phenylcarbamoyl)-cyclobutanecarboxylic acid. To a mixture of cyclobutane-1,1-dicarboxylic acid (10.0 g, 69.4 mmol, 1.0 eq.) in anhydrous THF (100 mL) under nitrogen was added dropwise triethylamine (7.02 g, 69.4 mmol, 1.0 eq.) with stirring for 30 minutes at 0°C, followed by the addition of thionyl chloride (8.25 g, 69.4 mmol, 1.0 eq.) with stirring for another 30 minutes at 0°C. To the resulting mixture under nitrogen was added dropwise a solution of 4-fluoroaniline (8.48 g, 76.3 mmol, 1.1 eq.) in anhydrous THF (50 mL) with stirring for 1.5 hours at 0°C. The reaction mixture was diluted with ethyl acetate and extracted with 2N NaOH. The aqueous phase was titrated with 2N HCl to pH 1-2 and then extracted with ethyl acetate. The organic phase was dried with sodium sulfate and concentrated in vacuo to give 1-(4-fluoro-phenylcarbamoyl)-cyclobutanecarboxylic acid as a light pink solid (5.75 g, 34.9%). ^1H NMR (400 MHz, CDCl_3 w/1drop CD_3OD): 7.53-7.48 (m, 2H), 7.06-7.00 (m, 2H), 2.81-2.63 (m, 4H), 2.14-2.02 (m, 2H).


Example 27



[0320] 1-Benzylcarbamoyl-cyclopropanecarboxylic acid. The title compound was prepared based on a modified procedure of Shih and Rankin [*Synthetic Communications*, 1996, 26(4), 833-836]: To a mixture of cyclopropane-1,1-dicarboxylic acid (5.0 g, 38.4


mmol, 1.0 eq.) in anhydrous THF (50 mL) under nitrogen was added dropwise triethylamine (3.89 g, 38.4 mmol, 1.0 eq.) with stirring for 30 minutes at 0°C, followed by the addition of thionyl chloride (4.57 g, 38.4 mmol, 1.0 eq.) with stirring for another 30 minutes at 0°C. To the resulting mixture under nitrogen was added dropwise a solution of benzylamine 5 (4.53 g, 42.3 mmol, 1.1 eq.) in anhydrous THF (25 mL) with stirring for 1.5 hours at 0°C. The reaction mixture was diluted with ethyl acetate and extracted with 2N NaOH (to pH 10). The aqueous phase was titrated with 2N HCl to pH 1-2 and then extracted with ethyl acetate. The organic phase was dried with sodium sulfate and concentrated in vacuo to give 1-Benzylcarbamoyl-cyclopropanecarboxylic acid as a white solid (4.39 g, 52.15%). ¹H NMR (400 MHz, CDCl₃): 8.44 (br s, 1H), 7.37-7.33 (m, 2H), 7.32-7.26 (m, 3H), 1.82-1.70 (m, 4H).

Example 28




[0321] 1-Phenylcarbamoyl-cyclopropanecarboxylic acid. To a mixture of cyclopropane-1,1-dicarboxylic acid (5.29 g, 40.7 mmol, 1.0 eq.) in anhydrous THF (50 mL) under nitrogen was added dropwise triethylamine (4.12 g, 40.7 mmol, 1.0 eq.) with stirring for 30 minutes at 0°C, followed by the addition of thionyl chloride (4.84 g, 40.7 mmol, 1.0 eq.) with stirring for another 30 minutes at 0°C. To the resulting mixture under nitrogen was added dropwise a solution of phenylamine 9 (4.17 g, 44.8 mmol, 1.1 eq.) in anhydrous THF (25 mL) with stirring for 1.5 hours at 0°C. The reaction mixture was diluted with ethyl acetate and extracted with 2N NaOH (to pH >10). The aqueous phase was titrated with 2N HCl to pH 1-2 and then extracted with ethyl acetate. The organic phase was dried with sodium sulfate and concentrated in vacuo to give 1-phenylcarbamoyl-cyclopropanecarboxylic acid as a white solid (5.08 g, 60.8%). ¹H NMR (400 MHz, CDCl₃): 10.50 (br s, 1H), 7.56-7.54 (m, 2H), 7.35-7.31 (m, 2H), 7.15-7.10 (m, 1H), 1.94-1.91 (m, 2H), 1.82-1.79 (m, 2H).

Example 29



[0322] 7-Benzyloxy-4-chloro-6-methoxy-quinoline. Dry DMF (8.0ml, 103mmol) was dissolved in dry CHCl₃ (40ml) and cooled in an ice bath. Oxalyl chloride (9.0ml, 105mmol) in CH₂Cl₂ (10ml) was added dropwise with stirring at 0°C. When the bubbling had ceased, this solution was added slowly to an ice-cold solution of 7-benzyloxy-6-methoxy-3H-quinazolin-4-one (10.0g, 35.4mmol) in dry CHCl₃ (60ml) and the mixture was then heated to reflux for 2-3hrs. After cooling to room temperature, H₂O (100ml) was added and the phases were separated. The aqueous phase was further extracted with CHCl₃ (2x). The combined CHCl₃ extractions were washed with sat'd NaCl (1x), dried (Na₂SO₄) and concentrated *in vacuo*. The resulting residue was purified by flash chromatography (1:1 hexanes:EtOAc, followed by 100%EtOAc) to give 7-benzyloxy-4-chloro-6-methoxy-quinoline (5.11g, 48%). LC/MS Calcd for [M+H]⁺ 301.1, found 301.1.


Example 30




[0323] Cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinolin-4-yloxy)-phenyl]-amide(4-fluoro-phenyl)-amide. To a solution of cyclopropane-1,1-dicarboxylic acid [4-(7-benzyloxy-6-methoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-amide (4-fluoro-phenyl)-amide (1.18g, 2.0 mmol) in EtOH (20 mL) was added 1,4-cyclohexadiene (2.0 mL, 20 mmol) and 10% Pd/C (300 mg). The reaction mixture was then heated to reflux and the stirring was continued for 2 h. It was cooled to room temperature, filtered through celite and washed with MeOH. The MeOH solution was then concentrated under reduced pressure. The residue was taken into EtOAc (200 mL).

The EtOAc solution was washed with water, and dried over Na₂SO₄. Removal of the solvent under reduced pressure gave 900 mg (89%) of the crude product (90% purity by analytical HPLC), which was used in the next reaction without further purification.

Example 31



[0324] N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methoxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. To a mixture of cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinolin-4-yloxy)-phenyl]-amide(4-fluoro-phenyl)-amide (186 mg, 0.36 mmol) in CH₂Cl₂ (10 mL) was added 2-(diethylamino)ethanol (63 mg, 0.54 mmol), and PPh₃ (141 mg, 0.54 mmol). DIAD (109 mg, 0.54 mmol) was then added as a CH₂Cl₂ (1 mL) solution. The resulted solution was stirred at room temperature for 2 h and the solvent was removed under reduced pressure. To the residue was added 1 N HCl (50 mL), and it was washed with EtOAc (50 mL x 2). The aqueous phase was basified by adding 15% NaOH aqueous solution until pH =11-13, and then extracted with ether (50 mL x 2). The combined organic layer was dried (MgSO₄), and concentrated in vacuo. The residue was purified on preparative HPLC to give N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methoxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluoro-phenyl)cyclopropane-1,1-dicarboxamide (74 mg, 34%) as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.40 (br s, 1 H), 10.02 (br s, 1 H), 8.47 (d, *J* = 5.2 Hz, 1 H), 7.91 (br d, *J* = 13.9 Hz, 1 H), 7.54-7.52 (m, 2 H), 7.55-7.50 (m, 1 H), 7.52 (s, 1 H), 7.50-7.40 (m, 1 H), 7.41 (s, 1 H), 7.16 (br t, *J* = 8.7 Hz, 2 H), 6.41 (br d, *J* = 4.7 Hz, 1 H), 4.18 (t, *J* = 6.0 Hz, 2 H), 3.94 (s, 3 H), 2.87 (br t, *J* = 6.3 Hz, 2 H), 2.59 (q, *J* = 7.1 Hz, 4 H), 1.47 (br s, 4 H), 1.00 (t, *J* = 7.0 Hz, 6 H).

Example 32

[0325] 1-(4-Benzyloxy-3-methoxyphenyl)ethanone. A solution of 4-hydroxy-3-methoxyacetophenone (40 g, 240 mmol), benzyl bromide (31.4 mL, 260 mmol) and potassium carbonate (99.6 g, 360 mmol) in DMF (800 mL) was heated to 40 °C overnight. The solution was cooled to room temperature, poured over ice and the resultant solid was filtered. This material was washed with water and dried to give 1-(4-benzyloxy-3-methoxyphenyl)ethanone (61 g, 99 %).

[0326] 1-(4-Benzyloxy-5-methoxy-2-nitrophenyl)ethanone. A stirred solution of 1-(4-benzyloxy-3-methoxyphenyl)ethanone (51.3 g, 200 mmol) in dichloromethane (750 mL) was cooled to 0 °C. Nitric acid (90 %, 14 mL, 300 mmol) was added dropwise to the cooled solution over 20 min. Sulfuric acid (96.2 %, 16.3 mL, 300 mmol) was then added dropwise over 40 min at 0 °C. Additional nitric acid (9.4 mL, 200 mmol) was added dropwise over 20 min. The reaction mixture was washed with water (3 x 200 mL), and saturated sodium bicarbonate (4 x 200 mL, or until neutral). The organic layer was dried over Na₂SO₄ and concentrated. The crude mixture was recrystallized from DMF to give 1-(4-benzyloxy-5-methoxy-2-nitrophenyl)ethanone (36 g, 60 %). ¹H NMR (400 MHz, CDCl₃): δ 7.65 (s, 1H), 7.45-7.33 (m, 5H), 6.74 (s, 1H), 5.21 (s, 2H), 3.97 (s, 3H), 2.49 (s, 3H).


[0327] 1-(2-Amino-4-benzyloxy-5-methoxyphenyl)ethanone. A mixture of iron powder (27 g, 0.48 g atoms), ammonium formate (31 g, 500 mmol), 1-(4-benzyloxy-5-methoxy-2-nitrophenyl)ethanone (36 g, 120 mmol), toluene (500 mL) and water (500 mL) was heated to reflux overnight. The mixture was filtered through celite and washed with ethyl acetate. The combined organic layers were washed with water and brine. The organic layer was

dried over Na_2SO_4 and concentrated to afford 1-(2-amino-4-benzyloxy-5-methoxyphenyl)ethanone (29.3 g, 90 %). ^1H NMR (CDCl_3): δ 7.41-7.30 (m, 5H), 7.13 (s, 1H), 6.16 (br s, 2H), 6.10 (s, 1H), 5.13 (s, 2H), 3.83 (s, 3H), 2.51 (s, 3H). LC/MS ($\text{M}+\text{H} = 272$).

[0328] 7-Benzyloxy-6-methoxyquinolin-4-ol. Sodium ethoxide (74.8 g, 1.1 mol) was added to a solution of 1-(2-amino-4-benzyloxy-5-methoxyphenyl)ethanone (29.3 g, 108 mmol) in DME (700 mL) and stirred for 30 min. Ethyl formate (44 mL, 540 mmol) was added and the mixture was stirred overnight (in case of incomplete reaction, additional sodium ethoxide can be added and the reaction monitored by LC/MS). After the reaction was complete, the mixture was diluted with water (40 mL) and acidified to neutral pH with 1M HCl. The solid was filtered, washed with water and dried to afford 7-benzyloxy-6-methoxyquinolin-4-ol (22 g, 72%). ^1H NMR (400 MHz, CDCl_3): δ 10.7 (br s, 1H), 7.70 (s, 1H), 7.49-7.46 (t, 1H), 7.43-7.41 (br d, 2H), 7.37-7.33 (t, 2H), 7.30-7.28 (d, 1H), 6.84 (s, 1H), 6.21-6.19 (d, 1H), 5.21 (s, 2H), 3.96 (s, 3H). LC/MS ($\text{M}+\text{H} = 282$).

[0329] 7-Benzyloxy-4-chloro-6-methoxyquinoline. Phosphorus oxychloride (300 mL) was added to 7-benzyloxy-6-methoxyquinolin-4-ol (40 g, 140 mmol) and the mixture heated to reflux for 2 h. The mixture was carefully poured into a mixture of ice and sodium carbonate. The solution was adjusted to pH 8 with the addition of solid sodium bicarbonate and stirred at room temperature overnight. The solid was filtered and washed with water and dried to give 7-benzyloxy-4-chloro-6-methoxyquinoline as a pale brown solid (40.2 g, 95%). ^1H NMR (400 MHz, d_6 -DMSO): δ 8.61 (s, 1H), 7.57-7.37 (m, 8H), 5.32 (s, 2H), 3.98 (s, 3H); ^{13}C NMR (100 MHz, d_6 -DMSO): δ 152.4, 151.5, 148.5, 146.2, 139.6, 137.0, 129.2, 128.8, 121.7, 120.4, 110.1, 101.9, 70.8, 56.5; IR (cm^{-1}): 2359, 2341, 1506, 1456, 1435, 1252, 1227, 1146, 999, 845, 752, 698, 667; LC/MS ($\text{M}+\text{H} = 300$).


Example 33



[0330] Trifluoromethanesulfonic acid 7-benzyloxy-6-methoxyquinolin-4-yl ester. To a dry 2L RBF containing 7-benzyloxy-6-methoxyquinolin-4-ol (75.3 g, 267 mmol) was added DCM (1 L), 4-dimethylaminopyridine (3.28 g, 26.8 mmol) and 2,6-lutidine (62 mL,


534 mmol). The mixture was cooled to -20°C by controlled addition of dry ice to an acetone bath. Trifluoromethanesulfonyl chloride (37 mL, 350 mmol) was added dropwise to the cooled solution with magnetic stirring over 25 minutes. After addition was complete, the mixture was stirred in bath for 20 minutes, then at room temperature for 3 hours. LCMS indicated reaction completion. The reaction mixture was concentrated *in vacuo* and placed under high vacuum to remove residual 2,6-lutidine. To the resulting brown solids was added methanol (3.5 L). The resulting slurry was stirred with mechanical stirrer for 30 min before adding water (1.5 L). The solids were isolated by filtration, followed by a water wash. The resulting solid was dried under high vacuum overnight yielding trifluoromethanesulfonic acid 7-benzyloxy-6-methoxy-quinolin-4-yl ester as a light brown solid (92.2 g, 83.8%). ^1H NMR (400MHz, DMSO, d_6): δ 8.82 (d, 1H), 7.67 (s, 1H), 7.59 (d, 1H), 7.54-7.52 (m, 2H), 7.46-7.42 (m, 2H), 7.39-7.36 (m, 1H), 7.23 (s, 1H), 5.35 (s, 2H), 3.97 (s, 3H). LC/MS: $M+H = 414$.

Example 34



[0331] Trifluoromethanesulfonic acid 6,7-dimethoxyquinolin-4-yl ester from 6,7-Dimethoxy-quinolin-4-ol. To a dry 1L RBF containing 6,7-dimethoxy-quinolin-4-ol (20.9 g, 102 mmol), which can be prepared according to the procedure of Riegel, B. (*J. Amer. Chem. Soc.* 1946, 68, 1264), was added DCM (500 mL), 4-dimethylaminopyridine (1.24 g, 10 mmol) and 2,6-lutidine (24 mL, 204 mmol). The mixture was vigorously stirred at RT. Trifluoromethanesulfonyl chloride (14 mL, 132 mmol) was added dropwise to the solution. After addition was complete, the mixture was stirred ice bath for 2 to 3 hrs. On LC/MS indicating the reaction completion, the reaction mixture was concentrated *in vacuo* and placed under high vacuum to remove residual 2,6-lutidine. To the resulting brown solids was added methanol (250 mL). The resulting slurry was stirred for 30 min before adding water (1 L). The solids were isolated by filtration, followed by a water wash. The resulting solid was dried under high vacuum overnight yielding trifluoromethanesulfonic acid 6, 7-dimethoxy-quinolin-4-yl ester as a light brown solid (27 g, 80%). ^1H NMR (400MHz, DMSO, d_6): δ 8.82 (d, 1H), 7.59 (m, 2H), 7.20 (s, 1H), 3.97 (d, 6H). LC/MS: $M+H = 338$.

Example 35



[0332] 1-Benzoyloxy-2-fluoro-4-nitrobenzene. A solution of 2-fluoro-4-nitrophenol (50.0 g, 318 mmol), benzyl bromide (42 mL, 350 mmol) and potassium carbonate (66.0 g, 478 mmol) in DMF (200 mL) was heated to 40 °C overnight. The solution was cooled to room temperature, poured over ice and the resultant solid was filtered. This material was washed with water and dried to give 1-benzoyloxy-2-fluoro-4-nitrobenzene (75.0 g, 95 %). ¹H NMR (400 MHz, *d*₆-DMSO): δ 8.19-8.11 (m, 2H), 7.53-7.37 (m, 6H), 5.36 (s, 2H); ¹³C NMR (100 MHz, *d*₆-DMSO): δ 152.8, 152.4, 149.9, 140.9, 136.1, 129.3, 129.1, 128.7, 122.0, 115.2, 112.8, 112.6, 71.6; IR (cm⁻¹): 1499, 1346, 1279, 1211, 1142, 1072, 986, 885, 812, 789, 754, 742, 700, 648, 577.

[0333] 4-Benzoyloxy-3-fluoroaniline. A mixture of iron powder (45.2 g, 0.809 g atoms), ammonium formate (53.6 g, 0.850 mol), 1-benzoyloxy-2-fluoro-4-nitrobenzene (50.0 g, 0.200 mol), toluene (400 mL) and water (400 mL) was heated to reflux overnight. The mixture was filtered through Celite and washed with hot ethyl acetate. The combined organic layers were washed with water and brine, then dried over sodium sulfate and concentrated to afford 4-benzoyloxy-3-fluoroaniline (44 g, 100 %). ¹H NMR (400 MHz, *d*₆-DMSO): δ 7.43-7.26 (m, 5H), 6.90 (dd, 1H), 6.49 (dd, 1H), 6.34 (m, 1H), 4.99 (br s, 2H), 4.98 (s, 2H); ¹³C NMR (100 MHz, *d*₆-DMSO): δ 171.1, 155.1, 152.7, 144.9, 138.0, 137.2, 129.6, 129.0, 128.5, 118.9, 110.0, 102.9, 72.5; IR (cm⁻¹): 1510, 1454, 1277, 1215, 1126, 1007, 957, 843, 800, 789, 739, 694, 604; LC/MS (M+H = 218).


[0334] Ethyl [(4-benzoyloxy-3-fluorophenyl)amino](oxo)acetate. Ethyl oxalyl chloride (44 mL, 390 mmol) was added to a solution of 4-benzoyloxy-3-fluoroaniline (44 g, 180 mmol) in diisopropylethylamine (69 mL, 400 mmol) and stirred at room temperature for 15 min. The mixture was extracted with dichloromethane and washed with water and brine. The organic layer was dried over sodium sulfate and concentrated to afford ethyl [(4-

benzyloxy-3-fluorophenyl)amino](oxo)acetate (58.4 g, 100 %). ^1H NMR (400 MHz, d_6 -DMSO): δ 10.87 (s, 1H), 7.73 (d, 1H), 7.69 (d, 1H), 7.53 (d, 1H), 7.46-7.40 (m, 4H), 5.17 (s, 2H), 4.31 (q, 2H), 1.31 (t, 3H); IR (cm^{-1}): 1732, 1705, 1558, 1541, 1508, 1456, 1273, 1186, 1167, 1101, 999, 858, 741, 694; LC/MS ($M+H = 318$).

[0335] *N*-(4-Benzyloxy-3-fluorophenyl)-*N'*-(2-phenylethyl)ethanediamide. Phenethylamine (33 mL, 520 mmol) was added to ethyl [(4-benzyloxy-3-fluorophenyl)amino](oxo)acetate (81 g, 260 mmol) and the mixture was sonicated at room temperature for 30 min. The resulting solid was filtered, washed with water and dried to give *N*-(4-benzyloxy-3-fluorophenyl)-*N'*-(2-phenylethyl)ethanediamide (100 g, 99 %). ^1H NMR (400 MHz, d_6 -DMSO): δ 10.72 (br s, 1H), 9.05 (m, 1H), 8.78 (m, 1H), 7.77 (m, 1H), 7.59 (m, 1H), 7.46-7.19 (m, 8H), 5.16 (m, 2H), 3.45 (m, 2H), 2.83 (m, 2H); IR (cm^{-1}): 2980, 2883, 1653, 1522, 1506, 1441, 1385, 1221, 1122, 951, 808, 746, 696, 584; LC/MS ($M+H = 393$).

[0336] *N*-(3-Fluoro-4-hydroxyphenyl)-*N'*-(2-phenylethyl)ethanediamide. A mixture of *N*-(4-benzyloxy-3-fluorophenyl)-*N'*-(2-phenylethyl)ethanediamide (40 g, 100 mmol) and 38% hydrobromic acid in acetic acid (250 mL) was stirred at room temperature overnight. The resulting solid was filtered, washed with water and dried to give *N*-(3-fluoro-4-hydroxyphenyl)-*N'*-(2-phenylethyl)ethanediamide as a slightly yellow solid (30.6 g, 99 %yield). ^1H NMR (400 MHz, d_6 -DMSO): δ 10.60 (s, 1H), 9.02 (t, 1H), 7.70 (d, 1H), 7.47 (d, 1H), 7.32-7.20 (m, 3H), 6.91 (t, 1H), 3.43 (m, 2H), 2.81 (m, 2H); ^{13}C NMR (100 MHz, d_6 -DMSO): δ 160.5, 158.8, 152.0, 149.6, 142.2, 139.8, 130.3, 129.3, 129.0, 126.8, 118.1, 117.4, 109.6, 109.3 IR (cm^{-1}): 3279, 1653, 1518, 1456, 1279, 1190, 742, 696, 584; LC/MS ($M+H = 303$).

Example 36



[0337] 1-Benzyloxy-2-fluoro-4-nitro-benzene. To a slurry of sodium hydride (60% dispersion in oil, 693 mmol, 27.7 g) and dimethylacetamide (600 ml) was added benzyl alcohol (462 mmol, 48 ml) dropwise with stirring under N₂. The mixture was stirred for 1 hour at RT and then cooled to 0°C. 3,4-difluoronitrobenzene (508 mmol, 56.2 ml) was added to the cooled solution and stirred for 1 hour. Reaction mixture poured onto saturated ammonium chloride solution (800 ml) and stirred for 30 minutes, filtered and washed with water. The solid was stirred in ethyl acetate (500 mL), and filtered to give 54g of product. The ethyl acetate filtrate, after concentrated in vacuo, was triturated with diethyl ether (500 mL), sonicated for 2 hours, and filtered to give another 30g of product. The ether layer was concentrated and column purified using 5% EtOAc/hexanes as eluent to give additional 15g of product. The total yield of 1-benzyloxy-2-fluoro-4-nitrobenzene was 95g (83%). (Note: the product contains ca. 5% of 3,4-Bis-benzyloxy-nitrobenzene, which is carried into the next step without further purification.) ¹H NMR (400MHz, CDCl₃): δ 8.04–8.00 (m, 2H), 7.43–7.37 (m, 5H), 7.08 (t, 1H), 5.26 (s, 2H).


[0338] 4-Benzyloxy-3-fluoro-phenylamine. A mixture of 1-benzyloxy-2-fluoro-4-nitrobenzene (44g, 178 mmol), toluene (400 ml), ammonium formate (35 g), iron (30 g), and water (400 ml) was heated to reflux with stirring overnight. The reaction mixture was filtered through celite and washed with ethyl acetate (400ml). The organic layer was separated and washed with brine (300 ml), dried over sodium sulfate and concentrated to give 4-benzyloxy-3-fluoro-phenylamine as an oil (33.7 g, 87%). ¹H NMR (400MHz, CDCl₃): δ 7.41–7.29 (m, 5H), 6.79 (t, 1H), 6.45 (dd, 1H), 6.14 (dd, 1H), 5.02 (s, 2H), 3.50 (s, 2H). LC/MS: (M+1) 218.

[0339] Cyclopropane-1,1-dicarboxylic acid (4-benzyloxy-3-fluoro-phenyl)-amide (4-fluoro-phenyl)-amide. To a stirred mixture of 4-benzyloxy-3-fluoro-phenylamine (155.3 mmol, 33.7 g), 1-(4-fluorophenylcarbonyl)-cyclopropanecarboxylic acid (170.8 mmol, 38.13 g) and anhydrous dichloromethane (600 ml) was added EDCI (233.9 mmol, 44.7 g) in portions. After stirring at RT for 1 hr, the reaction mixture was diluted with saturated sodium bicarbonate (400 ml) and stirred for 30 minutes. The precipitate was filtered and air dried to give the 1st crop of product. The biphasic filtrate was separated, and the organic phase was washed with brine (300 ml), dried over sodium sulfate, and concentrated. The residue was taken up in DCM (100 ml), stirred for 15 minutes, and filtered to give a 2nd crop of product. The combined yield of cyclopropane-1,1-dicarboxylic acid (4-benzyloxy-3-fluoro-phenyl)-amide (4-fluoro-phenyl)-amide was 64.5

g (98%). ^1H NMR (400MHz, CDCl_3): δ 8.92 (br s, 1H), 8.88 (br s, 1H), 7.50-7.32 (m, 8H), 7.06-7.02 (m, 3H), 6.97-6.92 (t, 1H), 5.13 (s, 2H), 1.65 (s, 4H). LC/MS: (M+1) 423.

[0340] Cyclopropane-1,1-dicarboxylic acid (3-fluoro-4-hydroxy-phenyl)-amide (4-fluoro-phenyl)-amide. A mixture of cyclopropane-1,1-dicarboxylic acid (4-benzyloxy-3-fluoro-phenyl)-amide (4-fluoro-phenyl)-amide (152.8 mmol, 64.5), ethanol (800 ml), cyclohexadiene (764 mmol, 71 ml), and 10% Pd/C (2 g) was refluxed for 2 hours. Reaction mixture cooled and filtered through celite and washed with methanol. The combined filtrate was concentrated and stirred in 10% EtOAc/ether (350 ml). The resulting precipitate was filtered and washed with ether to give a 1st crop of product. The filtrate was concentrated and stirred in DCM (150 ml) to give another precipitate, which was then filtered to give a 2nd crop of product. The combined yield of cyclopropane-1,1-dicarboxylic acid (3-fluoro-4-hydroxy-phenyl)-amide (4-fluoro-phenyl)-amide was 43 g (85%) in 95% purity by HPLC (UV @ 254 nm). ^1H NMR (400MHz, $\text{DMSO}-d_6$): δ 10.07 (br s, 1H), 9.92 (br s, 1H), 9.64 (br s, 1H), 7.64-7.60 (m, 2H), 7.55-7.51 (m, 1H), 7.17-7.12 (m, 3H), 6.89-6.84 (t, 1H), 1.43 (s, 4H). LC/MS: (M+1) 333.

Example 37




[0341] Cyclopropane-1,1-dicarboxylic acid (4-benzyloxy-phenyl)-amide (4-fluoro-phenyl)-amide. To a 0 °C suspension of 4-benzyloxyaniline hydrochloride (47.0 g, 200 mmol) and 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid (49.1 g, 220 mmol) in CH_2Cl_2 (400 mL) was added EDCI (38.2 g, 200 mmol). Stirring was continued at rt for 2-4 h until the reaction was complete. CH_2Cl_2 was removed under reduced pressure. H_2O (300 mL) and MeOH (200 mL) were added, and the resulting mixture was stirred at rt for 30 min. After filtration and wash with H_2O , the solid was transferred to another flask containing 300 mL of sat. aqueous NaHCO_3 solution. The mixture was stirred for another 30 min. The solid was filtered, washed with water, and dried over night

on a lyophilizer, affording cyclopropane-1,1-dicarboxylic acid (4-benzyloxy-phenyl)-amide (4-fluoro-phenyl)-amide (75.8g, 95% yield) as an off-white solid.


[0342] Cyclopropane-1,1-dicarboxylic acid (4-fluoro-phenyl)-amide (4-hydroxy-phenyl)-amide. To a refluxing mixture of cyclopropane-1,1-dicarboxylic acid (4-benzyloxy-phenyl)-amide (4-fluoro-phenyl)-amide (46 g, 113 mmol), 10% Pd/C (2 g) in EtOH (400 mL) was added dropwise 1,4-cyclohexadiene (62.7 mL, 678 mmol). Stirring was continued for 2-5 h until the reaction was complete. The mixture was cooled to rt, filtered through celite, and washed with EtOH. The solution was then concentrated under reduced pressure. To the flask containing the crude product was added CHCl_3 (200 mL). The resulting suspension was stirred for 15 min at rt. The solid was filtered, and dried in the air to give cyclopropane-1,1-dicarboxylic acid (4-fluoro-phenyl)-amide (4-hydroxy-phenyl)-amide (34.4 g, 95%, yield).

Example 38




[0343] Alternate Synthesis of Cyclopropane-1,1-dicarboxylic acid (4-fluoro-phenyl)-amide (4-hydroxy-phenyl)-amide. To a solution of 4-aminophenol (2.93 g, 26.9 mmol) and 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid (5.00 g, 22.4 mmol) in DMA (30 mL) was added EDCI (5.15 g, 26.9 mmol). The mixture was stirred vigorously until the reaction was complete (~ 3 h). With vigorous stirring, the reaction mixture was then poured into a flask containing sat. aqueous NaHCO_3 solution (200 mL). The stirring was continued for 1 h. The resulting suspension was then filtered. The solid was washed with water (50 mL), chloroform (50 mL) and dried under vacuum, affording 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid (6.22g, 88% yield) as a powder (>95% purity by HPLC and ¹H NMR).

Example 39



[0344] *N*-{4-[(7-benzyloxy-6-methoxyquinolin-4-yl)oxy]-3-fluorophenyl}-*N'*-(2-phenylethyl)ethanediamide. A mixture of 7-benzyloxy-4-chloro-6-methoxyquinoline (30 g, 100 mmol), *N*-(3-fluoro-4-hydroxyphenyl)-*N'*-(2-phenylethyl)ethanediamide (32 g, 106 mmol), DMAP (125 g, 1.02 mol) and bromobenzene (500 mL) was heated to reflux for 6 h. The mixture was cooled to room temperature and the bromobenzene was removed under reduced pressure. Methanol (500 mL) was added to the residue and the mixture was stirred at room temperature for 2 h. The resulting solid was filtered, washed with methanol and dried to give *N*-{4-[(7-benzyloxy-6-methoxyquinolin-4-yl)oxy]-3-fluorophenyl}-*N'*-(2-phenylethyl)ethanediamide (34 g, 61 %). ¹H NMR (400 MHz, *d*₆-DMSO): δ 11.05 (s, 1H), 9.15 (s, 1H), 8.47 (d, 1H), 8.05 (d, 1H), 7.84 (d, 1H), 7.56-6.36 (m, 13H), 6.46 (d, 1H), 5.32 (s, 2H), 3.97 (s, 3H), 3.47 (q, 2H), 2.86 (t, 2H); ¹³C NMR (100 MHz, *d*₆-DMSO): δ 160.5, 160.2, 159.9, 159.5, 155.2, 152.7, 152.2, 150.3, 149.6, 146.9, 139.7, 137.4, 137.3, 137.2, 137.1, 129.3, 129.2, 129.1, 129.0, 128.9, 128.7, 128.6, 126.9, 124.8, 117.9, 115.3, 109.9, 102.8, 99.8, 70.6, 56.5, 41.3, 35.2; IR (cm⁻¹): 1657, 1510, 1481, 1433, 1416, 1352, 1310, 1252, 1215, 1609, 986, 891, 868, 850, 742, 696; LC/MS (*M*+*H* = 566).


Example 40



[0345] *N*-{3-Fluoro-4-[(7-hydroxy-6-methoxyquinolin-4-yl)oxy]phenyl}-*N'*-(2-phenylethyl)ethanediamide. To a solution of *N*-{4-[(7-benzyloxy-6-methoxyquinolin-4-

yl)oxy]-3-fluorophenyl}-*N'*-(2-phenylethyl)ethanediamide (32 g, 56 mmol) in methanol (200 mL), DMF (100 mL), dichloromethane (100 mL), ethyl acetate (100 mL) and acetic acid (5 mL) was added palladium hydroxide (4.2 g) and the mixture was shaken on a Parr hydrogenator under a hydrogen pressure of 45 psi for 4 h. The resulting suspension was filtered through celite and the solid residue was washed with boiling dichloromethane (2 L) and acetone (2 L). The combined filtrates were evaporated to yield *N*-{3-fluoro-4-[(7-hydroxy-6-methoxyquinolin-4-yl)oxy]phenyl}-*N'*-(2-phenylethyl)ethanediamide as an off-white solid (25.6 g, 95 %). ¹H NMR (400 MHz, *d*₆-DMSO): δ 11.06 (s, 1H), 10.25 (br s, 1H), 9.12 (t, 1H), 8.40 (d, 1H), 8.01 (dd, 1H), 7.50-7.44 (m, 2H), 7.31-7.23 (m, 6H), 6.39 (d, 1H), 3.95 (s, 3H), 2.85 (t, 2H), 2.50 (m, 2H); IR (cm⁻¹): 1666, 1624, 1585, 1520, 1481, 1427, 1377, 1256, 1211, 1194, 1022, 880, 850, 839, 802, 750, 700; LC/MS (*M*+*H* = 476).


Example 41



[0346] *N*-(3-Fluoro-4-{[6-methoxy-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl]oxy}phenyl)-*N'*-(2-phenylethyl)ethanediamide. A solution of *N*-{3-fluoro-4-[(7-hydroxy-6-methoxyquinolin-4-yl)oxy]phenyl}-*N'*-(2-phenylethyl)ethanediamide (25.6 g, 54 mmol), *N*-(3-chloropropyl)morpholine hydrochloride (11.7 g, 592 mmol) and potassium carbonate (16.6 g, 120 mmol) in DMF (300 mL) was heated to 80 °C overnight. Upon cooling, a majority of the DMF (250 mL) was removed on a rotary evaporator, 5% aqueous LiCl (300 mL) was added and the mixture was sonicated at room temperature. The solid was filtered, suspended in 1N HCl and washed with ethyl acetate (2 x 300 mL). The solution was adjusted to pH 14 using 2N sodium hydroxide and subsequently extracted with dichloromethane (3 x 200 mL). The organic layer was dried over sodium sulfate, filtered and evaporated to give *N*-(3-fluoro-4-{[6-methoxy-7-(3-morpholin-4-ylpropoxy)quinolin-4-yl]oxy}phenyl)-*N'*-(2-phenylethyl)ethanediamide as a yellow solid (24 g, 74 %). ¹H NMR (400 MHz, CDCl₃): δ 9.37 (s, 1H), 8.46 (d, 1H), 7.81 (dd, 1H), 7.57 (t, 1H), 7.53 (s, 1H), 7.42 (s, 2H), 7.34-7.20 (m, 6H), 6.39 (d, 1H), 4.27 (t, 2H), 4.03 (s, 3H), 3.71 (m, 4H), 3.65 (q, 2H), 2.91 (t, 2H), 2.56 (br s, 4H), 2.13 (m, 2H); ¹³C NMR (100 MHz, *d*₆-DMSO): δ 160.1, 160.0, 159.5, 155.2, 152.7, 152.6, 150.2, 149.5, 147.1,


139.7, 137.3, 137.1, 129.3, 129.1, 126.9, 124.8, 117.9, 115.1, 109.2, 102.7, 99.6, 67.4, 66.9, 56.5, 55.5, 54.1, 41.3, 35.2, 26.4; IR (cm⁻¹): 1655, 1506, 1483, 1431, 1350, 1302, 1248, 1221, 1176, 1119, 864, 843, 804, 741, 700; LC/MS (M+H = 603).

Example 42




[0347] Cyclopropane-1,1-dicarboxylic acid [4-(7-benzyloxy-6-methoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-amide (4-fluoro-phenyl)-amide. To a flask containing cyclopropane-1,1-dicarboxylic acid (3-fluoro-4-hydroxy-phenyl)-amide (4-fluoro-phenyl)-amide (2.25 g, 6.7 mmol) and trifluoromethanesulfonic acid 7-benzyloxy-6-methoxy-quinolin-4-yl ester (1.87 g, 4.5 mmol) was added dry 2,6-lutidine (9 mL). The reaction mixture was heated to reflux (143°C) with vigorous stirring. The reaction progress was monitored by LC-MS. 2,6-Lutidine was removed under reduced pressure when the reaction was complete (about 6 h). The residue was treated with charcoal (1.5 g) in refluxing EtOAc (50 mL) for 15 min, and filtered through celite. The volume of the filtrate was reduced to about 20 mL and was added 20 mL of 1 N HCl. The crude product precipitated as the HCl salt, which was filtered and washed with EtOAc and H₂O (88% purity by analytical HPLC). The HCl salt was free-based with saturated aqueous NaHCO₃ solution. Further purification by column chromatography (hexans:EtOAc = 1:4) gave cyclopropane-1,1-dicarboxylic acid [4-(7-benzyloxy-6-methoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-amide (4-fluoro-phenyl)-amide as an off-white solid (1.3 g, 48% yield. ¹H NMR (400 MHz, DMSO, d₆): 10.41 (s, 1H), 10.02 (s, 1H), 8.48 (d, 1H), 7.92 (dd, 1H), 7.65 (m, 2H), 7.54 (m, 5H), 7.41 (m, 4H), 7.17 (m, 2H), 6.43 (d, 1H), 5.32 (s, 2H), 3.97 (s, 3H), 1.48 (m, 4H). LC/MS Calcd for [M+H]⁺ 596.2, found 596.3.

Example 43



[0348] Cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide. To a solution of the cyclopropane-1,1-dicarboxylic acid [4-(7-benzyloxy-6-methoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-amide (4-fluoro-phenyl)-amide (22.4 g, 37.6 mmol) in EtOH (340 mL) was added 1,4-cyclohexadiene (35 mL, 376 mmol) and 10% Pd/C (2.08 g). The reaction mixture was then heated at 65°C with stirring for 3 h (Caution: H₂ gas is released from the reaction). It was then allowed to cool to room temperature, and filtered through celite followed by a MeOH wash. The solution was then concentrated under reduced pressure. The yellow residue was taken into EtOAc (1 L). The EtOAc solution was washed with water (1X), brine (2X), dried over MgSO₄ and concentrated *in vacuo*. Cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide was obtained as a yellow solid (17.3 g, 91.1% yield), which were carried on to the next reaction without further purification. ¹H NMR (400 MHz, DMSO, d₆): 10.39 (s, 1H), 10.15 (s, 1H), 10.00 (s, 1H), 8.38 (d, 1H), 7.88 (dd, 1H), 7.63 (m, 2H), 7.50 (m, 2H), 7.40 (t, 1H), 7.27 (s, 1H), 7.14 (m, 2H), 6.33 (d, 1H), 3.95 (s, 3H), 1.47 (m, 4H). LC/MS Calcd for [M+H]⁺ 506.2, found 506.3. Anal. HPLC: 99.4% pure.


Example 44



[0349] N-[3-fluoro-4-({6-(methoxyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl)oxyphenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. To a mechanically stirred slurry of cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (16.6 g, 32.8 mmol) and

potassium carbonate (13.6 g, 98.6 mmol) in DMF (250 mL) was added 4-(3-chloropropyl)-morpholine hydrochloride (13, 7.92 g, 39.6 mmol). The resulting mixture was heated at 90°C for 5 hours (until phenol completely consumed). The reaction mixture was allowed to cool to room temperature, then dumped into water (900 mL), followed by extraction with EtOAc (3X). The combined extracts were washed with 5% LiCl (aq.) (3X) and brine (1X) followed by drying over MgSO₄ and concentration in vacuo. The crude (18.8g) obtained as brown solid was further purified by flash chromatography [silica gel, 4-stage gradient system: 1) EtOAc; 2) EtOAc:MeOH:7N NH₃/MeOH (95:5:0.5); 3) DCM:MeOH:7N NH₃/MeOH (95:5:0.5); 4) DCM:MeOH: 7N NH₃/MeOH (93:8:1)], affording N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide was obtained as an off white solid (15.0 g, 72% yield). ¹H NMR (400 MHz, DMSO-d₆): 10.41 (s, 1H), 10.02 (s, 1H), 8.47 (d, 1H), 7.91 (dd, 1H), 7.65 (m, 2H), 7.53 (m, 2H), 7.42 (t, 1H), 7.40 (s, 1H), 7.16 (m, 2H), 6.42 (d, 1H), 4.20 (t, 2H), 3.96 (s, 3H), 3.59 (t, 4H), 2.47 (t, 2H), 2.39 (br, s, 4H), 1.98 (m, 2H), 1.48 (m, 4H). LC/MS Calcd for [M+H]⁺ 633.3, found 633.0.


Example 45



[0350] Cyclopropane-1,1-dicarboxylic acid [4-(7-benzyloxy-6-methoxy-quinazolin-4-yloxy)-3-fluoro-phenyl]-amide (4-fluoro-phenyl)-amide: A mixture of 7-benzyloxy-4-chloro-6-methoxy-quinazoline (5 g, 16.67 mmol), cyclopropane-1,1-dicarboxylic acid (3-fluoro-4-hydroxy-phenyl)-amide (4-fluoro-phenyl)-amide (8.3 g, 25 mmol), potassium carbonate (125 mmol, 17.25 g), and dimethylacetamide (125 ml) was heated 50° C with stirring for 16h. Reaction mixture was poured onto ice/water (600 ml) and stirred for 30 minutes, and filtered. The solid was dissolved in ethyl acetate and washed with water (1x), brine, and concentrated. The crude was purified on silica get column eluting with 30% acetone in hexanes to yield cyclopropane-1,1-dicarboxylic acid [4-(7-benzyloxy-6-


methoxy-quinazolin-4-yloxy)-3-fluoro-phenyl]-amide (4-fluoro-phenyl)-amide (7.5 g, 76%). ^1H NMR (CDCl_3): 8.64 (1H, br. s), 8.55 (1H, s), 8.33 (1H, br. s), 7.74-7.71 (1H, dd), 7.54 (1H, s), 7.48-7.33 (8H, m), 7.31-7.24 (2H, m), 7.06-7.02 (2H, m), 5.32 (2H, s), 4.06 (3H, s), 1.77-1.74 (2H, m), 1.63-1.61 (2H, m).

Example 46



[0351] Cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinazolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide. To a mixture of cyclopropane-1,1-dicarboxylic acid [4-(7-benzyloxy-6-methoxy-quinazolin-4-yloxy)-3-fluoro-phenyl]-amide (4-fluoro-phenyl)-amide (7.5 g, 12.6 mmol), acetic acid (few drops), dichloromethane (50 ml) and methanol (100 ml) was added 10% Pd/C (700 mg). The mixture was agitated in hydrogen gas (40 psi) until the reaction was complete (ca. 4 hr). The solution was filtered through celite and concentrated to give a crude product as a solid. The crude product was triturated with ether, and filtered. The filter cake was dried *in vacuo* to yield cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinazolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (6.1 g, 95% yield). ^1H NMR (dmsd- d_6): 10.86 (1H, br. s), 10.34 (1H, br. s), 10.04 (1H, br. s), 8.46 (1H, s), 7.84-7.80 (1H, dd), 7.66-7.62 (2H, m), 7.55 (1H, s), 7.47-7.45 (1H, m), 7.41-7.37 (1H, m), 7.24 (1H, s), 7.18-7.13 (2H, t), 3.98 (3H, s), 1.46 (4H, s).


Example 47



[0352] N-[3-Fluoro-4-({6-(methoxy)-7-[(3-morpholin-4-yl)propyl]oxy}quinazolin-4-yl)oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. To a mixture of

cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinazolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (1.5 g, 2.96 mmol), 4-(3-hydroxypropyl)morpholine (0.623 mL, 4.5 mmol), triphenylphosphine (1.18 g, 4.5 mmol), and dichloromethane (50 mL) was added diisopropyl azodicarboxylate (0.886 mL, 4.5 mmol). The mixture was stirred at room temperature for 16 h, monitored by LCMS. After removal of solvent, the crude mixture was separated by flash column chromatography (silica), eluting with 5% methanol in dichloromethane to give N-[3-fluoro-4-({6-(methoxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarbox-amide (890 mg, 47% yield). ¹H NMR (400 MHz, DMSO-d₆): δ 10.36 (br s, 1H), 10.05 (br s, 1H), 8.55 (s, 1H), 7.83 (m, 1H), 7.64 (m, 2H), 7.57 (s, 1H), 7.44 (m, 3H), 7.18 (t, 2H), 4.27 (m, 2H), 3.99 (s, 3H), 3.61 (m, 6H), 2.40 (m, 4H), 2.01 (m, 2H), 1.47 (m, 4H). LC/MS Calcd for [M+H]⁺ 634.2, found 634.3.


Example 48



[0353] N-(4-([6,7-bis(methoxy)quinolin-4-yl]oxy)phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. To a solution of cyclopropane-1,1-dicarboxylic acid (4-fluoro-phenyl)-amide (4-hydroxy-phenyl)-amide (6.98 g, 22.2 mmol) in anhydrous 2,6-lutidine (50 mL) was added trifluoromethanesulfonic acid 6, 7-dimethoxy-quinolin-4-yl ester (5 g, 14.8 mmol). The reaction mixture was heated at 165°C in a sealed pressure tube with stirring for 18 h. The reaction mixture was concentrated on high vacuum to completely remove lutidine. The resulting solid material was dissolved in DCM (250 mL), and washed several times with 1 N sodium hydroxide to remove the excess phenol. The crude mixture was loaded on a silica gel flash column and eluted with 75% EtOAc-hexanes, affording N-(4-([6,7-bis(methoxy)quinolin-4-yl]oxy)phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (3.2 g, 44%). ¹H NMR (400 MHz, d₆-DMSO): δ 10.2 (s, 1H), 10.05 (s, 1H), 8.4 (s, 1H), 7.8 (m, 2H), 7.65 (m, 2H), 7.5 (s, 1H),


7.35 (s, 1H), 7.25 (m, 2H), 7.15(m, 2H), 6.4 (s, 1H), 4.0 (d, 6H), 1.5 (s, 4H). LC/MS: M+H= 502.

Example 49



[0354] 4,7-Dichloroquinoline. Phosphorus oxychloride (4mL, 429 mmol) was added to 7-chloro-4-hydroxyquinoline 2.86g, 15.9mmol) in a round bottom flask equipped with a reflux condenser. The mixture was heated to reflux for 2h, then allowed to cool to room temperature. The solution was concentrated *in vacuo* to a thick oil, then dumped over cracked ice. The resulting solution was neutralized with saturated NaHCO₃ (aq). The slurry was filtered and washed with water. The solids were dried under vacuum, afforded 4,7-dichloroquinoline as a white solid (2.79g, 88.5% yield).


Example 50



[0355] 4-[4-(2-Fluoro-4-{[1-(4-fluorophenyl)carbamoyl]-cyclopropanecarbonyl]-amino}-phenoxy)-6-methoxy-quinazolin-7-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester. Cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinazolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (325 mg, 0.64 mmol), 4-methanesulfonyloxymethyl-piperidine-1-carboxylic acid tert-butyl ester (193 mg, 0.66 mmol), K₂CO₃ (181 mg, 1.31 mmol) were combined in DMF (5 ml) and heated to 80°C overnight. The reaction was not complete and more 4-methanesulfonyloxymethyl-piperidine-1-carboxylic acid tert-butyl ester (90 mg, 0.31 mmol) and K₂CO₃ (90 mg, 0.65 mmol) were added and heating at 80°C continued for another night. The reaction mixture


was allowed to cool to room temperature, then diluted with EtOAc and washed with H₂O (3x), sat'd NaCl (1x), dried (Na₂SO₄) and concentrated *in vacuo*. The resulting crude material was purified by flash chromatography (1:1 hexanes:EtOAc, followed by 1:3 hexanes:EtOAc) to give 4-[4-(2-fluoro-4-{[1-(4-fluorophenylcarbamoyl)-cyclopropanecarbonyl]-amino}-phenoxy)-6-methoxy-quinazolin-7-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester (273 mg, 60%). LC/MS Calcd for [M+H]⁺ 704.3, found 704.4.

Example 51



[0356] Cyclopropane-1,1-dicarboxylic acid {3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-amide (4-fluoro-phenyl)-amide, TFA salt. 4-[4-(2-Fluoro-4-{[1-(4-fluorophenylcarbamoyl)-cyclopropanecarbonyl]-amino}-phenoxy)-6-methoxy-quinazolin-7-yloxymethyl]-piperidine-1-carboxylic acid tert-butyl ester (273 mg, 0.39 mmol) was dissolved in CH₂Cl₂ (8 ml) to which was added TFA (8 ml) and the mixture stirred at room temperature for 1hr. The reaction mixture was concentrated *in vacuo* and the resulting oil triturated with Et₂O. The resulting solids were filtered, washed with Et₂O and dried under high vacuum to give cyclopropane-1,1-dicarboxylic acid {3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-amide (4-fluoro-phenyl)-amide, TFA salt (222 mg, 80%). LC/MS Calcd for [M+H]⁺ 604.2, found 604.3.

Example 52



[0357] N-{3-Fluoro-4-[6-(methoxy)-7-[[1-(4-fluorophenyl)carbamoyl]-cyclopropanecarbonyl]-amino]-phenoxy]-6-methoxy-quinazolin-4-yl}oxyphenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide.

Cyclopropane-1,1-dicarboxylic acid {3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-amide (4-fluoro-phenyl)-amide, TFA salt (222 mg, 0.31 mmol), H₂O (3 ml), 37% formaldehyde in H₂O (0.18 ml) and acetic acid (27 drops) were combined in acetonitrile (9 ml) to which was slowly added triacetoxyborohydride (561 mg, 2.65 mmol). The mixture was stirred at room temperature for 1-2 hr, then diluted with 1N NaOH and H₂O and extracted with CH₂Cl₂ (3x). The combined CH₂Cl₂ extractions were washed with sat'd NaCl (1x), dried (Na₂SO₄) and concentrated *in vacuo*. The resulting residue was dissolved in a minimum of 1:1 dioxane:EtOAc to which was added 4M HCl in dioxane (1-2 ml). The resulting solids were filtered, washed with EtOAc and dried under high vacuum to give N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinazolin-4-yl)oxy] phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide, HCl salt (167 mg, 83%). ¹HNMR (400MHz, DMSO-d₆): δ 10.40 (s, 1H), 10.17 (br s, 1H) 10.07 (s, 1H), 8.61 (s, 1H), 7.85 (m, 1H), 7.65 (m, 2H), 7.48 (m, 2H), 7.42 (t, 1H), 7.16 (t, 2H), 4.12 (2, 2H), 4.00 (s, 3H), 3.46 (m, 2H), 2.99 (m, 2H), 2.73 (d, 3H), 2.13 (m, 1H), 2.01 (m, 2H), 1.63 (m, 2H), 1.47 (m, 4H). LC/MS Calcd for [M+H]⁺ 618.2, found 618.3.

Synthesis of Bridged Bicyclics

[0358] The following describes synthesis of bridged bicyclics with appended leaving groups for use as, for example, alkylating agents. In the context of this invention, these alkylating agents are used, for example, to alkylate the quinazoline or quinolines on the 6- or 7-oxygens to make compounds of the invention. The invention is not limited to alkylation chemistry to append such bridged bicyclics, but rather the aforementioned description is meant only to be illustrative of an aspect of the invention.

Example 53

[0359] 1,4:3,6-dianhydro-2-O-methyl-5-O-(methanesulfonyl)-D-glucitol: To a solution of 1,4:3,6-dianhydro-2-O-methyl-D-glucitol (1.19g, 7.4 mmol) in dichloromethane was added pyridine (1mL, 12.36 mmol) followed by methanesulfonyl chloride (0.69mL, 8.92 mmol) and the mixture was allowed to stir at room temperature over 12 hours. The solvent was removed and the amorphous residue was partitioned with ethyl acetate and 0.1M aqueous hydrochloric acid. The aqueous phase was extracted once with additional

ethyl acetate and the combined organic layers were washed with saturated aqueous sodium chloride then dried over anhydrous magnesium sulfate. Filtration and concentration followed by drying *in vacuo* afforded 1,4:3,6-dianhydro-2-*O*-methyl-5-*O*-(methylsulfonyl)-D-glucitol (1.67g, 94% yield) as a colorless oil. GC/MS calculated for $C_8H_{14}SO_6$: 238 (M^+).

Example 54

[0360] 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-D-fructose ethylene glycol acetal: A solution of 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-D-fructose (2.00g, 8.06 mmol), ethylene glycol (5.00g, 80.6 mmol), and *p*-toluenesulfonic acid (1.53g, 8.06 mmol) in benzene (100mL) was refluxed for 90 min using a Dean-Stark Trap apparatus. The reaction mixture was diluted with ethyl acetate (100mL), washed with saturated aqueous sodium bicarbonate (2 x 50mL) then brine (50mL), and dried over anhydrous sodium sulfate. Filtration, concentration and column chromatography on silica (1:1 hexane/ethyl acetate) provided 1.44g (61% yield) of 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-D-fructose ethylene glycol acetal as a colorless solid. 1H NMR (400 MHz; $CDCl_3$): 8.08 (m, 2H), 7.58 (m, 1H), 7.54 (m, 2H), 5.38 (dd, 1H), 4.97 (t, 1H), 4.21-4.02 (m, 7H), 3.86 (d, 1H), 3.75 (d, 1H).

Example 55

[0361] 1,4:3,6-dianhydro-D-fructose ethylene glycol acetal: To a solution of 1,4:3,6-dianhydro-5-*O*-(phenylcarbonyl)-D-fructose ethylene glycol acetal (1.44g, 4.93 mmol) in methanol (40mL) was added 50% aqueous sodium hydroxide (0.38 g, 4.75 mmol) and the mixture was stirred at room temperature for 30 minutes. Neutralization with 1M HCl, followed by concentration and column chromatography on silica (1:2 hexane/ethyl acetate) provided 0.74g (80% yield) of 1,4:3,6-dianhydro-D-fructose ethylene glycol acetal as a colorless solid. 1H NMR (400 MHz; $CDCl_3$): 4.60 (t, 1H), 4.32 (m, 1H), 4.14 (d, 1H), 4.05-3.98 (m, 5H), 3.82 (s, 2H), 3.62 (dd, 1H), 2.65 (d, 1H).

[0362] 1,4:3,6-dianhydro-5-*O*-(methylsulfonyl)-D-fructose ethylene glycol acetal: To a solution of 1,4:3,6-dianhydro-D-fructose ethylene glycol acetal (0.74g, 3.93 mmol) and triethylamine (1.20g, 11.86 mmol) in dichloromethane (40mL) was added

methanesulfonyl chloride (0.90g, 7.88 mmol) at 0°C under nitrogen. The solution was warmed to room temperature and stirred for 13 h. Dichloromethane (50mL) was added, and the organic layer was washed with saturated aqueous sodium bicarbonate (30mL), water (30mL), and brine (30mL) then dried over anhydrous sodium sulfate. Filtration and concentration provided 1.02g (97%) of 1,4:3,6-dianhydro-5-*O*-(methylsulfonyl)-D-fructose ethylene glycol acetal as a yellow oil. ¹H NMR (400 MHz; CDCl₃): 5.08 (m, 1H), 4.82 (t, 1H), 4.13 (dd, 1H), 4.04 (m, 4H), 3.93 (dd, 1H), 3.87 (d, 1H), 3.81 (d, 1H), 3.13 (s, 3H).

Example 56

[0363] 1,4:3,6-dianhydro-2-deoxy-2-methylidene-D-arabino-hexitol: To a solution of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-5-*O*-(phenylcarbonyl)-D-arabino-hexitol (329mg, 1.34 mmol) in methanol (10mL) was added 50% aqueous sodium hydroxide (95mg, 1.19 mmol) and the mixture was stirred at room temperature for 30 minutes. Neutralization with 4M hydrogen chloride in 1,4-dioxane followed by concentration and column chromatography on silica (1:1 hexane/ethyl acetate) provided 141mg (74%) of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-D-arabino-hexitol as a colorless solid. ¹H NMR (400 MHz; CDCl₃): 5.37 (m, 1H), 5.20 (m, 1H), 4.80 (m, 1H), 4.54 (m, 2H), 4.43 (m, 1H), 4.26 (m, 1H), 3.95 (dd, 1H), 3.54 (dd, 1H), 2.70 (d, 1H).

[0364] 1,4:3,6-dianhydro-2-deoxy-2-methylidene-5-*O*-(methylsulfonyl)-D-arabino-hexitol: To a solution of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-D-arabino-hexitol (135mg, 0.95 mmol) and triethylamine (288mg, 2.85 mmol) in dichloromethane (10mL) was added methanesulfonyl chloride (222mg, 1.94 mmol) at 0°C under nitrogen. The solution was warmed to room temperature and stirred for 18 h. Dichloromethane (50mL) was added and the organic layer was washed with saturated aqueous sodium bicarbonate (2 x 25mL), water (25mL) and brine (25mL) then dried over anhydrous sodium sulfate. Filtration and concentration provided 213mg (72%) of 1,4:3,6-dianhydro-2-deoxy-2-methylidene-5-*O*-(methylsulfonyl)-D-arabino-hexitol as a yellow oil. ¹H NMR (400 MHz; CDCl₃): 5.40 (m, 1H), 5.23 (m, 1H), 5.04 (m, 1H), 4.85 (m, 1H), 4.73 (t, 1H), 4.58 (m, 1H), 4.41 (m, 1H), 4.08 (dd, 1H), 3.86 (dd, 1H), 3.14 (s, 3H).

Example 57

[0365] 1,4:3,6-dianhydro-2-deoxy-5-O-(phenylcarbonyl)-L-arabino-hex-1-enitol: To a mixture of 1,4:3,6-dianhydro-5-O-(phenylcarbonyl)-(D)-glycitol (4.32g, 17.3 mmol), triethylamine (4.91 mL, 35.3 mmol) and 4-dimethylaminopyridine (0.63g, 5.2 mmol) in dichloromethane (50 mL) at -10 ° to -15° was added trifluoromethanesulfonic anhydride (3.48mL, 20.7 mmol) dropwise over ten minutes and the resulting mixture was stirred at this temperature for 3 hours. The mixture was poured into 100 mL of ice-water and extracted with dichloromethane (3 x 50 mL). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered then concentrated. The crude triflate was suspended in toluene (50 mL) followed by addition of 1,8-diazabicyclo[4,5,0]undec-7-ene (5.25 mL, 34.6 mmol) and the mixture was stirred at room temperature for 18 hours. The reaction mixture was poured into ice-water and partitioned then the aqueous portion was extracted with dichloromethane (3 x 50 mL). The combined organic portion was washed with brine, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by flashed chromatography (silica gel, 5-20% ethyl acetate-hexane) to give 1,4:3,6-dianhydro-2-deoxy-5-O-(phenylcarbonyl)-L-arabino-hex-1-enitol, as a white solid, 3.10g, 77% yield. ¹H NMR (400MHz; CDCl₃): 8.08-8.06 (m, 2H), 7.61-7.57 (m, 1H), 7.56-7.43 (m, 2H), 6.62-6.61 (d, 1H), 5.48-5.46 (m, 1H), 5.32-5.26 (m, 1H), 5.13-5.10 (m, 2H), 4.18-4.14 (tr, 1H), 3.61-3.56 (tr, 1H).

Example 58

[0366] Methyl 3,6-anhydro-5-O-(phenylcarbonyl)-β-L-glucofuranoside: To a solution of 1,4:3,6-dianhydro-2-deoxy-5-O-(phenylcarbonyl)-L-arabino-hex-1-enitol (1.00g, 4.3 mmol) in methanol (17 mL) at -4°C was added 3-chloroperoxybenzoic acid (85%, 1.35g, 8.6 mmol), and the resulting mixture was slowly warmed to room temperature and stirred for 18 hours. The reaction mixture was concentrated, diluted with dichloromethane (50 mL), washed with saturated aqueous sodium bicarbonate solution, dried over sodium sulfate, filtered and concentrated. The residue was purified by flash chromatography (silica gel, 25-60% ethyl acetate-hexane) to give methyl 3,6-anhydro-5-O-(phenylcarbonyl)-β-L-glucofuranoside as a white solid, 1.03g, 83% yield. ¹H NMR (400MHz; CDCl₃): 8.11-8.08 (d, 2H), 7.61-7.56 (tr, 1H), 7.48-7.44 (m, 2H), 5.24-5.17

(m, 2H), 4.96 (s, 1H), 4.57-4.56 (d, 1H), 4.27 (s, 1H), 4.22-4.18 (dd, 1H), 4.08-4.04 (dd, 1H) 3.36 (s, 3H).

[0367] Methyl 3,6-anhydro-2-O-methyl-5-O-(phenylcarbonyl)- β -L-glucofuranoside: A mixture of methyl 3,6-anhydro-5-O-(phenylcarbonyl)- β -L-glucofuranoside (1.03g, 3.7 mmol), silver (I) oxide (0.85g, 3.7 mmol) and methyl iodide (0.34 mL, 5.5 mmol) in DMF (2 mL) was heated at 60°C for 1 hour. After cooling to room temperature the reaction mixture was diluted with ethyl acetate (50 mL), filtered over celite, adsorbed on silica gel (10g) and purified by flash chromatography (silica gel, 5-30% ethyl acetate-hexane) to give methyl 3,6-anhydro-2-O-methyl-5-O-(phenylcarbonyl)- β -L-glucofuranoside as a colorless oil, 0.82g, 76% yield. ^1H NMR (400MHz; CDCl_3): 8.11-8.09 (d, 2H), 7.60-7.56 (m, 1H), 7.46-7.44 (m, 2H), 5.24-5.20 (m, 1H), 5.18-5.09 (tr, 1H), 4.99 (s, 1H), 4.61-4.60 (d, 1H), 4.21-4.17 (tr, 1H), 4.08-4.03 (tr, 1H), 3.81 (s, 1H), 3.40 (s, 3H), 3.57 (s, 3H).

[0368] Methyl 3,6-anhydro-2-O-methyl- α -D-idofuranoside: A solution of methyl 3,6-anhydro-2-O-methyl-5-O-(phenylcarbonyl)- β -L-glucofuranoside (820mg, 3.1mmol) and 50% sodium hydroxide (248 mg, 3.1 mmol) in methanol (10mL) was stirred at room temperature for 30 minutes. The material was adsorbed on silica gel (5g) and passed through a short column (15% ethyl acetate in hexanes to 5% methanol in ethyl acetate) to give methyl 3,6-anhydro-2-O-methyl- α -D-idofuranoside as a colorless oil, 420 mg, 85% yield. ^1H NMR (400MHz; CDCl_3): 5.04 (s, 1H), 5.84-5.81 (tr, 1H), 4.44-4.42 (tr, 1H), 4.25-4.19 (m, 1H), 3.85-3.75 (m, 1H), 3.49 (s, 3H), 3.43 (s, 3H), 2.75-2.72 (d, 1H).

[0369] Methyl 3,6-anhydro-2-O-methyl-5-O-(methylsulfonyl)- β -L-glucofuranoside: Methyl 3,6-anhydro-2-O-methyl- α -D-idofuranoside (420 mg, 2.6 mmol) was dissolved in dichloromethane (10 mL) and pyridine (0.36 mL, 3.7 mmol) at 0°C. Methanesulfonyl chloride (0.14 mL, 3.1 mmol) was added and the resulting mixture was stirred at 0°C for 1 hour then at room temperature for 2 hours. The reaction mixture was washed with water and saturated aqueous sodium bicarbonate solution, dried over anhydrous sodium sulfate, filtered and concentrated to give methyl 3,6-anhydro-2-O-methyl-5-O-(methylsulfonyl)- β -L-glucofuranoside as a colorless oil, 669mg, 95% yield, which was used without further purification.

Example 59

[0370] 3,6-anhydro-5-O-(phenylcarbonyl)- α -L-glucofuranose: A mixture of osmium tetroxide (4% in water, 0.25 mL, 0.03 mmol) and N-methylmorpholine (505 mg, 4.3 mmol) in 3 mL of 50% acetone in water was warmed to 60°C. A solution of 1,4:3,6-dianhydro-2-deoxy-5-O-(phenylcarbonyl)-L-arabino-hex-1-enitol (2.00g, 8.6 mmol) in 6 mL of 50% acetone in water was added over 3 hours. During this time an additional amount of N-methylmorpholine (1.01g, 8.6 mmol) was added in small portions periodically. Upon completion of the addition process the reaction was stirred for another hour and cooled to room temperature. The crude mixture was applied to a column of silica gel and flashed (0-6% methanol in 1:1 ethyl acetate:hexane) to give 3,6-anhydro-5-O-(phenylcarbonyl)- α -L-glucofuranose as a white solid, 1.5g, 65% yield. ¹H NMR (400MHz; DMSO-d₆): 8.01-7.95, (m, 2H), 7.68-7.66 (m, 1H), 7.57-7.53 (m, 2H), 5.18-5.11 (m, 2H), 4.85-4.81 (m, 1H, m), 4.37-4.35 (m, 1H), 4.05-3.96 (m, 2H), 3.85-3.83 (m, 1H).

[0371] 3,6-anhydro-2-O-methyl-5-O-(phenylcarbonyl)- α -L-glucofuranoside: 3,6-Anhydro-5-O-(phenylcarbonyl)- α -L-glucofuranose (576 mg, 2.2 mmol) was added to a mixture of sodium hydride (60% oil dispersion, 346 mg, 8.7 mmol) and methyl iodide (0.54mL, 8.7 mmol) in 5 mL of DMF at 0°C and the resulting mixture was stirred for 1 hour. The reaction mixture was diluted with ethyl acetate and quenched with water (5 mL). The aqueous portion was extracted with ethyl acetate (3 x 5 mL). The combined organic portion was washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The residue was purified by flashed chromatography (silica gel, 5-20% ethyl acetate in hexane) to give 3,6-anhydro-2-O-methyl-5-O-(phenylcarbonyl)- α -L-glucofuranoside as a white solid, 270 mg, 42% yield. ¹H NMR (400MHz; CDCl₃): 8.09-8.07 (m, 2H), 7.61-7.57 (m, 1H), 7.48-7.27 (m, 2H), 5.25-5.22 (m, 1H), 5.07-5.06 (d, 1H), 4.94-4.91 (m, 1H), 4.73-4.71 (m, 1H), 4.20-4.16 (m, 1H), 3.96-3.94 (m, 1H), 3.85-3.83 (tr, 1H), 3.50 (s, 3H), 3.42 (s, 3H).

[0372] Methyl 3,6-anhydro-2-O-methyl-5-O-(methylsulfonyl)- α -L-glucofuranoside: A solution of methyl 3,6-anhydro-2-O-methyl-5-O-(phenylcarbonyl)- α -L-glucofuranoside (230mg, 0.92 mmol) and 50% sodium hydroxide (74 mg, 0.92 mmol) in methanol (5 mL) was stirred at room temperature for 30 minutes. The mixture was adsorbed on silica gel (2g) and passed through a short column (15% ethyl acetate in hexanes to 5% methanol in ethyl acetate) to afford a colorless oil which was employed directly in the next step, 140

mg, 0.72 mmol, 95% yield. The alcohol was dissolved in dichloromethane (5 mL) and pyridine (121 μ L, 1.03 mmol) was added at 0°C. Methanesulfonyl chloride (27 μ L, 0.88 mmol) was added and the resulting mixture was stirred at 0°C for 1 hour then at room temperature for 2 hours. The reaction mixture was washed with water and saturated aqueous sodium bicarbonate solution, dried over sodium sulfate, filtered and concentrated to give methyl 3,6-anhydro-2-*O*-methyl-5-*O*-(methylsulfonyl)- α -L-glucofuranoside as a colorless oil, 190 mg, 96% yield.

Example 60

[0373] 3,6-Anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(phenylcarbonyl)- α -L-glucofuranose: A mixture of 3,6-anhydro-5-*O*-(phenylcarbonyl)- α -L-glucofuranose (1.00g), 2,2-dimethoxy propane (0.63 mL), *p*-toluenesulfonic acid (20 mg) and benzene (10 mL) was heated at reflux for 3 hours. The reaction mixture was cooled then adsorbed on silica gel (10g) and purified by flash chromatography (silica gel, 5-35% ethyl acetate in hexanes) to give 3,6-anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(phenylcarbonyl)- α -L-glucofuranose as colorless oil, 0.85g, 74% yield. ^1H NMR (400MHz; CDCl_3): 8.08-8.06 (d, 2H), 7.59-7.56 (tr, 1H), 7.46-7.42 (m, 2H), 5.99-5.98 (d, 1H), 5.35-5.31 (tr, 1H), 5.10-5.08 (d, 1H), 4.66-4.65 (d, 1H), 4.61-4.60 (d, 1H), 4.20-4.16 (dd, 1H), 3.91-3.74 (tr, 1H), 1.50 (s, 3H), 1.34 (s, 3H).

[0374] 3,6-Anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(methylsulfonyl)- α -L-glucofuranose: A solution of 3,6-anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(phenylcarbonyl)- α -L-glucofuranose (850mg) and 50% sodium hydroxide (111 mg) in methanol (10mL) was stirred at room temperature for 30 minutes. The mixture was then adsorbed on silica gel (5g) and passed through a short column (15% ethyl acetate in hexanes to 5% methanol in ethyl acetate) and the alcohol intermediate, 390 mg, 70% yield, was used immediately in the next step. The alcohol was dissolved in dichloromethane (10 mL) and pyridine (0.32 mL) at 0°C. Methanesulfonyl chloride (0.12 mL) was added and the resulting mixture was stirred at 0°C for 1 hour then at room temperature for 2 hours. The reaction mixture was washed with water and saturated aqueous sodium bicarbonate solution, dried over anhydrous sodium sulfate, filtered and concentrated to give 3,6-anhydro-1,2-*O*-(1-methylethylidene)-5-*O*-(methylsulfonyl)- α -L-glucofuranose as a colorless oil, 485 mg, 90% yield, which was immediately employed in the next step.

Example 61

[0375] (3S,8aS)-3-(Chloromethyl)hexahydro-1H-pyrrolo[2,1-c][1,4]oxazine: (S)-(+)-Prolinol (6.00 g, 59.3 mmol) was added to epichlorohydrin (47 mL, 600 mmol) at 0°C. The solution was stirred at 40°C for 0.5 h and then concentrated *in vacuo*. The residual oil was cooled in an ice bath and concentrated sulfuric acid (18 mL) was added dropwise with stirring. The mixture was heated at 170-180°C for 1.5 h, poured into ice (300 mL) and then basified with sodium carbonate to pH~8. The mixture was partitioned with ethyl acetate/hexanes and filtered. The filtrate was separated and the aqueous portion was extracted twice with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford oil that was purified by column chromatography (ethyl acetate for less polar product and then 30% methanol in ethyl acetate). (3S,8aS)-3-(Chloromethyl)hexahydro-1H-pyrrolo[2,1-c][1,4]oxazine (less polar product) (1.87 g, 10.7 mmol, 18% yield): ¹H NMR (400 MHz, CDCl₃): 4.06 (dd, 1H), 3.79-3.71 (m, 1H), 3.60-3.48 (m, 2H), 3.36 (dd, 1H), 3.15 (dd, 1H), 3.13-3.06 (m, 1H), 2.21-2.01 (m, 3H), 1.90-1.68 (m, 3H), 1.39-1.24 (m, 1H); MS (EI) for C₈H₁₄NOCl: 176 (MH⁺). (3R,8aS)-3-(Chloromethyl)hexahydro-1H-pyrrolo[2,1-c][1,4]oxazine (1.54 g, 8.77 mmol, 15% yield): ¹H NMR (400 MHz, CDCl₃): 3.94-3.77 (m, 4H), 3.55 (dd, 1H), 3.02-2.93 (m, 2H), 2.45 (dd, 1H), 2.29-2.15 (m, 2H), 1.88-1.64 (m, 3H), 1.49-1.38 (m, 1H); MS (EI) for C₈H₁₄NOCl: 176 (MH⁺).

[0376] Using the same or analogous synthetic techniques and/or substituting with alternative starting materials, the following were prepared:

[0377] (3R,8aR)-3-(Chloromethyl)hexahydro-1H-pyrrolo[2,1-c][1,4]oxazine: ¹H NMR (400 MHz, CDCl₃): 4.05 (dd, 1H), 3.79-3.70 (m, 1H), 3.61-3.48 (m, 2H), 3.35 (dd, 1H), 3.15 (dd, 1H), 3.13-3.07 (m, 1H), 2.21-2.01 (m, 3H), 1.89-1.67 (m, 3H), 1.39-1.25 (m, 1H); MS (EI) for C₈H₁₄NOCl: 176 (MH⁺).

[0378] (3S,8aR)-3-(Chloromethyl)hexahydro-1H-pyrrolo[2,1-c][1,4]oxazine: ¹H NMR (400 MHz, CDCl₃): 3.93-3.77 (m, 4H), 3.55 (dd, 1H), 3.02-2.93 (m, 2H), 2.45 (dd, 1H), 2.30-2.15 (m, 2H), 1.88-1.64 (m, 3H), 1.49-1.37 (m, 1H); MS (EI) for C₈H₁₄NOCl: 176 (MH⁺).

Example 62

[0379] (3S,8aS)-Hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl acetate: (3S,8aS)-3-(Chloromethyl)hexahydro-1H-pyrrolo[2,1-c][1,4]oxazine (2.30 g, 13.1 mmol) and potassium acetate (12.8 g, 131 mmol) were stirred in dimethylformamide (25 mL) at 140°C for 20 h. The mixture was partitioned between ethyl acetate and water. The organic portion was washed twice with water, then with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford (3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl acetate as a brown oil (2.53 g, 12.7 mmol, 97% yield). ¹H NMR (400 MHz, CDCl₃): 4.14-4.02 (m, 3H), 3.81-3.72 (m, 1H), 3.37-3.31 (m, 1H), 3.09 (dt, 1H), 3.00 (dd, 1H), 2.21-2.00 (m, 3H), 2.10 (s, 3H), 1.90-1.67 (m, 3H), 1.39-1.24 (m, 1H); MS (EI) for C₁₀H₁₇NO₃: 200 (MH⁺).

[0380] (3S,8aS)-Hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethanol: (3S,8aS)-Hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl acetate (2.36 g, 11.9 mmol) was treated with sodium methoxide (25 wt% solution in methanol; 2.7 mL) for 0.5 h. The mixture was cooled in an ice bath and a solution of 4M HCl in 1,4-dioxane (3 mL, 12.0 mmol) was added slowly. The mixture was stirred at room temperature for 5 minutes and then was concentrated *in vacuo* to afford a suspension which was diluted with dichloromethane, filtered and the filtrate was concentrated *in vacuo* to afford (3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethanol as a brown oil (1.93 g, >100% yield). ¹H NMR (400 MHz, CDCl₃): 4.05 (dd, 1H), 3.73-3.65 (m, 2H), 3.62-3.56 (m, 1H), 3.39-3.34 (m, 1H), 3.10 (dt, 1H), 3.00-2.95 (m, 1H), 2.24-1.98 (m, 4H), 1.97-1.70 (m, 3H), 1.44-1.28 (m, 1H); MS (EI) for C₈H₁₅NO₂: 158 (MH⁺).

[0381] (3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethyl methanesulfonate: (3S,8aS)-Hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-ylmethanol (1.00 g, 6.37 mmol) was dissolved in dichloromethane (10 mL) and triethylamine (2.4 mL, 17.3 mmol) was added at 0°C followed by dropwise addition of methanesulfonyl chloride (0.93 mL, 12.0 mmol). The solution was warmed to room temperature and stirred for 1.25 h and then was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate solution. The organic portion was washed with saturated sodium bicarbonate solution. The combined aqueous portion was extracted with ethyl acetate. The combined organic portion was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford (3S,8aS)-hexahydro-1H-pyrrolo[2,1-c][1,4]oxazin-3-

ylmethyl methanesulfonate as an orange-brown oil (1.20 g, 5.1 mmol, 80% yield). MS (EI) for $C_9H_{17}NO_4S$: 236 (MH^+).

Example 63

[0382] Octahydro-2H-quinolizin-3-ylmethanol: Ethyl octahydro-2H-quinolizine-3-carboxylate (2.35 g, 11.1 mmol) was added dropwise to a stirred suspension of lithium aluminum hydride (1 M solution in tetrahydrofuran, 33 mL, 33 mmol) in tetrahydrofuran (50 mL) at 0°C. The reaction was stirred at room temperature for 3 h. The mixture was cooled in an ice bath and ethyl acetate (6 mL) was added slowly, followed by water (1.25 mL), 15% aqueous sodium hydroxide solution (5 mL) and water (1.25 mL). The mixture was filtered through a pad of celite and washed with ether. The filtrate was concentrated *in vacuo* and dried rigorously to afford octahydro-2H-quinolizin-3-ylmethanol as a yellow oil (1.66 g, 9.82 mmol, 88% yield). MS (EI) for $C_{10}H_{19}NO$: 170 (MH^+).

[0383] Octahydro-2H-quinolizin-3-ylmethyl methanesulfonate: Octahydro-2H-quinolizin-3-ylmethanol (600 mg, 3.55 mmol) was dissolved in dichloromethane (8 mL) and triethylamine (1.5 mL, 10.8 mmol) was added at 0°C followed by dropwise addition of methanesulfonyl chloride (0.56 mL, 7.16 mmol). The solution was warmed to room temperature and stirred for 1.25 h and then was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and saturated sodium bicarbonate solution. The aqueous portion was extracted with ethyl acetate. The combined organic portion was washed with brine, dried over sodium sulfate, filtered and concentrated *in vacuo* to afford octahydro-2H-quinolizin-3-ylmethyl methanesulfonate as an orange oil (796 mg, 3.22 mmol, 91% yield). MS (EI) for $C_{11}H_{21}NO_3S$: 248 (MH^+).

Example 64

[0384] (3S,8aS)-3-(Hydroxymethyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one: A solution of methyl 1-[(2S)-3-hydroxy-2-({[(phenylmethyl)oxy]carbonyl}amino)propyl]-L-prolinate (3.50 g, 10.4 mmol) in methanol was added to 5% palladium on carbon (50 wt.% in water) in methanol and treated with hydrogen at 40 psi for 1 h. The mixture was filtered and the filtrate was brought to reflux briefly and then cooled and concentrated *in vacuo* to afford (3S,8aS)-3-(hydroxymethyl)hexahydropyrrolo[1,2-a]pyrazin-1(2H)-one as

a colorless solid (1.50 g, 8.83 mmol, 85% yield). ^1H NMR (400 MHz, CDCl_3): 7.28-7.22 (m, 1H), 3.83-3.75 (m, 1H), 3.69 (dd, 1H), 3.56 (dd, 1H), 3.31 (t, 1H), 3.08 (dd, 1H), 2.92 (dt, 1H), 2.76-2.70 (m, 1H), 2.66 (dd, 1H), 2.28-2.16 (m, 1H), 2.02-1.73 (m, 3H); MS (EI) for $\text{C}_8\text{H}_{14}\text{N}_2\text{O}_2$: 171 (MH^+).

[0385] (3*S*,8*aS*)-3-([(1,1-Dimethylethyl)(dimethyl)silyl]oxy)methyl)hexahydro-pyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: To a solution of (3*S*,8*aS*)-3-(hydroxymethyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one (1.49 g, 8.82 mmol) in dimethylformamide (20 mL) was added triethylamine (2.45 mL, 17.6 mmol) and 4-dimethylaminopyridine (90 mg, 0.882 mmol). The solution was cooled in an ice bath and *tert*-butyldimethylsilyl chloride (2.66 g, 17.6 mmol) was added. The mixture was warmed to room temperature and stirred for 14 h. The mixture was concentrated *in vacuo* and the residue was partitioned between ethyl acetate and water. The aqueous portion was extracted twice with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford a pale brown solid which was triturated with ethyl acetate to afford (3*S*,8*aS*)-3-([(1,1-dimethylethyl)(dimethyl)silyl]oxy)methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as an off-white solid (1.74 g, 5.84 mmol, 66% yield). ^1H NMR (400 MHz, CDCl_3): 6.09-5.90 (m, 1H), 3.86-3.76 (m, 1H), 3.63 (dd, 1H), 3.44 (dd, 1H), 3.25 (t, 1H), 3.10 (ddd, 1H), 2.98-2.90 (m, 1H), 2.68-2.60 (m, 1H), 2.52 (dd, 1H), 2.28-2.18 (m, 1H), 2.06-1.95 (m, 1H), 1.93-1.74 (m, 2H), 0.90 (s, 9H), 0.07 (s, 6H); MS (EI) for $\text{C}_{14}\text{H}_{28}\text{N}_2\text{O}_2\text{Si}$: 285 (MH^+).

[0386] (3*S*,8*aS*)-3-([(1,1-Dimethylethyl)(dimethyl)silyl]oxy)methyl)-2-methylhexahydro-pyrrolo[1,2-*a*]pyrazin-1(2*H*)-one: (3*S*,8*aS*)-3-([(1,1-Dimethylethyl)(dimethyl)silyl]oxy)methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one (1.51 g, 5.32 mmol) in dimethylformamide (8 mL) was added to an ice-cooled suspension of sodium hydride (60 wt.% dispersion in oil; 213 mg, 5.32 mmol) in dimethylformamide (8 mL). The mixture was stirred at 0°C for 0.25 h and then iodomethane (0.332 mL, 5.32 mmol) was added dropwise. The mixture was stirred at room temperature for 0.5 h and then was stirred at 70°C for 2 h. The mixture was concentrated *in vacuo* and the residue was partitioned between ethyl acetate and water. The aqueous portion was extracted with ethyl acetate. The combined organic portion was dried over sodium sulfate, filtered and concentrated *in vacuo* to afford (3*S*,8*aS*)-3-([(1,1-dimethylethyl)(dimethyl)silyl]oxy)methyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as a yellow oil (1.552 g, 5.21 mmol) which was dissolved in tetrahydrofuran (20 mL) and treated with tetrabutylammonium

fluoride (1.0M solution in tetrahydrofuran; 10.4 mL, 10.4 mmol) for 2 h at room temperature. The mixture was concentrated *in vacuo* and purified by column chromatography (10% methanol in dichloromethane) to afford (3*S*,8*aS*)-3-(hydroxymethyl)-2-methylhexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one as a yellow oil (496mg, 2.70mmol, 51% yield from (3*S*,8*aS*)-3-([(1,1-dimethylethyl)(dimethyl)silyl]oxy) methyl)hexahydropyrrolo[1,2-*a*]pyrazin-1(2*H*)-one). ¹H NMR (400 MHz, CDCl₃): 3.98-3.93 (m, 1H), 3.86 (dd, 1H), 3.61-3.55 (m, 1H), 3.29-3.25 (m, 1H), 3.09-3.03 (m, 1H), 3.03-2.97 (m, 1H), 3.02 (s, 3H), 2.93 (dd, 1H), 2.87-2.79 (m, 1H), 2.32-2.21 (m, 1H), 2.00-1.86 (m, 2H), 1.83-1.64 (m, 1H); MS (EI) for C₉H₁₆N₂O₂: 185 (MH⁺).

Example 65

[0387] 1,2-Dideoxy-1-[(2*S*)-2-(methoxycarbonyl)-1-pyrrolidinyl]-2-[(phenylmethoxy)carbonyl]amino]-D-glycero-hexitol: To a solution of 2-deoxy-2-([(phenylmethoxy)carbonyl]amino)-D-glycero-hexopyranose (5.0 g, 0.016 mol) in methanol (500 mL) was added L-proline methyl ester hydrochloride (2.8 g, 0.022 mol) and sodium cyanoborohydride (3.4 g, 0.054 mol). The solution was heated to 64 °C for 14 h. After cooling to room temperature, the reaction mixture was concentrated *in vacuo* to afford 1,2-dideoxy-1-[(2*S*)-2-(methoxycarbonyl)-1-pyrrolidinyl]-2-[(phenylmethoxy)carbonyl]amino]-D-glycero-hexitol (6.81 g, 100%) as a clear and colorless oil. MS (EI) for C₂₀H₃₁N₂O₈: 427 (MH⁺).

Example 66

[0388] Methyl 1-[(2*S*)-3-hydroxy-2-([(phenylmethyl)oxy]carbonyl)amino]propyl]-L-prolinate: 1,2-dideoxy-1-[(2*S*)-2-(methoxycarbonyl)-1-pyrrolidinyl]-2-[(phenylmethoxy)carbonyl]amino]-D-glycero-hexitol (6.81 g, 0.016 mol) was taken into water (100 mL) and the resulting solution was cooled to 0°C. Sodium periodate (14.8 g, 0.069 mol) dissolved in water was added dropwise and the resulting mixture was stirred at 0°C for 2 h. The reaction mixture was partitioned with dichloromethane (3x100 mL), dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The residue was taken up in methanol (200 mL) and the resulting solution was cooled to 0°C. Sodium borohydride

(1.98 g, 0.052 mol) was added and the reaction mixture was stirred for 1 h at 0°C. The reaction mixture was concentrated *in vacuo* and partitioned with dichloromethane and saturated aqueous ammonium chloride. The organic layer was dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo*. The resulting crude product was purified by column chromatography (5% methanol in dichloromethane) to yield methyl 1-[(2*S*)-3-hydroxy-2-([(phenylmethyl)oxy]carbonyl)amino)propyl]-L-prolinate (4.9 g, 92%) as a white solid. MS (EI) for C₁₇H₂₅N₂O₅: 337 (MH⁺).

[0389] Methyl 1-[(2*S*)-3-[(methylsulfonyl)oxy]-2-([(phenylmethyl)oxy]carbonyl)amino)propyl]-L-prolinate: Methyl 1-[(2*S*)-3-hydroxy-2-([(phenylmethyl)oxy]carbonyl)amino)propyl]-L-prolinate (200 mg, 0.594 mmol) was dissolved in dichloromethane (5 mL) followed by the addition of 4-(dimethylamino)pyridine (3.6 mg, 0.039 mmol) and triethylamine (0.125 mL, 0.891 mmol) and the resulting mixture was cooled to 0 °C. Methanesulfonyl chloride (0.060 mL, 0.773 mmol) was added dropwise and the reaction mixture was stirred for 1 h at 0°C. The mixture was partitioned between dichloromethane and saturated aqueous sodium bicarbonate. The organic layer was dried over anhydrous magnesium sulfate, filtered and concentrated *in vacuo* to afford methyl 1-[(2*S*)-3-[(methylsulfonyl)oxy]-2-([(phenylmethyl)oxy]carbonyl)amino)propyl]-L-prolinate (246 mg, 100%) as a clear and colorless oil. MS (EI) for C₁₈H₂₇N₂O₇S: 415 (MH⁺).

Example 67

[0390] 1,1-Dimethylethyl (3*aR*,6*aS*)-5-(hydroxymethyl)hexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate: Under a nitrogen atmosphere, borane tetrahydrofuran complex (1M in THF, 42 mL, 41.9 mmol) was diluted with tetrahydrofuran (42 mL) and cooled with an ice bath. Neat 2,3-dimethylbut-2-ene (5.0 mL, 41.9 mmol) was added in portions over 0.25 h and the solution was stirred at 0°C for 3 h. A solution of 1,1-dimethylethyl (3*aR*,6*aS*)-5-methylidenehexahydrocyclopenta[*c*]pyrrole-2(1*H*)-carboxylate (1.98 g, 8.88 mmol) in tetrahydrofuran (10 mL) was added slowly, and the solution was warmed to room temperature and stirred 12 h. After cooling to 0°C, 10% aqueous sodium hydroxide (17 mL, 41.7 mmol) was added slowly, followed by 30% aqueous hydrogen peroxide (13 mL, 128 mmol) and the solution was warmed to room temperature. The solvent was removed *in vacuo* and the solution was partitioned between water and diethyl ether. The layers were separated and the aqueous layer was further extracted (3 x 50 mL diethyl ether). The

combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo* to provide 2.04 (95%) of 1,1-dimethylethyl (3aR,6aS)-5-(hydroxymethyl)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate, which was used without purification. ¹H NMR (400 MHz, CDCl₃): 8.50 (broad s, 1H), 3.66-3.46 (m, 3H), 3.20-3.00 (m, 2H), 2.70-2.59 (m, 2H), 2.37-2.18 (m, 1H), 2.04 (m, 1H), 1.84 (broad s, 1H), 1.70-1.55 (m, 1H), 1.46 (s, 9H), 1.17 (m, 1H), 0.93 (m, 1H).

[0391] 1,1-Dimethylethyl (3aR,6aS)-5-[(methylsulfonyl)oxy]methyl}hexahydrocyclopenta [c]pyrrole-2(1H)-carboxylate: Methanesulfonyl chloride (0.2mL, 2.48mmol), was added dropwise to a solution of 1,1-dimethylethyl (3aR,6aS)-5-(hydroxymethyl)hexahydrocyclopenta[c] pyrrole-2(1H)-carboxylate (0.40 g, 1.65 mmol) and triethylamine (0.69 mL, 4.95 mmol) in 20 mL dichloromethane at 0°C and the reaction mixture was stirred for 1 h at room temperature. The solvent was evaporated, the resulting crude mixture was diluted with 100 mL ethyl acetate and washed with water (30 mL), 1M aqueous sodium hydroxide, brine, 1M aqueous hydrochloric acid and brine again. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. The resulting 1,1-dimethylethyl (3aR,6aS)-5-[(methylsulfonyl)oxy]methyl}hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate was used without further purification. MS (EI) for C₁₄H₂₅NO₅S: 320 (MH⁺), 264 (M-tBu).

Example 68

[0392] 1,1-Dimethylethyl (3aR,6aS)-5-(hydroxy)-hexahydrocyclopenta[c] pyrrole-2(1H)-carboxylate: Sodium borohydride (0.15 g, 4.00 mmol), was added to a solution of 1,1-dimethylethyl (3aR,6aS)-5-oxo-hexahydrocyclopenta[c] pyrrole-2(1H)-carboxylate (0.45 g, 2.00 mmol) in 10 mL methanol at 0°C and the reaction mixture was stirred for 1 h at this temperature. The solvent was evaporated, the crude mixture was diluted with 100 mL ethyl acetate and washed with water (30 mL), 1M aqueous hydrochloric acid and brine. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated to give 1,1-dimethylethyl (3aR,6aS)-5-(hydroxy)-hexahydrocyclopenta[c] pyrrole-2(1H)-carboxylate (0.44g, 98%). ¹H NMR (400 MHz, d₆-DMSO): 4.08 (m, 1H), 3.40 (m, 2H), 3.30 (m, 2H), 2.50 (m, 2H), 1.98 (m, 2H), 1.40 (s, 9H), 1.30 (m, 2H). MS (EI) for C₁₂H₂₁NO₃: 228 (MH⁺).

[0393] 1,1-Dimethylethyl (3aR,6aS)-5-[[[(methylsulfonyl)oxy]]hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate: Methanesulfonyl chloride (0.18 mL, 2.33 mmol), was added dropwise to a solution of 1,1-dimethylethyl (3aR,6aS)-5-(hydroxy)-hexahydrocyclopenta[c] pyrrole-2(1H)-carboxylate (0.44 g, 1.94 mmol) and triethylamine (0.81 mL, 5.81 mmol) in 10 mL dichloromethane at 0°C and the reaction mixture was stirred for 1 h at room temperature. The solvent was evaporated, the resulting crude mixture was diluted with 100 mL ethyl acetate and washed with water (30 mL), brine, 1M aqueous hydrochloric acid and brine again. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated. The resulting crude 1,1-dimethylethyl (3aR,6aS)-5-[[[(methylsulfonyl)oxy]]hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate was used without further purification. MS (EI) for C₁₃H₂₃NO₅S: 306 (MH⁺).

Example 69

[0394] 3-(Chloromethyl)hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazine: A solution of (3R)-morpholin-3-ylmethanol (4.21 g, 36.0 mmol) in 2-(chloromethyl)oxirane (28.2 mL, 0.360 mol) was heated to 40°C for 3 h and then the solution was concentrated *in vacuo*. The intermediate was cooled in an ice bath and treated with 30.0 mL of concentrated sulfuric acid. The mixture was heated to 170°C for 2 h and then allowed to cool to room temperature. The mixture was poured into ice-water and solid sodium bicarbonate was carefully added until the solution was basic. 10% methanol in ethyl acetate was added and the biphasic mixture was filtered. The layers were separated and the aqueous layer was extracted (3 x 100 mL 10% methanol in ethyl acetate). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo*. Column chromatography (SiO₂, 2:5 hexanes:ethyl acetate) provided 3-(chloromethyl)hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazine 2.44g (35%) as two separated diastereomers. (3R,9aS)-3-(chloromethyl)hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazine: (0.886 g, 13% yield): ¹H NMR (400 MHz, CDCl₃): 3.91 (m, 3H), 3.82 (m, 1H), 3.68 (dt, 1H), 3.61 (dd, 1H), 3.47 (dd, 1H), 3.35 (t, 1H), 3.19 (t, 1H), 2.80 (d, 1H), 2.54 (m, 2H), 2.40 (m, 2H); MS (EI) for C₈H₁₄NO₂Cl: 192 (MH⁺). (3S,9aS)-3-(chloromethyl)hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazine: (1.55 g, 22% yield): ¹H NMR (400 MHz, CDCl₃): 3.85 (m, 2H), 3.73 (m, 3H), 3.50 (m, 2H), 3.29 (t, 1H), 3.18 (t, 1H), 2.85 (dd, 1H), 2.64 (dd, 1H), 2.40 (m, 2H), 2.17 (t, 1H); MS (EI) for C₈H₁₄NO₂Cl: 192 (MH⁺).

[0395] Hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl acetate: A suspension of (3R,9aS)-3-(chloromethyl)hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazine (1.97 g, 10.3 mmol) and potassium acetate (10.1 g, 102 mmol) in DMF (20.0 mL) was stirred at 140°C for 16 h, and then at 150°C for another 12 h. The reaction mixture was partitioned between water (250 mL) and ethyl acetate (250 mL), the organic layer was washed with 5% lithium chloride (2 x 100 mL) and brine (100 mL) then dried over anhydrous sodium sulfate and concentrated *in vacuo*. Column chromatography (SiO₂, 1:1 hexane:ethyl acetate, then 100% ethyl acetate) afforded 0.92 g (42%) of hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl acetate as a yellow oil. Distinct diastereomers as described above were converted in this step to give: (3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl acetate: ¹H NMR (400 MHz, CDCl₃): 4.18 (dd, 1H), 4.00 (m, 1H), 3.80 (dd, 1H), 3.68 (dt, 1H), 3.60 (dd, 1H), 3.46 (m, 2H), 3.22 (t, 1H), 2.64 (dd, 1H), 2.53 (m, 2H), 2.43-2.35 (m, 2H), 2.10 (s, 3H), and (3S,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl acetate: ¹H NMR (400 MHz, CDCl₃): 4.09 (d, 2H), 3.90-3.82 (m, 2H), 3.75-3.64 (m, 3H), 3.27 (t, 1H), 3.18 (t, 1H), 2.69 (dd, 1H), 2.63 (m, 1H), 2.46-2.33 (m, 2H), 2.16 (t, 1H), 2.10 (s, 3H).

[0396] (3R,9aS)-Hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl methanesulfonate: To a solution of (3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl acetate (0.922 g, 4.28 mmol) in methanol (14.0 mL) was added 1.03 mL (4.50 mmol) of sodium methoxide (25% wt. in methanol) dropwise at room temperature. After 5 min., 1.6 mL (6.43 mmol) of 4.0M hydrogen chloride in dioxane was added and a pink precipitate formed. The solution was concentrated *in vacuo* and the pink solid was taken up in 30.0 mL dichloromethane. This slurry was cooled in an ice bath and triethylamine (3.0 mL, 21.5 mmol) was added, followed by methanesulfonyl chloride (0.37 mL, 4.71 mmol). The resultant yellow solution was stirred for 30 minutes at room temperature. The mixture was then partitioned between dichloromethane and saturated aqueous sodium bicarbonate then the aqueous layer was extracted (3 x 50 mL dichloromethane). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo* to provide crude (3R,9aS)-hexahydro-1H-[1,4]oxazino[3,4-c][1,4]oxazin-3-ylmethyl methanesulfonate which was taken on to the following reaction without purification.


Example 70

[0397] (8aR)-6-(Chloromethyl)tetrahydro-1H-[1,3]thiazolo[4,3-c][1,4]oxazine: A solution of (4R)-1,3-thiazolidin-4-ylmethanol (0.300 g, 2.52 mmol) in 2-(chloromethyl)oxirane (2.0 mL, 25.5 mmol) was heated under nitrogen to 40°C for 12 h. The solution was then cooled to room temperature and 2-(chloromethyl)oxirane was removed *in vacuo*. The crude intermediate was cooled in ice, and was taken up in 2.0 mL of concentrated sulfuric acid. The resulting mixture was heated to 200°C for 0.5 h then poured carefully onto wet ice, which was allowed to melt. The aqueous solution was carefully made basic using solid sodium bicarbonate and the resulting mixture was filtered using water and 10% methanol in ethyl acetate as eluent. The layers were separated and the aqueous layer was extracted with 10% methanol in ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated *in vacuo* to give 11.6 mg (2.4% yield) of crude (8aR)-6-(chloromethyl)tetrahydro-1H-[1,3]thiazolo[4,3-c][1,4]oxazine as a mixture of diastereomers which was directly taken on to the next step.

Example 71

[0398] 1,1-Dimethylethyl (3-endo)-3-{2-[(methanesulfonyl)oxy]ethyl}-8-azabicyclo[3.2.1]octane-8-carboxylate: To a solution of 1,1-dimethylethyl (3-endo)-3-(2-hydroxyethyl)-8-azabicyclo[3.2.1]octane-8-carboxylate (30.3 mg, 1.19 mmol) in dichloromethane (4.0 mL), was added triethylamine (0.5 mL, 3.56 mmol) and the solution was cooled to 0°C under nitrogen. Methanesulfonyl chloride (0.11 mL, 1.42 mmol) was added slowly and mixture was allowed to warm to room temperature and stirred for 1h. The reaction mixture was partitioned between dichloromethane and water. The aqueous phase was extracted with dichloromethane (2 x 100 mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo* to provide 35.1 mg (89%) of 1,1-dimethylethyl (3-endo)-3-{2-[(methanesulfonyl)oxy]ethyl}-8-azabicyclo[3.2.1]octane-8-carboxylate, which was carried forward for alkylation without purification.

Example 72




[0399] N-[3-Fluoro-4-((7-(methyloxy)-6-((3-morpholin-4-ylpropyl)oxy)quinazolin-4-yl)oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. Crude cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(6-hydroxy-7-methoxy-quinazolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (333mg, 0.66mmol), PS-PPh₃ resin, (loading = 2.33mmol/g, 797mg, 1.86mmol), 3-morpholin-4-yl-propan-1-ol (0.26ml, 1.88mmol), and DEAD (0.31ml, 1.91mmol) were combined in CH₂Cl₂ (10ml) and stirred at room temperature for 1-2hrs. The reaction mixture was filtered and the resin thoroughly washed with CH₂Cl₂. The filtrate was concentrated *in vacuo* and the resulting residue was dissolved in EtOAc and washed with H₂O (4x) and sat'd NaCl (1x) and then extracted with 1N HCl (3x). The combined 1N HCl extractions were washed with EtOAc (2x). The acidic aqueous phase was then basified with 1N NaOH and extracted with EtOAc (3x). The combined EtOAc extractions were washed with H₂O (1x), sat'd NaCl (1x), dried (Na₂SO₄), and concentrated *in vacuo*. The resulting residue was purified by preparative reverse phase HPLC (25mM NH₄OAc/acetonitrile) and the pure fractions were lyophilized to give cyclopropane-1,1-dicarboxylic acid {3-fluoro-4-[7-methoxy-6-(3-morpholin-4-yl-propoxy)-quinazolin-4-yloxy]-phenyl}-amide (4-fluoro-phenyl)-amide (42.6mg, 10%). ¹H NMR (400MHz, DMSO-d₆): δ 10.37 (br s, 1H), 10.05 (br s, 1H), 8.55 (s, 1H), 7.84 (m, 1H), 7.65 (m, 2H), 7.58 (s, 1H), 7.43 (m, 3H), 7.16 (t, 2H), 4.27 (m, 2H), 4.00 (s, 3H), 3.60 (m, 6H), 2.39 (m, 4H), 1.99 (m, 2H), 1.47 (m, 4H). LC/MS Calcd for [M+H]⁺ 634.2, found 634.1.

[0400] Using the same or analogous synthetic techniques and/or substituting with alternative starting materials, the following were prepared:

[0401] N-{3-fluoro-4-[(7-(methyloxy)-6-[[1-methylpiperidin-4-yl)methyl]oxy]quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide: ¹H NMR (400MHz, CDCl₃): δ 9.67 (s, 1H), 8.59 (s, 1H), 8.43 (s, 1H), 7.75 (d, 1H), 7.52 (s, 1H),


7.46 (m, 2H), 7.31 (s, 1H), 7.20 (m, 2H), 7.06 (t, 2H), 4.04 (d, 2H), 4.03 (s, 3H), 2.98 (d, 2H), 2.34 (s, 3H), 2.12-2.1.95 (m, 5H), 1.76 (m, 2H), 1.64 (m, 2H), 1.57 (m, 2H).

Example 73



[0402] Preparation of 1-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropanecarboxylic acid. To the cyclopropyl di-carboxylic acid (449 mg, 3.45 mmol) in THF (3.5 mL) was added TEA (485 μ L, 3.45 mmol). The resulting solution was stirred at room temperature under a nitrogen atmosphere for 40 minutes before adding thionyl chloride (250 μ L, 3.44 mmol). The reaction was monitored by LCMS for the formation of mono acid chloride (quenched the sample with MeOH and looked for corresponding mono methyl ester). After 3 hours stirring at room temperature, 4-(6,7-dimethoxy-quinolin-4-yloxy)-phenylamine (1.02 g, 3.44 mmol) was added as a solid, followed by more THF (1.5 mL). Continued to stir at room temperature for 16 hours. The resulting thick slurry was diluted with EtOAc (10 mL) and extracted with 1N NaOH. The biphasic slurry was filtered and the aqueous phase was acidified with conc. HCl to pH = 6 and filtered. Both solids were combined and washed with EtOAc, then dried under vacuum. The desired product, 1-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropanecarboxylic acid, was obtained (962 mg, 68.7% yield, 97% pure) as a white solid. ^1H NMR ($\text{D}_2\text{O}/\text{NaOH}$): 7.97 (d, 1H), 7.18 (d, 2H), 6.76 (m, 4H), 6.08 (d, 1H), 3.73 (s, 3H), 3.56 (s, 3H), 1.15 (d, 4H).

Example 74



[0403] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[(4-fluorophenyl)methyl]cyclopropane-1,1-dicarboxamide. To a solution of 1-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropanecarboxylic acid (74.3 mg, 0.182 mmol), 4-Fluorobenzylamine (25 μ L, 0.219 mmol), DIEA (90.0 μ L, 0.544 mmol) in DMA (1.0 mL) was added HATU (203 mg, 0.534 mmol). The resulting solution was stirred at room temperature for 1 hour before adding dropwise to water (10 mL) with stirring. The slurry was sonicated, filtered and the solids were washed with 1 N NaOH followed by water. After air drying, the solids were further purified by prep HPLC, affording N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[(4-fluorophenyl)methyl]cyclopropane-1,1-dicarboxamide (33 mg, 35% yield, 98% pure) as a white solid. ^1H NMR (DMSO, d_6): 10.82 (s, 1H), 8.80 (d, 1H), 8.50 (t, 1H), 7.83 (d, 2H), 7.74 (s, 1H), 7.56 (s, 1H), 7.30-7.38 (m, 4H), 7.15 (t, 2H), 6.80 (d, 1H), 4.32 (d, 2H), 4.04 (s, 3H), 4.03 (s, 3H), 1.42 (s, 4H).

[0404] The following compounds were prepared, in a similar manner as above, from the coupling of 1-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropane carboxylic acid with a corresponding alkylamine or arylamine.

[0405] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide. ^1H NMR (DMSO- d_6): 10.62 (s, 1H), 8.79 (d, 1H), 8.24 (t, 1H), 7.83 (d, 2H), 7.72 (s, 1H), 7.58 (s, 1H), 7.37 (d, 2H), 6.76 (d, 1H), 4.04 (s, 3H), 4.03 (s, 3H), 3.98 (m, 2H), 3.66 (m, 2H), 3.49 (m, 4H), 3.25 (t, 2H), 3.13 (br., 2H), 1.42 (d, 4H).

[0406] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide. ^1H NMR (DMSO- d_6): 10.78 (s, 1H), 10.53 (s, 1H), 8.43 (d, 1H), 8.12 (d, 1H), 7.82 (d, 2H), 7.49 (s, 1H), 7.37 (s, 1H), 7.20-7.28 (m, 3H), 7.15 (dd, 1H), 7.01 (td, 1H), 6.35 (d, 1H), 3.93 (s, 3H), 3.92 (s, 3H), 3.47 (s, 2H), 2.17 (br., 4H), 1.49 (m, 4H), 1.41 (m, 4H), 1.32 (br., 2H).

[0407] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide. ^1H NMR (DMSO- d_6): 10.98 (s, 1H), 10.56 (s, 1H), 8.42 (d, 1H), 8.10 (dd, 1H), 7.81 (m, 2H), 7.49 (s, 1H), 7.37 (s, 1H), 7.17-7.27 (m, 4H), 7.01 (td, 1H), 6.35 (d, 1H), 3.93 (s, 3H), 3.92 (s, 3H), 3.61 (s, 2H), 2.30 (br., 4H), 1.47 (br., 4H), 1.43 (m, 4H).

[0408] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide. ^1H NMR (DMSO- d_6): 10.12 (s, 1H),


10.03 (s, 1H), 8.44 (d, 1H), 7.74 (d, 2H), 7.57 (s, 1H), 7.53 (d, 1H), 7.48 (s, 1H), 7.37 (s, 1H), 7.21 (m, 3H), 6.98 (d, 1H), 6.40 (d, 1H), 3.93 (s, 3H), 3.92 (s, 3H), 3.56 (t, 4H), 3.41 (s, 2H), 2.34 (br., 4H), 1.48 (s, 4H).


[0409] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide. ¹H NMR (DMSO-d₆): 10.54 (s, 1H), 10.47 (s, 1H), 8.43 (d, 1H), 8.08 (d, 1H), 7.78 (d, 2H), 7.49 (s, 1H), 7.37 (d, 1H), 7.18-7.30 (m, 4H), 7.03 (t, 1H), 6.37 (d, 1H), 3.94 (s, 3H), 3.93 (s, 3H), 3.50 (s, 2H), 3.44 (br., 4H), 2.20 (br., 4H), 1.48 (d, 4H).

[0410] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide. ¹H NMR (DMSO-d₆): 10.0-10.2 (br., 2H), 8.46 (d, 1H), 7.76 (d, 2H), 7.53 (m, 3H), 7.39 (s, 1H), 7.24 (m, 3H), 6.98 (d, 1H), 6.43 (d, 1H), 3.95 (s, 3H), 3.93 (s, 3H), 3.37 (s, 2H), 2.31 (br., 4H), 1.48 (m, 8H), 1.39 (br., 2H).

[0411] N-(4-{[6,7-Bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide. ¹H NMR (DMSO-d₆): 10.0-10.2 (br., 2H), 8.46 (d, 1H), 7.77 (d, 2H), 7.59 (s, 1H), 7.53 (d, 1H), 7.51 (s, 1H), 7.39 (s, 1H), 7.23 (m, 3H), 6.99 (d, 1H), 6.43 (d, 1H), 3.95 (s, 3H), 3.93 (s, 3H), 3.52 (s, 2H), 2.42 (br., 4H), 1.69 (br, 4H), 1.48 (s, 4H).

Example 75





[0412] Synthesis of N-(4-{[6,7-bis(methoxy)-2-(methylthio)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide Commercially


available 5-(bis-methylsulfanyl-methylene)-2,2-dimethyl-[1,3]dioxane-4,6-dione (3.5 g, 14 mmol) and 3,4-dimethoxyaniline (2.2 g, 14 mmol) were reflux in EtOH (20 mL) for 2 hours. The EtOH was removed under reduced pressure and EtOAc was added to the residue. The product was filtered and washed with cold EtOAc (3X). 5-[(3,4-dimethoxy-phenylamino)-methylsulfanyl-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione was obtained as a white solid (1.7 g, 47% yield) and used without further purification. LCMS: m/z 352 (M-H)⁻.

[0413] A mixture of 5-[(3,4-dimethoxy-phenylamino)-methylsulfanyl-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione (1.7 g, 6.6 mmol) and diphenylether (3.5 g, 21 mmol) were heated at 260 °C for 10 minutes. The mixture was cooled to room temperature and heptane was added. 6,7-Dimethoxy-2-methylsulfanyl-quinolin-4-ol was filtered and isolated as an orange solid and used without further purification (1.4 g, 83% yield). LCMS: m/z 352 (M+H)⁺.

[0414] A mixture of 6,7-dimethoxy-2-methylsulfanyl-quinolin-4-ol (1.0 g, 4.0 mmol), 3,4-difluoronitrobenzene (0.48 mL, 4.3 mmol), cesium carbonate (2.6 g, 8.0 mmol), and DMF (15 mL) was stirred at room temperature for 12 hours, after which time, the mixture was filtered. The filtrate was extracted with DCM, washed with 10% LiCl_(aq.), water, (1X) and brine (1X), followed by drying over Na₂SO₄ and concentration *in vacuo*. The crude solids were purified by flash chromatography (silica gel, 5% MeOH in DCM), affording the nitroquinoline (1.3 g, 85.8% yield) as an orange solid. LCMS: m/z 391 (M+H)⁺. A mixture of nitroquinoline (0.33 g, 0.85 mmol), 5% Pt/S on carbon (0.050 g), ammonium formate (0.40 g, 6.3 mmol) in EtOH (5 mL) was heated at 80 °C for 1 hour. The mixture was cooled to room temperature and the solvent removed under reduced pressure. The residue was dissolved in DCM, the mixture filtered, and the precipitate discarded. Removal of the organic solvent afforded 4-(6,7-dimethoxy-2-methylsulfanyl-quinolin-4-yl)oxy-3-fluoro-phenylamine as an orange oil (220 mg, 73% yield). LCMS: m/z 361 (M+H)⁺.

[0415] To a mixture of 4-(6,7-dimethoxy-2-methylsulfanyl-quinolin-4-yloxy)-3-fluorophenylamine (0.22 g, 0.61 mmol) and 1-(4-Fluorophenylcarbonyl)-cyclopropanecarboxylic acid (0.16 g, 0.73 mmol) in DMF (5 mL) was added TEA (0.25 mL, 1.8 mmol) followed by HATU (0.57 g, 1.5 mmol). The resulting solution was stirred overnight at room temperature. The reaction mixture was dumped into water and extracted with DCM (2X). The combined extracts were washed with 5% LiCl_(aq) (3X), water, (1X) and brine (1X), followed by drying over Na₂SO₄ and concentration *in vacuo*. The crude solids were purified by preparatory HPLC with ammonium acetate, affording N-(4-{[6,7-bis(methyloxy)-2-(methylthio)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide (0.39 g, 11% yield) as a white solid. ¹H NMR (DMSO-*d*₆) δ 10.34 (s, 1H), 9.94 (s, 1H), 7.83 (d, 1H), 7.59 (m, 2H), 7.56 (m, 1H), 7.40 (m, 2H), 7.23 (s, 1H), 7.09 (t, 2H), 6.12 (s, 1H), 3.88 (s, 3H), 3.85 (s, 3H), 2.48 (s, 3H), 1.40 (m, 4H).

Example 76




[0416] Synthesis of N-(4-{[2-amino-6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. A mixture of 5-[(3,4-dimethoxy-phenylamino)-methylsulfanyl-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione (1.0 g, 2.8 mmol), 30% ammonium hydroxide (8.5 mL), HgCl₂ (0.76 g, 2.8 mmol) in THF (5 mL) was stirred at room temperature for 30 minutes. The mixture was extracted with DCM and water (3X) and dried with Na₂SO₄. Concentration *in vacuo* afforded 5-[amino-(3,4-dimethoxy-phenylamino)-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione as a white solid (0.90 g, 97% yield) and this compound was used without further purification. LCMS: m/z 321 (M-H)⁻.

[0417] A mixture of 5-[amino-(3,4-dimethoxy-phenylamino)-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione (0.90 g, 2.8 mmol) and diphenylether (3.0 g, 18 mmol) was heated at 260 °C for 30 minutes. The mixture was cooled to room temperature and heptane was added. Product 2-amino-6,7-dimethoxy-quinolin-4-ol was filtered and isolated as an orange solid and used without further purification (0.31 g, 33% yield). LCMS: m/z 221 ($M+H$)⁺.

[0418] 4-(4-Amino-2-fluoro-phenoxy)-6,7-dimethoxy-quinolin-2-ylamine was synthesized from 2-amino-6,7-dimethoxy-quinolin-4-ol in a similar manner as 4-(6,7-dimethoxy-2-methylsulfanyl-quinolin-4-yloxy)-3-fluoro-phenylamine, and obtained as a white solid (4.0% yield). LCMS: m/z 330 ($M+H$)⁺.

[0419] N-(4-{[2-amino-6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide was synthesized from 4-(4-amino-2-fluoro-phenoxy)-6,7-dimethoxy-quinolin-2-ylamine in a similar manner as N-(4-{[6,7-bis(methyloxy)-2-(methylthio)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. It was purified by preparatory HPLC using ammonium acetate and isolated as a white solid (4.0% yield). ¹H NMR (DMSO-*d*₆) δ 10.34 (s, 1H), 9.95 (s, 1H), 7.82 (d, 1H), 7.58 (m, 2H), 7.44 (d, 1H), 7.33 (t, 1H), 7.25 (s, 1H), 7.09 (t, 2H), 7.07 (s, 1H), 6.17 (br s, 2H), 5.66 (s, 1H), 3.79 (s, 3H), 3.77 (s, 3H), 1.40 (d, 4H). LCMS: m/z 535 ($M+H$)⁺.

Example 77



[0420] Synthesis of 'N-(3-fluoro-4-{[2-(methylamino)-6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. A mixture of 5-


[(3,4-dimethoxy-phenylamino)-methylsulfanyl-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione (0.50 g, 1.4 mmol), methylamine (2 M in THF, 0.75 mL, 1.5 mmol), HgCl_2 (0.38 g, 1.4 mmol) in THF (5 mL) was stirred at room temperature for 30 minutes. The mixture was extracted with DCM and water (3X) and dried with Na_2SO_4 . Concentration *in vacuo* afforded 5-[(3,4-dimethoxy-phenylamino)-methylamino-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione as a yellow solid (0.48 g, 99% yield) and this compound was used without further purification. LCMS: m/z 335 ($\text{M}-\text{H}^-$).

[0421] A mixture of 5-[(3,4-dimethoxy-phenylamino)-methylamino-methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione (0.40 g, 2.8 mmol) and diphenylether (3.0 g, 18 mmol) was heated at 260°C for 15 minutes. The mixture was cooled to room temperature and heptane was added. Product 6,7-dimethoxy-2-methylamino-quinolin-4-ol was filtered and isolated as a tan solid and used without further purification (0.30 g, quantitative yield). LCMS: m/z 235 ($\text{M}+\text{H}^+$).

[0422] [4-(4-Amino-2-fluoro-phenoxy)-6,7-dimethoxy-quinolin-2-yl]-methyl-amine was synthesized from 6,7-dimethoxy-2-methylamino-quinolin-4-ol in a similar manner as 4-(4-Amino-2-fluoro-phenoxy)-6,7-dimethoxy-quinolin-2-ylamine, and isolated as a yellow oil (58% yield). LCMS: m/z 330 ($\text{M}+\text{H}^+$).


[0423] N-(3-fluoro-4-{[2-(methylamino)-6,7-bis(methoxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide was synthesized from [4-(4-amino-2-fluoro-phenoxy)-6,7-dimethoxy-quinolin-2-yl]-methyl-amine in a similar manner as N-(4-{[6,7-bis(methoxy)-2-(methylthio)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. It was purified by preparatory HPLC using ammonium acetate and isolated as a white solid (6.0 mg, 4.0% yield). ^1H NMR ($\text{DMSO}-d_6$) δ 10.42 (s, 1H), 9.91 (s, 1H), 7.88 (dd, 1H), 7.56 (m, 2H), 7.44 (m, 4H), 7.09 (t, 2H), 5.90 (s, 1H), 3.88 (s, 3H), 3.85 (s, 3H), 3.39 (br s, 1H), 2.92 (s, 3H), 1.41 (dt, 4H). LCMS: m/z 535 ($\text{M}+\text{H}^+$).

Example 78



[0424] 'N-(4-{[6-{[3-(diethylamino)propyl]oxy}-7-(methoxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. To a slurry of cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(6-hydroxy-7-methoxy-quinolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (0.12 g, 0.23 mmol), hydroxypropyldiethylamine (0.090 mL, 0.61 mmol), triphenylphosphine (0.20 g, 0.76 mmol) in DCM (10 mL) was added DIAD (0.17 mL, 0.86 mmol). The resulting mixture was stirred at room temperature for 12 hours, after which time, the solvent was removed under reduced pressure. The residue was extracted with EtOAc and 1N HCl (6X) and brine (1X) followed by drying with Na₂SO₄. Concentration of the organic fraction *in vacuo* afforded 'N-(4-{[6-{[3-(diethylamino)propyl]oxy}-7-(methoxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide as a yellow oil (0.18 g, wet, ca 95% purity by analytical HPLC). Further purification by preparatory HPLC using ammonium acetate afforded the product in 99% purity by analytical HPLC. LCMS: *m/z* 619 (M+H)⁺. ¹H NMR (DMSO-*d*₆) δ 10.37 (br s, 1H), 10.00 (s, 1H), 8.44 (d, 1H), 7.87 (d, 1H), 7.62 (m, 2H), 7.49 (m, 2H), 7.41 (m, 2H), 7.13 (t, 2H), 6.40 (d, 1H), 4.17 (t, 2H), 3.93 (s, 3H), 2.59 (t, 2H), 2.49 (m, 6H), 1.91 (m, 4H), 0.94 (t, 6H).

Example 79




- [0425] N-(4-fluorophenyl)-N'-(4-{[2-methyl-6,7-bis(methyloxy)quinazolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide. Commercially available 2-amino-4,5-dimethoxy-benzoic acid methyl ester (3g, 0.014 mol) and acetic anhydride (4.03 mL, 0.0426 mol) were heated in heptane at 100°C for 3 hours. After removal of heptane *in vacuo*, the crude product of 2-acetylamino-4,5-dimethoxy-benzoic acid methyl ester was obtained and used without further purification. LC/MS: m/z 254 (M+H).
- [0426] To the crude 2-acetylamino-4,5-dimethoxy-benzoic acid methyl ester obtained above was added ammonium acetate (7.98g, 0.104 mol) and acetic acid (10 mL). The mixture was heated at reflux in a pressure tube until the formation of the desired cyclization product, as indicated by LC/MS: m/z 221 (M+H). After cooling to RT, the reaction mixture was diluted with water, and extracted with EtOAc 3 times. The combined organic phase was basified with aq. NaOH solution, and washed 3 times with EtOAc. The aqueous layer was then acidified with aq. HCl and extracted three times with EtOAc. The combined organic extract was dried over Na₂SO₄ and concentrated in vacuo, affording 6,7-dimethoxy-2-methyl-quinazolin-4-ol (0.15g), which was used without further purification. LC/MS: m/z 221 (M+H).
- [0427] A mixture of 6,7-dimethoxy-2-methyl-quinazolin-4-ol obtained from previous step (0.15g, 0.68 mmol) and POCl₃ (1.59 mL, 17.04 mmol) was heated at reflux for 48 hours. The reaction mixture was poured into ice water, neutralized with NaHCO₃, and adjusted to basic with K₂CO₃. The mixture was cooled to 0°C with stirring. The resulting precipitate was filtered, giving 4-chloro-6,7-dimethoxy-2-methyl-quinazoline (0.094g), which was used without further purification.
- [0428] A mixture of the chloro quinazoline (0.094g, 0.397 mmol) obtained above, 4-nitrophenol (0.11g, 0.795 mmol) and bromobenzene (3mL) was heated at 160°C for 48 hours. The solvent was then removed and the reaction was taken up in MeOH. Et₂O was added and the reaction stirred 30 min and the precipitate was filtered, affording 6,7-dimethoxy-2-methyl-4-(4-nitro-phenoxy)-quinazoline (0.081g) as a very light yellow solid. LC/MS: m/z 342 (M+H).
- [0429] A mixture of 6,7-dimethoxy-2-methyl-4-(4-nitro-phenoxy)-quinazoline (0.081g, 0.236 mmole), Pt/S (0.008g, 15 mol%), ammonium formate (0.098g, 1.56 mmol) and EtOH (3mL) was heated with stirring at 70°C for 3 hours. The reaction mixture was then filtered while hot and washed with hot EtOH. The crude product of 4-(6,7-dimethoxy-2-

methyl-quinazolin-4-yloxy)-phenylamine (0.924g) was obtained as a yellow solid, which was used in the next reaction without further purification. LC/MS: m/z 312 (M+H).

[0430] To a mixture of 4-(6,7-dimethoxy-2-methyl-quinazolin-4-yloxy)-phenylamine (0.100g, 0.321 mmol) and 1-(4-fluoro-phenylcarbamoyl)-cyclopropanecarboxylic acid (0.056g, 0.386 mmol) in DMF was added DIEA (0.168 mL, 0.963 mmol), followed by HATU (0.183g, 0.482 mmol). The reaction mixture was stirred at RT for 15 hours. The mixture was diluted with EtOAc, washed with 5% LiCl aq solution three times, dried over Na₂SO₄, and concentrated *in vacuo*. The crude product was purified on preparative HPLC to give 'N-(4-fluorophenyl)-N'-(4-{[2-methyl-6,7-bis(methyloxy)quinazolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide (3.2 mg) as a white solid. ¹H NMR (DMSO-d₆) 10.15 (bs, 1H), 10.01 (bs, 1H), 7.69-7.75 (m, 2H), 7.61-7.68 (m, 2H); 7.52 (s, 1H), 7.32 (s, 1H), 7.23-7.29 (m, 2H), 7.12-7.19 (m, 2H), 3.93 (d, 6H), 2.43 (s, 3H), 1.53 (s, 4H).


Example 80



[0431] Preparation of 1-(4-Fluoro-phenylcarbamoyl)-2-methyl-cyclopropanecarboxylic acid. 2-Methylcyclopropane-1,1-dicarboxylic acid methyl ester was prepared by following the literature procedure (Baldwin, J. E.; Adlington, R. M.; Rawlings, B. J. Tetrahedron Lett. 1985, 481.) The carboxylic acid (700 mg, 4.4 mmol) was dissolved in CH₂Cl₂ (10 mL). To the resulting solution was added 4-fluoroaniline (590 mg, 5.3 mmol), HOBT (890 mg, 6.6 mmol) and EDCI (2.5 g, 13.2 mmol). The stirring was continued for 3 h at rt. CH₂Cl₂ (30 mL) was added to the reaction mixture, and the resulting solution was washed with brine, and dried over Na₂SO₄. CH₂Cl₂ was removed under reduced pressure. Further purification by column chromatography gave 635 mg (57%) of the desired amide.

[0432] The methyl ester obtained above was then treated with $\text{LiOH} \cdot \text{H}_2\text{O}$ (116 mg, 2.78 mmol, 1.1 equiv.) in THF (2 mL) and H_2O (1 mL) for 3h at rt. THF was removed under reduced pressure. The aqueous solution was diluted with 20 mL of H_2O , washed with ether (10 mL), and acidified with 1 N HCl. The solid was filtered, dissolved in EtOAc, and dried over Na_2SO_4 . Removal of EtOAc gave the crude product of 1-(4-fluorophenylcarbamoyl)-2-methyl-cyclopropane-carboxylic acid, which was used in the next reaction. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.99 (br s, 1 H), 10.33 (br s, 1 H), 7.59 (dd, J = 9.0, 5.0 Hz, 2 H), 7.11 (dd, J = 9.0, 9.0 Hz, 2 H), 1.86-1.78 (m, 1 H), 1.43 (dd, J = 9.0, 4.2 Hz, 1 H), 1.30 (dd, J = 7.8, 4.3 Hz, 1 H), 1.19 (d, J = 6.3 Hz, 3 H).

Example 81




[0433] Synthesis of (1S,2R)-N-[3-fluoro-4-((6-(methoxy)-7-[(3-morpholin-4-yl)propyl]oxy)quinolin-4-yl)oxy]phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide. To a solution of 4-(7-benzyloxy-6-methoxyquinolin-4-yloxy)-3-fluorophenylamine (150 mg, 0.38 mmol) in CH_2Cl_2 (3 mL) was added DIEA (341 mg, 2.64 mmol), 1-(4-fluorophenylcarbamoyl)-2-methyl-cyclopropanecarboxylic acid (120 mg, 0.49 mmol) and PyBOP (686 mg, 1.32 mmol). The reaction mixture was stirred at rt for 6 h. After standard workup, the crude product was purified by column chromatography.

[0434] The coupling product (130 mg, 0.21 mmol) obtained above was dissolved in EtOH (2 mL). 1,4-cyclohexadiene (170 mg, 2.1 mmol) and 10% Pd/C (10 mg) were added. The

mixture was stirred for 2 h under reflux. After cooling, the mixture was filtered through Celite, and washed with MeOH. Removal of the solvents gave the crude product (136 mg), which was used in the next reaction.

[0435] To a solution of the 7-hydroxyquinoline (136 mg, 0.26 mmol) in DMF (2 mL) was added 4-(3-chloropropyl)morpholine hydrochloride (70 mg, 0.35 mmol) and K_2CO_3 (69 mg, 0.50 mmol). The reaction mixture was then stirred at 80 °C for 5 h. After cooling, EtOAc (20 mL) was added. The EtOAc solution was washed twice with brine, and dried over Na_2SO_4 . Removal of EtOAc and purification by column chromatography (CH_2Cl_2 : MeOH = 10:1) gave '(1S,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide. The product was then dissolved in ethyl ether, and treated with 1.5 equiv. of 1 N HCl/ether. Filtration and lyophilization gave the HCl salt of '(1S,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide: 1H NMR (400 MHz, $DMSO-d_6$) δ 10.49 (br s, 1 H), 10.26 (br s, 1 H), 10.15 (br s, 1 H), 8.74 (br s, 1 H), 7.95 (br d, J = 13.2 Hz, 1 H), 7.8-7.5 (m, 6 H), 7.16 (t, J = 8.9 Hz, 2 H), 6.82 (br s, 1 H), 4.34 (t, J = 5.9 Hz, 2 H), 4.02 (s, 3 H), 3.99 (br s, 2 H), 3.77 (br t, J = 12.0 Hz, 2 H), 3.56-3.30 (m, 4 H), 3.17-3.07 (m, 2 H), 2.40-2.30 (m, 2 H), 2.04-1.95 (m, 1 H), 1.45 (dd, J = 7.2, 4.7 Hz, 1 H), 1.36 (dd, J = 8.5, 4.5 Hz, 1 H), 1.09 (d, J = 6.2 Hz, 3 H).


Example 82



- [0436] Synthesis of (1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide. To a solution of 4-(7-benzyloxy-6-methoxy-quinolin-4-yloxy)-3-fluorophenylamine (322 mg, 0.82 mmol) and 2-Methyl-cyclopropane-1,1-dicarboxylic acid methyl ester (195 mg, 1.23 mmol) in CH₂Cl₂ (4 mL) was added HOBt (61 mg, 0.32 mmol) and EDCI (211 mg, 1.64 mmol). The stirring was continued for 12 h at rt. The reaction mixture was then diluted with EtOAc and washed with brine. Removal of organic solvents *in vacuo* and further purification by column chromatography gave the desired coupling product (153 mg).
- [0437] The product (153 mg, 0.29 mmol) obtained above was treated with LiOH•H₂O (15 mg, 0.35 mmol) in THF (1 mL) and H₂O (1 mL) for 2 h. THF was removed. 10 mL of H₂O was added to the mixture. The aqueous solution was washed with ether, and acidified with 1 N HCl. The solid was then filtered and dried under vacuum.
- [0438] The crude carboxylic acid (118 mg, 0.23 mmol) and 4-fluoroaniline (111 mg, 0.27 mmol) were dissolved in DMF (2 mL). To this solution was added DIEA (178 mg, 1.38 mmol) and PyBOP (358 mg, 0.69 mmol). The mixture was stirred overnight at rt. It was then diluted with EtOAc, washed twice with brine. Removal of EtOAc and column chromatography gave the desired product.
- [0439] The product (66 mg, 0.11 mmol) obtained above was dissolved in EtOH (2 mL). 1,4-cyclohexadiene (80 mg, 1.1 mmol) and 10% Pd/C (10 mg) were added. The mixture was stirred for 2 h under reflux. After cooling, the mixture was filtered through Celite, and washed with MeOH. Removal of the solvents gave the crude product (70 mg), which was used in the next reaction.
- [0440] To a solution of the 7-hydroxyquinoline (80 mg, 0.15 mmol) in DMF (2 mL) was added 4-(3-chloropropyl)morpholine hydrochloride (62 mg, 0.31 mmol) and K₂CO₃ (64 mg, 0.46 mmol). The reaction mixture was then stirred at 80 °C for 5 h. After cooling, EtOAc (20 mL) was added. The EtOAc solution was washed twice with brine, and dried over Na₂SO₄. Removal of EtOAc and purification by column chromatography (CH₂Cl₂ : MeOH = 10:1) gave '(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide. The product was then dissolved in ethyl ether, and treated with 1.5 equiv. of 1 N HCl/ether. Filtration and lyophilization gave the HCl salt of '(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-

fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide: ^1H NMR (400 MHz, DMSO- d_6) δ 10.65 (br s, 1 H), 10.54 (br s, 1 H), 9.74 (s, 1 H), 8.75 (br s, 1 H), 8.01 (br d, $J = 12.9$ Hz, 1 H), 7.80-7.50 (m, 6 H), 7.20-7.10 (m, 2 H), 6.84 (br s, 1 H), 4.34 (br t, $J = 5$ Hz, 2 H), 4.04 (s, 3 H), 4.05-3.95 (m, 2 H), 3.77 (br t, $J = 11$ Hz, 2 H), 3.52 (br d, $J = 12.7$ Hz, 4 H), 3.12 (br q, $J = 9.0$ Hz, 2 H), 2.40-2.30 (m, 2 H), 2.10-1.95 (m, 1 H), 1.40-1.30 (m, 2 H), 1.10 (d, $J = 6.2$ Hz, 3 H).

Example 83



[0441] Synthesis of '(2R,3R)-N-[3-fluoro-4-({6-(methoxy)-7-[(3-morpholin-4-ylpropyl)oxy]-quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide. 2,3-trans-Dimethyl-cyclopropane-1,1-dicarboxylic acid diethyl ester was prepared by following the literature procedure. (Ohishi, J. Synthesis, 1980, 690.) To a solution of 2,3-trans-dimethyl-cyclopropane-1,1-dicarboxylic acid diethyl ester (6.75 g, 31.5 mmol) in MeOH (30 mL) was added 33 mL of 1 N NaOH aqueous solution. The mixture was stirred at 85 °C for 5 h. MeOH was removed under reduced pressure; the residue was diluted with 40 mL of H₂O. The aqueous solution was washed with 20 mL of


ether, and acidified with 1 N HCl. Filtration and drying under vacuum gave 4.72 g 80 %) of the desired carboxylic acid.

[0442] The aniline (1.08 g, 2.78 mmol) and the carboxylic acid (518 mg, 2.78 mmol) prepared above were dissolved in CH_2Cl_2 (15 mL). HATU (2.11 g, 5.56 mmol) and DIEA (1.8 mL, 11.1 mmol) were added. The reaction mixture was stirred at rt overnight. It was then concentrated and diluted with EtOAc. The EtOAc solution was then washed with 5% NaOH and brine. Removal of EtOAc gave the crude coupling product, which was hydrolyzed to the corresponding carboxylic acid by treatment with $\text{LiOH}\cdot\text{H}_2\text{O}$ (175 mg, 4.17 mmol) in THF (100 mL) - H_2O (50 mL) at 60 °C for 10 h.

[0443] The carboxylic acid (850 mg, 1.60 mmol) and 4-fluoroaniline (355 mg, 3.20 mmol) were dissolved in DMF (8 mL). HATU (3.89 g, 3.2 mmol) and DIEA (1.1 mL, 6.4 mmol) were added. The reaction mixture was stirred at rt overnight. H_2O (10 mL) was added to the reaction, and a precipitate formed. The solid was filtered, washed with aqueous sat. Na_2CO_3 and ether. Further purification by column chromatography gave 596 mg (60%) of the desired product. Debenzylation was done by following the standard procedure.

[0444] To a solution of the 7-hydroxyquinoline (261 mg, 0.49 mmol) in DMF (5 mL) was added 4-(3-chloropropyl)morpholine hydrochloride (195 mg, 0.98 mmol) and K_2CO_3 (202 mg, 1.46 mmol). The reaction mixture was then stirred at 80 °C for 4 h. After cooling, EtOAc (20 mL) was added. The EtOAc solution was washed twice with brine, and dried over Na_2SO_4 . Removal of EtOAc and purification by column chromatography (CH_2Cl_2 : MeOH = 10:1) gave 122 mg (37%) of '(2R,3R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide. ^1H NMR (400 MHz, CDCl_3) δ 8.44 (d, J = 5.1 Hz, 1 H), 8.11 (br s, 1 H), 7.77-7.70 (m, 2 H), 7.53 (s, 1 H), 7.50-7.44 (m, 2 H), 7.40 (s, 1 H), 7.22-7.16 (m, 2 H), 7.06-6.98 (m, 2 H), 6.36 (br d, J = 5.1 Hz, 1 H), 4.26 (t, J = 7.0 Hz, 2 H), 4.02 (s, 3 H), 3.72 (t, J = 4.4 Hz, 4 H), 2.57 (t, J = 7.3 Hz, 2 H), 2.50-2.42 (m, 4 H), 2.18-2.10 (m, 2 H), 1.80-1.66 (m, 2 H), 1.30-1.24 (m, 6 H).

Example 84




[0445] Synthesis of 'N-(4-{{[6,7-bis(methoxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)-1-(phenylmethyl)azetidine-3,3-dicarboxamide.

1-Benzyl-azetidine-3,3-dicarboxylic acid was prepared by following the literature procedure (Miller, R. A.; et al. Syn. Comm. 2003, 33, 3347). To a solution of 4-(6,7-dimethoxy-quinolin-4-yloxy)-phenylamine (4.2 mmol, 1 equiv.) and 4-fluoroaniline (4.2 mmol, 1 equiv.) in DMF (20 mL) was charged with DIEA (12.6 mmol, 3 equiv.) and a solution of 1-benzyl-azetidine-3,3-dicarboxylic acid (4.2 mmol, 1 equiv.) in DMF (10 mL). The reaction mixture was allowed to stir at RT and monitored by LCMS. The reaction was complete in 6 h. The reaction mixture was diluted with ethyl acetate and washed with 10% LiCl (3x), brine (3x), dried with sodium sulfate, filtered and the solvent was reduced *in vacuo*. The crude product was purified by silica gel chromatography eluting with 2% of MeOH in EtOAc. The fractions containing the desired product were further purified using preparative HPLC to give 'N-(4-{{[6,7-bis(methoxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)-1-(phenylmethyl)azetidine-3,3-dicarboxamide (300 mg, 12% yield) as a white solid. ¹HNMR (DMSO-d₆): 10.0 (s, 1H), 9.90 (s, 1H), 8.45 (d, 1H), 7.80 (d, 2H), 7.70 (m, 2H), 7.50 (s, 1H), 7.40 (s, 1H), 7.48-7.15 (m, 9H), 3.95 (s, 6H), 3.70 (s, 4H), 3.60 (s, 2H). LCMS (POS): 607.2 (M+H).

[0446] N-(4-{{[6,7-bis(methoxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)azetidine-3,3-dicarboxamide. To a solution of 'N-(4-{{[6,7-bis(methoxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)-1-(phenylmethyl)azetidine-3,3-dicarboxamide (300 mg, 0.5 mmol) in MeOH (50 mL) was charged with Pd/C (50% wet, 10% mmol, 265 mg) and acetic acid (2 mL). The reaction mixture was subjected to hydrogenolysis condition under H₂ (50 psi) on a Parr Hydrogenator for 16 hr. The reaction mixture was filtered through celite and washed with MeOH. After removal of solvent in *vacuo*, the crude product was purified using preparative HPLC (solvent system: MeCN/H₂O/NH₄OAc), affording N-(4-{{[6,7-bis(methoxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)


azetidine-3,3-dicarboxamide (82 mg, 32% yield) as a white solid. ^1H NMR (DMSO- d_6): 8.46 (d, 1H), 7.84 (d, 2H), 7.70 (m, 2H), 7.50 (s, 1H), 7.40 (s, 1H), 7.24 (d, 2H), 7.20 (t, 2H), 6.44 (d, 1H), 4.03 (s, 4H), 3.95 (s, 6H), 1.90 (s, 3H, acetate salt). LCMS (POS): 517.3 (M+H).

Example 85




[0447] N-{3-fluoro-4-[(6-(methoxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide. To a solution of cyclopropane-1,1-dicarboxylic acid {3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yl]oxy}phenyl}-amide (4-fluoro-phenyl)-amide, TFA salt (~500 mg, 0.71 mmol) in $\text{ClCH}_2\text{CH}_2\text{Cl}$ (8 mL) were added 30% formaldehyde (4 mL) and $\text{NaBH}(\text{OAc})_3$ (752 mg, 3.55 mmol). The reaction mixture was stirred overnight. It was then quenched with aqueous sat. NaHCO_3 , extracted with EtOAc. The organic phase was washed with brine and dried over Na_2SO_4 . Drying salts were filtered, washed with EtOAc and the filtrate concentrated in vacuo to give 210mg of crude product. The resulting residue was redissolved in EtOAc and any insoluble material filtered. To the filtrate was added 4M HCl in dioxane (200 μL) and the mixture was stirred at room temperature for 1 hour. Solids were filtered, washed with EtOAc, dried under high vacuum, dissolved in 50% aqueous AcCN and lyophilized to give 'N-{3-fluoro-4-[(6-(methoxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide, HCl salt (113 mg, ~25% yield). ^1H NMR (400MHz, DMSO- d_6): δ 10.51 (s, 1H), 10.30 (br. s, 1H), 10.04 (s, 1H), 8.80 (d, 1H), 7.99 (dd, 1H), 7.55 (m, 2H), 7.67-7.53 (m, 4H), 7.16 (t, 2H), 6.89 (d, 1H), 4.13 (d, 2H), 4.05 (s, 3H), 3.47 (m, 2H), 3.00 (m, 2H), 2.74 (d, 3H), 2.17 (m, 1H), 2.03 (m, 2H), 1.68 (m, 2H), 1.49 (m, 4H). LC/MS Calcd for $[\text{M}+\text{H}]^+$ 617.3, found 617.4. Anal. HPLC (8 min gradient): 98% pure, 3.11 min.

Example 86




[0448] (1R,2R,3S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methoxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide. 2,3-Dimethyl-cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxy-quinolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (210 mg, 0.39mmol), DMA (2mls), (2-chloro-ethyl)-diethyl-amine, HCl salt (73 mg, 0.42mmol) and K_2CO_3 (136 mg, 0.98mmol) were combined and heated at 80C overnight. The reaction mixture was then diluted with H_2O and sonicated. The resulting solids were filtered, washed with H_2O and dried under high vacuum. The crude product was then purified by preparative HPLC using an ammonium acetate buffer system and lyophilized to give '(1R,2R,3S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methoxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide (39 mg, 16% yield). ^1H NMR (400MHz, DMSO-d_6): δ 10.14 (s, 1H), 9.61 (s, 1H), 8.46 (d, 1H), 7.87 (dd, 1H), 7.67 (m, 2H), 7.57 (m, 1H), 7.51 (s, 1H), 7.42 (s, 1H), 7.39 (m, 1H), 7.15 (t, 2H), 6.41 (d, 1H), 4.20 (m, 2H), 3.94 (s, 3H), 2.87 (m, 2H), 2.60 (m, 4H), 1.80 (m, 2H), 1.18 (s, 3H), 1.17 (s, 3H), 1.01 (m, 6H). Note: 0.5eq of AcOH is present by NMR. LC/MS Calcd for $[\text{M}+\text{H}]^+$ 633.3, found 633.4. Anal. HPLC (25 min gradient): 96% pure, 18.52 min.

Example 87



[0449] N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide. 2,2-Dimethyl-cyclopropane-1,1-dicarboxylic acid [3-fluoro-4-(7-hydroxy-6-methoxyquinazolin-4-yloxy)-phenyl]-amide (4-fluoro-phenyl)-amide (203 mg, 0.38mmol), DMA (2mls), (2-chloro-ethyl)-diethyl-amine, HCl salt (73 mg, 0.42mmol) and K₂CO₃ (146 mg, 1.05mmol) were combined and heated at 80C overnight. The reaction mixture was then diluted with H₂O and extracted with CH₂Cl₂ (3x). The combined CH₂Cl₂ extractions were washed with sat'd NaHCO₃ (1x), sat'd NaCl (1x), dried (Na₂SO₄), and concentrated in vacuo. The resulting crude product was purified by flash chromatography (Silica Gel 60, 100% EtOAc, followed by 10% MeOH, 1% triethylamine in EtOAc), then dissolved in 50% aqueous AcCN and lyophilized to give N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide (70 mg, 29% yield). ¹HNMR (400MHz, DMSO): δ 10.24 (s, 1H), 10.00 (s, 1H), 8.54 (s, 1H), 7.84 (dd, 1H), 7.66 (m, 2H), 7.56 (s, 1H), 7.51 (m, 1H), 7.43 (m, 2H), 7.18 (t, 2H), 4.26 (m, 2H), 3.98 (s, 3H), 2.88 (m, 2H), 2.59 (m, 4H), 1.58 (m, 2H), 1.18 (s, 6H), 1.00 (t, 6H). LC/MS Calcd for [M+H]⁺ 634.3, found 634.4. Anal. HPLC (25 min gradient): 94% pure, 24.08 min.

Example 88



[0450] Synthesis of N-[3-(aminomethyl)phenyl]-N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide. (4-Nitro-benzyl)-carbamic acid tert-butyl ester. 4-Nitro-benzylamine, HCl salt (5.19g, 27.5mmol) was dissolved in dioxane (100mls). NaOH (3.4g, 85.0mmol) in H₂O (20mls) was added, followed by Boc

anhydride (7.6g, 34.8mmol). The mixture was stirred at room temperature. After 3hrs, the reaction mixture was diluted with EtOAc and washed with H₂O (3x), sat'd NaCl (1x), dried (Na₂SO₄), and concentrated in vacuo. The resulting residue was triturated with hexanes, the resulting solids filtered, washed with hexanes and dried under vacuum to give (4-nitro-benzyl)-carbamic acid tert-butyl ester (6.34g, 91% yield). LC/MS Calcd for [M+H]⁺ 253.1, found 197.0 (minus t-butyl).

[0451] (4-Amino-benzyl)-carbamic acid tert-butyl ester. (4-Nitro-benzyl)-carbamic acid tert-butyl ester (6.34g, 25.1mmol), iron powder (6.5g, 116mmol), ammonium formate (13.0g, 206mmol), H₂O (75mls), and toluene (75mls) were combined and heated to reflux. After 3hrs the reaction mixture was allowed to cool and filtered through Celite with thorough washing with EtOAc. The filtrate was transferred to a separatory funnel and the phases separated. The organic phase was further washed with H₂O (1x), sat'd NaCl (1x), dried (Na₂SO₄), and concentrated in vacuo to give (4-amino-benzyl)-carbamic acid tert-butyl ester (5.02g, 90% yield). LC/MS Calcd for [M+H]⁺ 223.1, found 167.1 (minus t-butyl).

[0452] [3-({1-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropanecarbonyl}-amino)-benzyl]-carbamic acid tert-butyl ester. 1-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropanecarboxylic acid (254 mg, 0.62mmol), (4-amino-benzyl)-carbamic acid tert-butyl ester (164 mg, 0.74mmol), dry DMA (10mls), HATU (714 mg, 1.88mmol), and DIEA (325ml, 1.86mmol) were combined and stirred at room temperature. After 2hrs, the reaction mixture is diluted with H₂O and the resulting solids are filtered, washed with H₂O, followed by sat'd NaHCO₃, and dried under high vacuum to give crude [3-({1-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropanecarbonyl}-amino)-benzyl]-carbamic acid tert-butyl ester (301 mg, 79% yield) which was used in the next reaction without further purification. LC/MS Calcd for [M+H]⁺ 613.3, found 613.1.

[0453] N-[3-(Aminomethyl)phenyl]-N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide, TFA salt. [3-({1-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-phenylcarbamoyl]-cyclopropanecarbonyl}-amino)-benzyl]-carbamic acid tert-butyl ester (50 mg, 0.081mmol) was dissolved in 50% TFA in CH₂Cl₂ (10mls) and stirred at room temperature. After 2hrs, the reaction mixture was concentrated in vacuo and the resulting residue was triturated with Et₂O. The resulting solids were filtered, washed with Et₂O and dried under high vacuum to give 'N-[3-(aminomethyl)phenyl]-N'-(4-{[6,7-

bis(methyloxy)quinolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide as the TFA salt (54 mg, 100%). ¹H NMR (400MHz, DMSO-d₆): δ 10.28 (s, 1H), 10.19 (s, 1H), 8.77 (m, 1H), 8.21 (m, 3H), 7.84 (m, 2H), 7.76 (m, 1H), 7.71 (m, 1H), 7.58 (m, 2H), 7.38 (m, 3H), 7.19 (m, 1H), 6.76 (m, 1H), 4.03 (s, 6H), 3.39 (m, 2H), 1.53 (m, 4H). Note: all peaks are very broad and unresolved. LC/MS Calcd for [M+H]⁺ 513.2, found 513.4. Anal. HPLC (25 min gradient): 88% pure, 12.39 min.

[0454] Table 3 contains ¹H-NMR data for selected compounds of the invention.

Table 3

Entry	Name	¹ H-NMR
1	N-[3-fluoro-4-((6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.52 (s, 1H), 10.02 (s, 1H), 9.38 (br., 3H), 8.79 (d, 1H), 7.98 (dd, 1H), 7.74 (s, 1H), 7.65 (m, 3H), 7.54 (m, 2H), 7.15 (t, 2H), 6.86 (d, 1H), 4.33 (t, 2H), 4.04 (s, 3H), 3.17-3.50 (m, 9H), 2.27 (br., 2H), 1.79 (m, 1H), 1.48 (m, 4H). Note: The peak at δ9.38 includes 2 TFA equivalents.
2	N-{3-fluoro-4-[(6-(methyloxy)-7-[[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.41 (s, 1H), 10.03 (s, 1H), 8.47 (d, 1H), 7.90 (dd, 1H), 7.64 (m, 2H), 7.52 (s, 2H), 7.42 (t, 1H), 7.39 (s, 1H), 7.16 (t, 2H), 6.41 (d, 1H), 4.18 (t, 2H), 3.95 (s, 3H), 2.47 (t, 2H), 2.6-2.8 (br., 8H), 2.17 (s, 3H), 1.97 (m, 2H), 1.48 (s, 4H).
3	N-{3-fluoro-4-[(6-(methyloxy)-7-[[1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (400MHz, DMSO-d ₆): d 10.51 (s, 1H), 10.30 (br. s, 1H), 10.04 (s, 1H), 8.80 (d, 1H), 7.99 (dd, 1H), 7.55 (m, 2H), 7.67-7.53 (m, 4H), 7.16 (t, 2H), 6.89 (d, 1H), 4.13 (d, 2H), 4.05 (s, 3H), 3.47 (m, 2H), 3.00 (m, 2H), 2.74 (d, 3H), 2.17 (m, 1H), 2.03 (m, 2H), 1.68 (m, 2H), 1.49 (m, 4H).
4	N-(4-fluorophenyl)-N'-[4-((6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl)oxy)phenyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆): 8.47 (d, 1H), 8.30 (m, 1H), 8.15 (m, 1H), 7.8 (m, 2H), 7.62 (s, 1H), 7.45 (m, 2H), 7.2 (m, 3H), 7.10 (m, 2H), 6.7 (d, 1H), 4.5 (m, 2H), 4.3 (m, 2H), 4.01(s, 3H), 3.5 (br, 2H), 3.3 (m, 2H), 3.1 (m, 2H), 2.51 (m, 2H), 1.9 (m, 2H) 1.6 (m, 4H).
5	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆): 10.58 (s, 1H), 10.31 (bs, 1H), 10.04 (s, 1H), 8.75 (d, 1H), 7.99 (d, 1H), 7.74 (s, 1H), 7.63 (m, 4H), 7.19 (t, 2H), 6.91 (m, 1H), 4.39 (t, 2H), 4.19 (s, 3H), 3.21 (m, 7H), 2.29 (m, 2H), 1.46 (d, 4H), 1.15 (t, 6H).

Table 3

Entry	Name	¹ H-NMR
6	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-chloro-5-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 11.56 (s, 1H), 9.77 (s, 1H), 8.50 (d, 1H), 8.32 (d, 1H), 7.82 (d, 1H), 7.59 (m, 2H), 7.51 (s, 1H), 7.42 (s, 1H), 7.20 (t, 2H), 6.55 (d, 1H), 3.95 (s, 3H), 3.94 (s, 3H), 1.73 (m, 2H), 3.65 (m, 2H).
7	N-(4-{[6,7-bis(methyloxy)-2-(methylthio)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) d 10.34 (s, 1H), 9.94 (s, 1H), 7.83 (d, 1H), 7.59 (m, 2H), 7.56 (m, 1H), 7.40 (m, 2H), 7.23 (s, 1H), 7.09 (t, 2H), 6.12 (s, 1H), 3.88 (s, 3H), 3.85 (s, 3H), 2.48 (s, 3H), 1.40 (m, 4H).
8	N-(4-fluorophenyl)-N'-(4-{[2-methyl-6,7-bis(methyloxy)quinazolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) 10.15 (bs, 1H), 10.01 (bs, 1H), 7.69-7.75 (m, 2H), 7.61-7.68 (m, 2H), 7.52 (s, 1H), 7.32 (s, 1H), 7.23-7.29 (m, 2H), 7.12-7.19 (m, 2H), 3.93 (d, 6H), 2.43 (s, 3H), 1.53 (s, 4H).
9	N-(4-{[2-amino-6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) d 10.34 (s, 1H), 9.95 (s, 1H), 7.82 (d, 1H), 7.58 (m, 2H), 7.44 (d, 1H), 7.33 (t, 1H), 7.25 (s, 1H), 7.09 (t, 2H), 7.07 (s, 1H), 6.17 (br s, 2H), 5.66 (s, 1H), 3.79 (s, 3H), 3.77 (s, 3H), 1.40 (d, 4H).
10	N-(3-fluoro-4-{[2-(methylamino)-6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) d 10.42 (s, 1H), 9.91 (s, 1H), 7.88 (dd, 1H), 7.56 (m, 2H), 7.44 (m, 4H), 7.09 (t, 2H), 5.90 (s, 1H), 3.88 (s, 3H), 3.85 (s, 3H), 3.39 (br s, 1H), 2.92 (s, 3H), 1.41 (dt, 4H).
11	(1S,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆) d 10.49 (br s, 1 H), 10.26 (br s, 1 H), 10.15 (br s, 1 H), 8.74 (br s, 1 H), 7.95 (br d, J = 13.2 Hz, 1 H), 7.8-7.5 (m, 6 H), 7.16 (t, J = 8.9 Hz, 2 H), 6.82 (br s, 1 H), 4.34 (t, J = 5.9 Hz, 2 H), 4.02 (s, 3 H), 3.99 (br s, 2 H), 3.77 (br t, J = 12.0 Hz, 2 H), 3.56-3.30 (m, 4 H), 3.17-3.07 (m, 2 H), 2.40-2.30 (m, 2 H), 2.04-1.95 (m, 1 H), 1.45 (dd, J = 7.2, 4.7 Hz, 1 H), 1.36 (dd, J = 8.5, 4.5 Hz, 1 H), 1.09 (d, J = 6.2 Hz, 3 H).

Table 3

Entry	Name	¹ H-NMR
12	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆) d 10.65 (br s, 1 H), 10.54 (br s, 1 H), 9.74 (s, 1 H), 8.75 (br s, 1 H), 8.01 (br d, J = 12.9 Hz, 1 H), 7.80-7.50 (m, 6 H), 7.20-7.10 (m, 2 H), 6.84 (br s, 1 H), 4.34 (br t, J = 5 Hz, 2 H), 4.04 (s, 3 H), 4.05-3.95 (m, 2 H), 3.77 (br t, J = 11 Hz, 2 H), 3.52 (br d, J = 12.7 Hz, 4 H), 3.12 (br q, J = 9.0 Hz, 2 H), 2.40-2.30 (m, 2 H), 2.10-1.95 (m, 1 H), 1.40-1.30 (m, 2 H), 1.10 (d, J = 6.2 Hz, 3 H).
13	N-(4-{{6-{{3-(diethylamino)propyl}oxy}-7-(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) d 10.37 (br s, 1H), 10.00 (s, 1H), 8.44 (d, 1H), 7.87 (d, 1H), 7.62 (m, 2H), 7.49 (m, 2H), 7.41 (m, 2H), 7.13 (t, 2H), 6.40 (d, 1H), 4.17 (t, 2H), 3.93 (s, 3H), 2.59 (t, 2H), 2.49 (m, 6H), 1.91 (m, 4H), 0.94 (t, 6H).
14	N-(4-{{6-{{2-(diethylamino)ethyl}oxy}-7-(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) d 10.36 (br s, 1H), 9.99 (s, 1H), 8.44 (d, 1H), 7.88 (dd, 1H), 7.62 (m, 2H), 7.57 (m, 2H), 7.41 (m, 2H), 7.13 (t, 2H), 6.40 (d, 1H), 4.17 (t, 2H), 3.93 (s, 3H), 2.85 (t, 2H), 2.56 (q, 4H), 2.49 (m, 4H), 0.98 (t, 6H).
15	1,1-dimethylethyl 4-(3-{{4-{{2-fluoro-4-{{1-{{4-fluorophenyl}amino}carbonyl}cyclopropyl}carbonyl}amino}phenyl}oxy}-6-(methyloxy)quinolin-7-yl}oxy}propyl)piperazine-1-carboxylate	¹ H NMR (400 MHz, CDCl ₃): 10.05 (s, 1H), 8.49-8.27 (t, 1H), 7.79-7.76 (d, 1H), 7.57 (s, 1H), 7.47-7.43 (m, 3H), 7.27-7.20 (m, 1H), 7.09-7.04 (m, 2H), 6.40-6.39 (d, 1H), 4.28-4.25 (t, 2H), 3.50 (s, 3H), 3.47-3.44, (t, 4H), 2.62-2.59 (t, 2H), 2.46-2.44 (t, 4H), 2.18-2.11 (m, 2H), 2.09 (s, 1H), 1.83-1.81 (t, 2H), 1.64-1.61 (t, 2H), 1.47 (s, 9H).
16	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) d 10.40 (s, 1H), 9.65 (s, 1H), 8.45 (s, 1H), 7.79 (dd, 1H), 7.53 (m, 2H), 7.47 (s, 1H), 7.36 (m, 1H), 7.31 (m, 2H), 7.05 (t, 2H), 4.17 (t, 2H), 3.91 (s, 3H), 3.51 (t, 4H), 2.40 (t, 2H), 2.36 (m, 4H), 1.90 (m, 3H), 1.30 (m, 2H), 1.02 (d, 3H).
17	(1R,2R)-N-(4-{{7-{{2-(diethylamino)ethyl}oxy}-6-(methyloxy)quinazolin-4-yl}oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) d 10.49 (s, 1H), 9.73 (s, 1H), 8.52 (s, 1H), 7.85 (dd, 1H), 7.61 (m, 2H), 7.54 (s, 1H), 7.41 (m, 3H), 7.12 (t, 2H), 7.23 (t, 2H), 3.96 (s, 3H), 2.86 (t, 2H), 2.56 (q, 4H), 1.98 (m, 1H), 1.34 (m, 2H), 1.07 (d, 3H), 0.97 (t, 6H).

Table 3

Entry	Name	¹ H-NMR
18	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆) 8.51 (s, 1H), 7.78-7.84 (m, 1H), 7.58-7.64 (m, 2H), 7.53 (s, 1H), 7.34-7.48 (m, 3H), 7.13 (t, 2H), 4.22 (t, 2H), 3.98 (s, 3H), 2.84 (t, 2H), 2.55 (q, 4H), 1.48 (s, 4H), 1.39 (t, 6H).
19	N-(4-{[7-{[3-(4-acetylpiperazin-1-yl)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, CDCl ₃): 10.04 (s, 1H), 8.48-8.47 (d, 1H), 8.21 (s, 1H), 7.79-7.76 (d, 1H), 7.57 (s, 1H), 7.52-7.44 (m, 3H), 7.28-7.20 (m, 2H), 7.09-7.05 (t, 2H), 6.40-6.39 (d, 1H), 4.30-4.26 (t, 2H), 4.04 (s, 3H), 3.64-3.62 (t, 2H), 3.49-3.47 (t, 2H), 2.62-2.58 (t, 2H), 2.50-2.44 (m, 4H), 2.17-2.12 (m, 2H), 2.10 (s, 3H), 1.84-1.81 (t, 2H), 1.64-1.61 (t, 2H).
20	1,1-dimethylethyl 4-(3-{[4-{(2-fluoro-4-{[(1R,2R)-1-{[(4-fluorophenyl)amino]carbonyl}-2-methylcyclopropyl)carbonyl]amino}phenyl]oxy}-6-(methyloxy)quinolin-7-yl]oxy}propyl)piperazine-1-carboxylate	¹ H NMR (400 MHz, DMSO-d ₆): 10.54 (s, 1H), 9.72 (s, 1H), 8.47-8.46 (d, 1H), 7.96-7.93 (dd, 1H), 7.63-7.61 (m, 2H), 7.52 (br s, 2H), 7.42-7.40 (d, 2H), 7.17-7.12 (t, 2H), 6.44-6.42 (d, 1H), 4.22-4.18 (t, 2H), 3.95 (s, 3H), 3.42-3.40 (m, 2H), 2.36-2.26 (m, 8H), 2.00-1.98 (m, 3H), 1.58-1.54 (m, 2H), 1.40 (s, 9H), 1.10-1.09 (d, 3H).
21	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)-1-(phenylmethyl)azetidine-3,3-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.0 (s, 1H), 9.9 (s, 1H), 8.45 (d, 1H), 7.8 (d, 2H), 7.7 (m, 2H), 7.5 (s, 1H), 7.4 (s, 1H), 7.48-7.15 (m, 9H), 3.95 (s, 6H), 3.7 (s, 4H), 3.6 (s, 2H).
22	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)azetidine-3,3-dicarboxamide	¹ H NMR (DMSO-d ₆): 8.46 (d, 1H), 7.84 (d, 2H), 7.70 (m, 2H), 7.50 (s, 1H), 7.40 (s, 1H), 7.24 (d, 2H), 7.20 (t, 2H), 6.44 (d, 1H), 4.03 (s, 4H), 3.95 (s, 6H), 1.90 (s, 3H, acetate salt).
23	(1R,2S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆): 10.26 (s, 1H), 9.75 (s, 1H), 8.47-8.46 (d, 1H), 7.91-7.87 (dd, 1H), 7.70-7.66 (m, 2H), 7.56-7.51 (m, 2H), 7.43-7.38 (m, 2H), 6.42-6.41 (d, 1H), 4.20-4.16 (t, 2H), 3.95 (s, 3H), 2.47-2.43 (m, 2H), 2.40-2.24 (m, 5H), 2.14 (s, 3H), 2.03-1.93 (m, 3H), 1.89 (s, 3H), 1.45-1.42 (m, 1H), 1.38-1.35 (m, 1H), 1.10-1.08 (d, 3H).

Table 3

Entry	Name	¹ H-NMR
24	(1R,2R)-N-{3-fluoro-4-[(6-(methyloxy)-7-[[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl]oxy]phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): 10.56 (s, 1H), 9.75 (s, 1H), 8.47-8.46 (d, 1H), 7.96-7.93 (d, 1H), 7.68-7.61 (m, 2H), 7.53-7.52 (m, 2H), 7.44-7.39 (m, 2H), 7.18-7.12 (m, 2H), 6.44-6.42 (d, 1H), 4.20-4.17 (t, 2H), 3.95 (s, 3H), 3.42-3.30 (m, 3H), 2.46-2.44 (m, 2H), 2.33 (br s, 2H), 2.15 (s, 3H), 2.05-1.94 (m, 2H), 1.89 (s, 5H), 1.40-1.35 (m, 1H), 1.10-1.09 (m, 3H).
25	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-yl)propyl]oxy}quinolin-4-yl]oxy]phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): 10.55 (s, 1H), 9.72 (s, 1H), 8.47-8.46 (d, 1H), 7.96-7.93 (d, 1H), 7.68-7.61 (m, 2H), 7.52 (br s, 2H), 7.44-7.40 (m, 2H), 7.17-7.12 (m, 2H), 6.44-6.43 (d, 1H), 4.21-4.18 (t, 2H), 3.95 (s, 3H), 2.79 (br s, 4H), 2.47-2.44 (t, 2H), 2.38 (br s, 3H), 2.04-1.95 (m, 3H), 1.40-1.35 (m, 2H), 1.11-1.09 (m, 5H).
26	N-(3-fluoro-4-{[7-({3-[4-(1-methylethyl)piperazin-1-yl]propyl}oxy)-6-(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): 10.40 (s, 1H), 10.02 (s, 1H), 8.47-8.46 (d, 1H), 7.92-7.89 (d, 1H), 7.66-7.63 (m, 2H), 7.52-7.51 (d, 2H), 7.44-7.39 (m, 2H), 7.19-7.14 (m, 2H), 6.42-6.41 (d, 1H), 4.20-4.17 (t, 2H), 3.95 (s, 3H), 2.70-2.68 (m, 1H), 2.62-2.55 (m, 2H), 2.46-2.33 (m, 8H), 1.99-1.94 (m, 2H), 1.47 (s, 4H), 1.00-0.95 (m, 6H).
27	N-(4-{[7-{{3-(diethylamino)propyl}oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆) 10.34 (s, 1H), 10.01 (s, 1H), 8.50 (s, 1H), 7.81 (dd, 1H), 7.55-7.68 (m, 2H), 7.51-7.55 (m, 2H), 7.33-7.48 (m, 3H), 7.12 (t, 2H), 4.22 (t, 2H), 3.94 (s, 3H), 2.52-2.61 (m, 2H), 2.49-2.51 (m, 4H), 1.83-1.94 (m, 2H), 1.42 (s, 4H), 0.95 (t, 6H).
28	(1R,2R)-N-(4-{[7-{{3-(diethylamino)propyl}oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆) d 10.52 (s, 1H), 9.70 (s, 1H), 8.44 (d, 1H), 7.92 (dd, 1H), 7.61 (m, 2H), 7.50 (m, 2H), 7.43 (m, 2H), 7.12 (t, 2H), 6.41 (d, 1H), 4.17 (t, 2H), 3.93 (s, 3H), 2.55 (m, 2H), 2.31 (m, 4H), 1.98 (m, 1H), 1.88 (m, 2H), 1.35 (m, 2H), 1.07 (d, 3H), 0.94 (t, 6H).
29	(1R,2R)-N-(4-{[7-{{2-(diethylamino)ethyl}oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆) d 10.52 (s, 1H), 9.70 (s, 1H), 8.44 (d, 1H), 7.78 (dd, 1H), 7.61 (m, 2H), 7.51 (m, 2H), 7.41 (m, 2H), 7.12 (t, 2H), 6.41 (d, 1H), 4.17 (t, 2H), 3.93 (s, 3H), 2.85 (t, 2H), 2.57 (q, 4H), 1.98 (m, 1H), 1.34 (m, 2H), 1.07 (d, 3H), 0.98 (t, 6H).

Table 3

Entry	Name	¹ H-NMR
30	(1R,2S)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆) d 10.17 (s, 1H), 9.97 (s, 1H), 8.39 (d, 1H), 7.80 (dd, 1H), 7.62 (m, 2H), 7.45 (m, 2H), 7.31 (m, 2H), 7.09 (t, 2H), 6.34 (d, 1H), 4.12 (t, 2H), 3.88 (s, 3H), 2.46 (m, 2H), 2.40 (m, 4H), 1.92 (m, 1H), 1.84 (m, 2H), 1.37 (m, 1H), 1.29 (m, 1H), 1.01 (d, 3H), 0.89 (t, 6H).
31	(1R,2S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆) d 10.22 (s, 1H), 10.01 (s, 1H), 8.44 (d, 1H), 7.86 (dd, 1H), 7.66 (m, 2H), 7.49 (m, 2H), 7.39 (m, 2H), 7.14 (m, 2H), 6.39 (d, 1H), 4.18 (m, 2H), 3.92 (s, 3H), 2.85 (t, 2H), 2.57 (q, 4H), 1.97 (m, 1H), 1.42 (m, 1H), 1.35 (m, 1H), 1.06 (d, 3H), 0.98 (t, 6H).
32	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	¹ H NMR (CDCl ₃) 8.57 (s, 1H), 8.12 (s, 1H), 7.73-7.81 (m, 2H), 7.48-7.53 (m, 2H), 7.32 (s, 1H), 6.98-7.08 (m, 3H), 4.28 (t, 2H), 4.04 (s, 3H), 3.25 (t, 2H), 2.76 (q, 4H), 2.67 (q, 4H), 2.01-2.15 (m, 2H), 1.10 (t, 6H).
33	(1R,2S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): 10.28 (s, 1H), 9.80 (s, 1H), 8.47-8.46 (d, 1H), 7.90-7.88 (d, 1H), 7.70-7.62 (m, 2H), 7.56-7.52 (m, 2H), 7.44-7.39 (m, 2H), 7.18-7.12 (m, 2H), 6.44-6.41 (t, 1H), 4.20-4.17 (t, 2H), 3.95 (s, 3H), 2.74-2.72 (t, 3H), 2.46-2.42 (m, 1H), 2.35 (br s, 3H), 2.03-1.93 (m, 3H), 1.87 (s, 4H), 1.43-1.35 (m, 2H), 1.09-1.08 (m, 3H).
34	(1r,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆): 1H (10.12 ppm, s), 1H (9.6 ppm, s), 1H (8.46 ppm, d), 1H (7.88 ppm, dd), 2H (7.68 ppm, m), 1H (7.56 ppm, d), 1H (7.51 ppm, s), 2H (7.4 ppm, m), 2H (7.13 ppm, t), 1H (6.4 ppm, d), 2H (4.2 ppm, t), 3H (3.94 ppm, s), 4H (3.6 ppm, t), 2H (2.45 ppm, t), 4H (2.37 ppm, m), 2H (1.97 ppm, t), 2H (1.8 ppm, m), 6 H (1.28 ppm, d).
35	(1r,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆): 1H (10.12 ppm, s), 1H (9.6 ppm, s), 1H (8.46 ppm, d), 1H (7.88 ppm, dd), 2H (7.69 ppm, m), 1H (7.58 ppm, d), 1H (7.51 ppm, s), 2H (7.4 ppm, m), 2H (7.13 ppm, t), 1H (6.4 ppm, d), 2H (4.2 ppm, t), 3H (3.95 ppm, s), 10H (2.35 ppm, m), 3H (2.14 ppm, s), 2H (1.97 ppm, t), 2H (1.8 ppm, m), 6H (1.28 ppm, d).

Table 3

Entry	Name	¹ H-NMR
36	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆): 8.45 (d, 2 H), 8.15 (d, 1H), 7.8 (d, 1H), 7.45 (m, 3H), 7.25 (m, 3H), 7.0 (m, 2H), 4.20 (t, 2H), 4.0 (s, 3H), 3.7 (m, 4H), 2.67 (m, 4H), 2.45 (m, 6H), 2.0 (m, 4H).
37	(2R,3R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, CDCl ₃) d 8.44 (d, J = 5.1 Hz, 1 H), 8.11 (br s, 1 H), 7.77-7.70 (m, 2 H), 7.53 (s, 1 H), 7.50-7.44 (m, 2 H), 7.40 (s, 1 H), 7.22-7.16 (m, 2 H), 7.06-6.98 (m, 2 H), 6.36 (br d, J = 5.1 Hz, 1 H), 4.26 (t, J = 7.0 Hz, 2 H), 4.02 (s, 3 H), 3.72 (t, J = 4.4 Hz, 4 H), 2.57 (t, J = 7.3 Hz, 2 H), 2.50-2.42 (m, 4 H), 2.18-2.10 (m, 2 H), 1.80-1.66 (m, 2 H), 1.30-1.24 (m, 6 H).
38	(2R,3R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆) d 10.34 (s, 1 H), 10.05 (s, 1 H), 8.46 (br s, 1 H), 7.93 (br d, J = 4.7 Hz, 1 H), 7.54-7.52 (m, 2 H), 7.52-7.50 (m, 2 H), 7.50-7.30 (m, 2 H), 7.20-7.10 (m, 2 H), 6.47 (br s, 1 H), 4.30-4.20 (m, 2 H), 3.95 (s, 3 H), 3.40-3.10 (m, 6 H), 2.60-2.40 (m, 2 H), 1.90-1.80 (m, 2 H), 1.30-1.10 (m, 12 H).
39	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400MHz, DMSO-d ₆): d 10.47 (s, 1H), 10.16 (s, 1H), 8.43 (d, 1H), 7.92 (dd, 1H), 7.67 (m, 2H), 7.58 (m, 1H), 7.52 (s, 1H), 7.41 (m, 2H), 7.15 (t, 2H), 6.44 (d, 1H), 4.25 (t, 2H), 3.95 (s, 3H), 3.10 (m, 6H), 2.17 (m, 2H), 1.91 (s, 3H, acetate salt), 1.52 (m, 2H), 1.18 (m, 12H).
40	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400MHz, DMSO-d ₆): d 10.21 (s, 1H), 9.97 (s, 1H), 8.51 (s, 1H), 7.81 (dd, 1H), 7.64 (m, 2H), 7.54 (s, 1H), 7.48 (m, 1H), 7.41 (m, 1H), 7.38 (s, 1H), 7.15 (t, 2H), 4.24 (t, 2H), 3.97 (s, 3H), 3.58 (m, 4H), 2.45 (t, 2H), 2.38 (m, 4H), 1.97 (m, 2H), 1.58 (m, 2H), 1.18 (s, 3H), 1.17 (s, 3H).
41	(1R,2R,3S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400MHz, DMSO-d ₆): d 10.14 (s, 1H), 9.61 (s, 1H), 8.46 (d, 1H), 7.87 (dd, 1H), 7.67 (m, 2H), 7.57 (m, 1H), 7.51 (s, 1H), 7.42 (s, 1H), 7.39 (m, 1H), 7.15 (t, 2H), 6.41 (d, 1H), 4.20 (m, 2H), 3.94 (s, 3H), 2.87 (m, 2H), 2.60 (m, 4H), 1.80 (m, 2H), 1.18 (s, 3H), 1.17 (s, 3H), 1.01 (m, 6H). Note: 0.5eq of AcOH is present by NMR.

Table 3

Entry	Name	¹ H-NMR
42	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400MHz, DMSO): d 10.24 (s, 1H), 10.00 (s, 1H), 8.54 (s, 1H), 7.84 (dd, 1H), 7.66 (m, 2H), 7.56 (s, 1H), 7.51 (m, 1H), 7.43 (m, 2H), 7.18 (t, 2H), 4.26 (m, 2H), 3.98 (s, 3H), 2.88 (m, 2H), 2.59 (m, 4H), 1.58 (m, 2H), 1.18 (s, 6H), 1.00 (t, 6H).
43	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400MHz, DMSO-d ₆): d 10.21 (s, 1H), 9.97 (s, 1H), 8.51 (s, 1H), 7.82 (dd, 1H), 7.64 (m, 2H), 7.54 (s, 1H), 7.48 (m, 1H), 7.41 (m, 1H), 7.37 (s, 1H), 7.15 (t, 2H), 4.23 (t, 2H), 3.97 (s, 3H), 2.56 (m, 2H), 2.46 (m, 4H), 1.91 (m, 2H), 1.58 (m, 2H), 1.18 (s, 6H), 0.96 (t, 6H).
44	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	¹ H NMR (CDCl ₃): 8.57 (s, 1H), 8.50 (s, 1H), 8.11 (s, 1H), 7.81 (dd, 1H), 7.53 (m, 3H), 7.28 (m, 4H), 7.04 (t, 2H), 4.24 (t, 2H), 4.04 (s, 3H), 2.95 (t, 2H), 2.84 (q, 4H), 2.75 (m, 4H), 2.21 (m, 2H), 2.02 (m, 2H), 1.18 (t, 6H).
45	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	¹ H NMR (CDCl ₃): 8.57 (s, 1H), 8.49 (s, 1H), 8.10 (s, 1H), 7.80 (d, 1H), 7.70 (br., 1H), 7.52 (m, 3H), 7.31 (m, 3H), 7.04 (t, 2H), 4.26 (t, 2H), 4.04 (s, 3H), 2.62-2.77 (m, 14H), 2.40 (s, 3H), 2.13 (m, 2H), 2.01 (m, 2H).
46	(2R,3R)-N-[3-fluoro-4-[(6-(methyloxy)-7-[(3-morpholin-4-yl)propyl]oxy]quinazolin-4-yl)oxy]phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, CDCl ₃) d 8.59 (s, 1 H), 8.11 (br s, 1 H), 7.80-7.76 (m, 2 H), 7.53 (s, 1 H), 7.50-7.46 (m, 2 H), 7.34 (s, 1 H), 7.26-7.24 (m, 2 H), 7.06-7.00 (m, 2 H), 4.28 (t, J = 6.6 Hz, 2 H), 4.05 (s, 3 H), 3.73 (br t, J = 4.4 Hz, 4 H), 2.57 (t, J = 7.0 Hz, 2 H), 2.52-2.45 (m, 4 H), 2.18-2.10 (m, 2 H), 1.80-1.68 (m, 2 H), 1.28-1.20 (m, 6 H).
47	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO-d ₆): 9.98 (s, 1H), 9.72 (s, 1H), 8.45-8.43 (d, 1H), 7.97-7.94 (dd, 1H), 7.73-7.69 (m, 2H), 7.65-7.52 (m, 3H), 7.44-7.39 (m, 1H), 7.18-7.14 (m, 2H), 6.43-6.42 (d, 1H), 4.20-4.19 (t, 2H), 3.95 (s, 3H), 2.70-2.66 (m, 6H), 2.45 (br s, 2H), 1.91-1.84 (m, 6H), 0.98 (br s, 6H).

Table 3

Entry	Name	¹ H-NMR
48	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆): 9.99 (s, 1H), 9.73 (s, 1H), 8.45-8.43 (d, 1H), 7.97-7.93 (dd, 1H), 7.73-7.69 (m, 2H), 7.65-7.52 (m, 2H), 7.44-7.38 (m, 2H), 7.18-7.14 (m, 2H), 6.43-6.42 (d, 1H), 4.19-4.16 (t, 3H), 3.95 (s, 3H), 2.70-2.66 (m, 4H), 2.47-2.33 (m, 8H), 2.15 (s, 3H), 1.98-1.94 (m, 2H), 1.90-1.84 (m, 4H).
49	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, CDCl ₃) d 8.59 (br s, 1 H), 8.29 (br s, 1 H), 7.93 (s, 1 H), 7.77 (d, J = 10.8 Hz, 1 H), 7.53 (s, 1 H), 7.50-7.45 (m, 2 H), 7.32 (s, 1 H), 7.26-7.22 (m, 2 H), 7.05-6.99 (m, 2 H), 4.27 (t, J = 6.6 Hz, 2 H), 4.04 (s, 3 H), 3.03 (t, J = 6.5 Hz, 2 H), 2.67 (q, J = 7.0 Hz, 4 H), 1.80-1.70 (m, 2 H), 1.22 (br t, J = 5.3 Hz, 6 H), 1.09 (br t, J = 7.2 Hz, 6 H).
50	(2R,3R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, CDCl ₃) d 8.58 (s, 1 H), 8.40-8.36 (m, 1 H), 8.02-7.96 (m, 1 H), 7.80-7.75 (m, 1 H), 7.53 (s, 1 H), 7.52-7.50 (m, 2 H), 7.31 (s, 1 H), 7.28-7.20 (m, 2 H), 7.02 (t, J = 8.5 Hz, 2 H), 4.25 (t, J = 6.3 Hz, 2 H), 4.04 (s, 3 H), 3.00-2.90 (m, 2 H), 2.88-2.80 (m, 4 H), 2.30-2.20 (m, 2 H), 1.76-1.68 (m, 2 H), 1.25-1.15 (m, 12 H).
51	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	¹ H NMR (400 MHz, CDCl ₃) d 8.47 (d, J = 5.2 Hz, 1 H), 8.17 (br s, 1 H), 7.80-7.74 (m, 2 H), 7.55 (s, 1 H), 7.52-7.46 (m, 2 H), 7.42 (s, 1 H), 7.24-7.20 (m, 2 H), 7.05 (t, J = 8.6 Hz, 2 H), 6.38 (br d, J = 5.4 Hz, 1 H), 4.27 (t, J = 6.4 Hz, 2 H), 4.03 (s, 3 H), 3.04 (br t, J = 7.2 Hz, 2 H), 2.68 (q, J = 6.8 Hz, 4 H), 1.80-1.68 (m, 2 H), 1.26 (d, J = 6.4 Hz, 6 H), 1.09 (br t, J = 7.2 Hz, 6 H).
52	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[(4-fluorophenyl)methyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆): 10.82 (s, 1H), 8.80 (d, 1H), 8.50 (t, 1H), 7.83 (d, 2H), 7.74 (s, 1H), 7.56 (s, 1H), 7.30-7.38 (m, 4H), 7.15 (t, 2H), 6.80 (d, 1H), 4.32 (d, 2H), 4.04 (s, 3H), 4.03 (s, 3H), 1.42 (s, 4H).
53	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(2-morpholin-4-ylethyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO- <i>d</i> ₆): 10.62 (s, 1H), 8.79 (d, 1H), 8.24 (t, 1H), 7.83 (d, 2H), 7.72 (s, 1H), 7.58 (s, 1H), 7.37 (d, 2H), 6.76 (d, 1H), 4.04 (s, 3H), 4.03 (s, 3H), 3.98 (m, 2H), 3.66 (m, 2H), 3.49 (m, 4H), 3.25 (t, 2H), 3.13 (br., 2H), 1.42 (d, 4H).

Table 3

Entry	Name	¹ H-NMR
54	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.78 (s, 1H), 10.53 (s, 1H), 8.43 (d, 1H), 8.12 (d, 1H), 7.82 (d, 2H), 7.49 (s, 1H), 7.37 (s, 1H), 7.20-7.28 (m, 3H), 7.15 (dd, 1H), 7.01 (td, 1H), 6.35 (d, 1H), 3.93 (s, 3H), 3.92 (s, 3H), 3.47 (s, 2H), 2.17 (br., 4H), 1.49 (m, 4H), 1.41 (m, 4H), 1.32 (br., 2H).
55	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.98 (s, 1H), 10.56 (s, 1H), 8.42 (d, 1H), 8.10 (dd, 1H), 7.81 (m, 2H), 7.49 (s, 1H), 7.37 (s, 1H), 7.17-7.27 (m, 4H), 7.01 (td, 1H), 6.35 (d, 1H), 3.93 (s, 3H), 3.92 (s, 3H), 3.61 (s, 2H), 2.30 (br., 4H), 1.47 (br., 4H), 1.43 (m, 4H).
56	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.12 (s, 1H), 10.03 (s, 1H), 8.44 (d, 1H), 7.74 (d, 2H), 7.57 (s, 1H), 7.53 (d, 1H), 7.48 (s, 1H), 7.37 (s, 1H), 7.21 (m, 3H), 6.98 (d, 1H), 6.40 (d, 1H), 3.93 (s, 3H), 3.92 (s, 3H), 3.56 (t, 4H), 3.41 (s, 2H), 2.34 (br., 4H), 1.48 (s, 4H).
57	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.54 (s, 1H), 10.47 (s, 1H), 8.43 (d, 1H), 8.08 (d, 1H), 7.78 (d, 2H), 7.49 (s, 1H), 7.37 (d, 1H), 7.18-7.30 (m, 4H), 7.03 (t, 1H), 6.37 (d, 1H), 3.94 (s, 3H), 3.93 (s, 3H), 3.50 (s, 2H), 3.44 (br., 4H), 2.20 (br., 4H), 1.48 (d, 4H).
58	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-phenylcyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.14 (s, 1H), 10.03 (s, 1H), 8.44 (d, 1H), 7.74 (d, 2H), 7.62 (d, 2H), 7.48 (s, 1H), 7.37 (s, 1H), 7.27-7.31 (m, 2H), 7.19-7.23 (m, 2H), 7.05 (t, 1H), 6.41 (d, 1H), 3.93 (s, 6H), 3.92 (s, 3H), 1.48 (s, 4H).
59	N-[3-(aminomethyl)phenyl]-N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide	¹ H NMR (400MHz, DMSO-d ₆): d 10.28 (s, 1H), 10.19 (s, 1H), 8.77 (m, 1H), 8.21 (m, 3H), 7.84 (m, 2H), 7.76 (m, 1H), 7.71 (m, 1H), 7.58 (m, 2H), 7.38 (m, 3H), 7.19 (m, 1H), 6.76 (m, 1H), 4.03 (s, 6H), 3.39 (m, 2H), 1.53 (m, 4H). Note: all peaks are very broad and unresolved.
60	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.0-10.2 (br., 2H), 8.46 (d, 1H), 7.76 (d, 2H), 7.53 (m, 3H), 7.39 (s, 1H), 7.24 (m, 3H), 6.98 (d, 1H), 6.43 (d, 1H), 3.95 (s, 3H), 3.93 (s, 3H), 3.37 (s, 2H), 2.31 (br., 4H), 1.48 (m, 8H), 1.39 (br., 2H).

Table 3

Entry	Name	¹ H-NMR
61	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	¹ H NMR (DMSO-d ₆): 10.0-10.2 (br., 2H), 8.46 (d, 1H), 7.77 (d, 2H), 7.59 (s, 1H), 7.53 (d, 1H), 7.51 (s, 1H), 7.39 (s, 1H), 7.23 (m, 3H), 6.99 (d, 1H), 6.43 (d, 1H), 3.95 (s, 3H), 3.93 (s, 3H), 3.52 (s, 2H), 2.42 (br., 4H), 1.69 (br, 4H), 1.48 (s, 4H).

Assays

[0455] Kinase assays were performed by measurement of incorporation of γ -³³P ATP into immobilized myelin basic protein (MBP). High binding white 384 well plates (Greiner) were coated with MBP (Sigma #M-1891) by incubation of 60ul/well of 20μg/ml MBP in Tris-buffered saline (TBS; 50mM Tris pH 8.0, 138mM NaCl, 2.7mM KCl) for 24 hours at 4° C. Plates were washed 3X with 100μl TBS. Kinase reactions were carried out in a total volume of 34μl in kinase buffer (5mM Hepes pH 7.6, 15mM NaCl, 0.01% bovine gamma globulin (Sigma #I-5506), 10mM MgCl₂, 1mM DTT, 0.02% TritonX-100). Compound dilutions were performed in DMSO and added to assay wells to a final DMSO concentration of 1%. Each data point was measured in duplicate, and at least two duplicate assays were performed for each individual compound determination. Enzyme was added to final concentrations of 10nM or 20nM, for example. A mixture of unlabeled ATP and γ -³³P ATP was added to start the reaction (2x10⁶ cpm of γ -³³P ATP per well (3000Ci/mmol) and either 10μM or 30μM unlabeled ATP, typically. The reactions were carried out for 1 hour at room temperature with shaking. Plates were washed 7x with TBS, followed by the addition of 50μl/well scintillation fluid (Wallac). Plates were read using a Wallac Trilux counter. This is only one format of such assays, various other formats are possible, as known to one skilled in the art.

[0456] The above assay procedure can be used to determine the IC₅₀ for inhibition and/or the inhibition constant, K_i. The IC₅₀ is defined as the concentration of compound required to reduce the enzyme activity by 50% under the conditions of the assay. Exemplary compositions have IC₅₀'s of, for example, less than about 100 μM, less than about 10 μM, less than about 1 μM, and further for example having IC₅₀'s of less than about 100 nM, and still further, for example, less than about 10 nM. The K_i for a compound may be

determined from the IC₅₀ based on three assumptions. First, only one compound molecule binds to the enzyme and there is no cooperativity. Second, the concentrations of active enzyme and the compound tested are known (i.e., there are no significant amounts of impurities or inactive forms in the preparations). Third, the enzymatic rate of the enzyme-inhibitor complex is zero. The rate (i.e., compound concentration) data are fitted to the equation:

$$V = V_{\max} E_0 \left[I - \frac{(E_0 + I_0 + K_d) - \sqrt{(E_0 + I_0 + K_d)^2 - 4E_0 I_0}}{2E_0} \right]$$

where V is the observed rate, V_{max} is the rate of the free enzyme, I₀ is the inhibitor concentration, E₀ is the enzyme concentration, and K_d is the dissociation constant of the enzyme-inhibitor complex.

Kinase specificity assays:

[0457] Kinase activity and compound inhibition are investigated using one or more of the three assay formats described below. The ATP concentrations for each assay are selected to be close to the Michaelis-Menten constant (K_M) for each individual kinase. Dose-response experiments are performed at 10 different inhibitor concentrations in a 384-well plate format. The data are fitted to the following four-parameter equation:

$$Y = \text{Min} + (\text{Max} - \text{Min}) / (1 + (X/\text{IC}_{50})^H)$$

where Y is the observed signal, X is the inhibitor concentration, Min is the background signal in the absence of enzyme (0% enzyme activity), Max is the signal in the absence of inhibitor (100% enzyme activity), IC₅₀ is the inhibitor concentration at 50% enzyme inhibition and H represents the empirical Hill's slope to measure the cooperativity. Typically H is close to unity.

c-Met Assay

[0458] c-Met biochemical activity was assessed using a Luciferase-Coupled Chemiluminescent Kinase assay (LCCA) format as described above. Again, kinase activity was measured as the percent ATP remaining following the kinase reaction.

Remaining ATP was detected by luciferase-luciferin-coupled chemiluminescence. Specifically, the reaction was initiated by mixing test compounds, 1 μ M ATP, 1 μ M poly-EY and 10nM c-Met (baculovirus expressed human c-Met kinase domain P948-S1343) in a 20 μ L assay buffer (20mM Tris-HCL pH7.5, 10mM MgCl₂, 0.02% Triton X-100, 100mM DTT, 2mM MnCl₂). The mixture is incubated at ambient temperature for 2hours after which 20 μ L luciferase-luciferin mix is added and the chemiluminescent signal read using a Wallac Victor² reader. The luciferase-luciferin mix consists of 50 mM HEPES, pH 7.8, 8.5 μ g/mL oxalic acid (pH 7.8), 5 (or 50) mM DTT, 0.4% Triton X-100, 0.25 mg/mL coenzyme A, 63 μ M AMP, 28 μ g/mL luciferin and 40,000 units of light/mL luciferase.

KDR Assay

[0459] KDR biochemical activity was assessed using a Luciferase-Coupled Chemiluminescent Kinase assay (LCCA) format. Kinase activity was measured as the percent ATP remaining following the kinase reaction. Remaining ATP was detected by luciferase-luciferin-coupled chemiluminescence. Specifically, the reaction was initiated by mixing test compounds, 3 μ M ATP, 1.6 μ M poly-EY and 5 nM KDR (baculovirus expressed human KDR kinase domain D807-V1356) in a 20 μ L assay buffer (20mM Tris-HCL pH7.5, 10mM MgCl₂, 0.01% Triton X-100, 1mM DTT, 3mM MnCl₂). The mixture is incubated at ambient temperature for 4 hours after which 20 μ L luciferase-luciferin mix is added and the chemiluminescent signal read using a Wallac Victor² reader. The luciferase-luciferin mix consists of 50 mM HEPES, pH 7.8, 8.5 μ g/mL oxalic acid (pH 7.8), 5 (or 50) mM DTT, 0.4% Triton X-100, 0.25 mg/mL coenzyme A, 63 μ M AMP, 28 μ g/mL luciferin and 40,000 units of light/mL luciferase.

flt-4 Assay

[0460] Biochemical activity for flt-4 was assessed using an Alphascreen Tyrosine Kinase protocol. AlphaScreenTM (Perkin Elmer) technology is a proximity assay employing microparticles. Singlet oxygen derived from a donor bead following laser excitation results in chemiluminescence when in proximity (100 Å) to an acceptor bead due to biomolecular interactions. For the Flt-4 assay, donor beads coated with streptavidin and acceptor beads coated with PY100 anti-phosphotyrosine antibody were used (Perkin Elmer). Biotinylated poly(Glu,Tyr) 4:1 (Perkin Elmer) was used as the substrate.

Substrate phosphorylation was measured by addition of donor/acceptor beads by chemiluminescence following donor-acceptor bead complex formation. Test compounds, 5 μ M ATP, 3 nM biotinylated poly(Glu, Tyr) and 1 nM Flt-4 (baculovirus expressed human Flt-4 kinase domain D725-R1298) were combined in a volume of 20 μ L in a 384-well white, medium binding microtiter plate (Greiner). Reaction mixtures were incubated for 1 hr at ambient temperature. Reactions were quenched by addition of 10 μ L of 15-30 mg/mL AlphaScreen bead suspension containing 75 mM Hepes, pH 7.4, 300 mM NaCl, 120 mM EDTA, 0.3% BSA and 0.03% Tween-20. After 2-16 hr incubation at ambient temperature plates were read using an AlphaQuest reader (Perkin Elmer). IC₅₀ values correlate well with those determined by radiometric assays.

flt-3 Assay

[0461] Biochemical activity for flt-3 was assessed using a Luciferase-Coupled Chemiluminescent Kinase assay (LCCA) format. Kinase activity was measured as the percent ATP remaining following the kinase reaction. Remaining ATP was detected by luciferase-luciferin-coupled chemiluminescence. Specifically, the reaction was initiated by mixing test compounds, 5 μ M ATP, 3 μ M poly-EY and 5 nM Flt-3 (baculovirus expressed human Flt-3 kinase domain R571-S993) in a 20 μ L assay buffer (20mM Tris-HCL pH7.5, 10mM MgCl₂, 0.01% Triton X-100, 1mM DTT, 2mM MnCl₂). The mixture is incubated at ambient temperature for 3 hours after which 20 μ L luciferase-luciferin mix is added and the chemiluminescent signal read using a Wallac Victor² reader. The luciferase-luciferin mix consists of 50 mM HEPES, pH 7.8, 8.5 μ g/mL oxalic acid (pH 7.8), 5 (or 50) mM DTT, 0.4% Triton X-100, 0.25 mg/mL coenzyme A, 63 μ M AMP, 28 μ g/mL luciferin and 40,000 units of light/mL luciferase.

c-Kit Assay

[0462] c-Kit biochemical activity was assessed using AlphaScreen TM (Perkin Elmer) technology, described above. Test compounds, ATP, biotinylated poly(Glu, Tyr) and c-Kit kinase were combined in a volume of 20 μ L in a 384-well white, medium binding microtiter plate (Greiner). Reaction mixtures were incubated for 1 hr at ambient temperature. Reactions were quenched by addition of 10 μ L of 15-30 mg/mL AlphaScreen bead suspension containing 75 mM Hepes, pH 7.4, 300 mM NaCl, 120 mM

EDTA, 0.3% BSA and 0.03% Tween-20. After 16 hr incubation at ambient temperature plates were read using an AlphaQuest reader (Perkin Elmer).

Structure Activity Relationships

[0463] Table 4 shows structure activity relationship data for selected compounds of the invention. Inhibition is indicated as IC₅₀ with the following key: A = IC₅₀ less than 50 nM, B = IC₅₀ greater than 50 nM, but less than 500 nM, C = IC₅₀ greater than 500 nM, but less than 5000 nM, and D = IC₅₀ greater than 5,000 nM. Depending upon the functionality about the quinazoline or quinoline, exemplary compounds of the invention exhibit selectivity for any of c-Met, KDR, c-Kit, flt-3, and flt-4. Abbreviations for enzymes listed in Tables 2-3 are defined as follows: c-Met refers to hepatocyte growth factor receptor kinase; KDR refers to kinase insert domain receptor tyrosine kinase; flt-4, fms-like tyrosine kinase-4, representative of the FLK family of receptor tyrosine kinases; c-Kit, also called stem cell factor receptor or steel factor receptor; and flt-3, fms-like tyrosine kinase-3. Empty cells in the tables indicate lack of data only.

Table 4

Entry	Name	c-Met	KDR	c-Kit	flt3	flt4
1	N-[(3-fluoro-4-[(6-(methyloxy)-7-[(3aR,6aS)-octahydrocyclopenta[c]pyrrol-5-ylmethyl]oxy)quinazolin-4-yl]oxy]phenyl)amino]carbonothioyl]-2-phenylacetamide		A			A
2	N-[(3-fluoro-4-[(7-[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-yl]oxy]phenyl)amino]carbonothioyl]-2-phenylacetamide	A	A	A		A
3	N-[(4-[(6,7-bis(methyloxy)quinolin-4-yl]oxy)-3-fluorophenyl)(methyl)amino]carbonothioyl]-2-phenylacetamide	C				
4	1-(4-[(6,7-bis(methyloxy)quinolin-4-yl]oxy)-3-fluorophenyl)imidazolidin-2-one	C		C	C	
5	1-(4-[(6,7-bis(methyloxy)quinolin-4-yl]oxy)-3-fluorophenyl)-3-(phenylmethyl)imidazolidin-2-one	C		B	C	
6	1-(4-[(6,7-bis(methyloxy)quinolin-4-yl]oxy)-3-fluorophenyl)-3-(phenylacetyl)imidazolidin-2-one	B				

7	ethyl [(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)amino](oxo)acetate	B	B	C		B
8	N-{[(4-{[6,7-bis(methyloxy)quinazolin-4-yl]amino}-3-fluorophenyl)amino]carbonothioyl}-2-phenylacetamide	A	B	C		B
9	N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N-methyl-N-(2-phenylethyl)sulfamide	C				
10	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-(phenylmethyl)-1,2,4-oxadiazol-5-amine	C		B	C	
11	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)piperidin-2-one	C		C	C	
12	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(phenylmethyl)ethanediamide	B	B	B	C	B
13	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-4-phenyl-1,3-thiazol-2-amine	C		C	B	
14	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-phenylethyl)ethanediamide	A	A	A	C	A
15	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-1-phenylmethanesulfonamide	C		C	B	
16	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-2-phenylethanesulfonamide	C		C	C	
17	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(phenylmethyl)benzenesulfonamide	C		C	C	
18	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(phenylmethyl)benzenesulfonamide	C		C	C	
19	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(2-phenylethyl)benzenesulfonamide	C		C	C	
20	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(2-phenylethyl)benzenesulfonamide	C		C	C	
21	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(3-phenylpropyl)benzenesulfonamide	C				
22	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)pyrrolidin-2-one	C		C	B	
23	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl (phenylmethyl)carbamate	C				
24	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl (2-phenylethyl)carbamate	C				
25	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(3-phenylpropyl)benzenesulfonamide	C				
26	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-phenylethanediamide	B	D	C		C
27	4-{[6,7-bis(methyloxy)quinolin-4-yl]amino}-N-(3-phenylpropyl)benzamide	C				

28	N-[[[3-fluoro-4-{[7-[[2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}phenyl]amino]carbonothioyl]-2-phenylacetamide	A	A	A	A	A
29	N-[(Z)-[(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)amino](imino)methyl]-2-phenylacetamide	C				
30	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-[2-(phenyloxy)ethyl]benzenesulfonamide	C				
31	This type of multiplicative nomenclature is not supported in current version!	C				
32	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-phenylpropane-1-sulfonamide	C				
33	N~2~{[(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)sulfonyl]-N-phenylglycinamide	C				
34	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-2-phenylacetamide	C				
35	N-[[[6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl]amino]carbonothioyl]-2-phenylacetamide	A	C	D		C
36	6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-1,3-benzothiazol-2-amine	C		C	C	
37	6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-amine	C		C	C	
38	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-2-phenylacetamide	B	C	D		B
39	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-morpholin-4-ylethyl)ethanediamide	C		B	B	
40	1,1-dimethylethyl {2-[(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)amino]-2-oxoethyl}(phenylmethyl)carbamate	C				
41	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N~2~(phenylmethyl)glycinamide	B				
42	N~2~acetyl-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N~2~(phenylmethyl)glycinamide	C				
43	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-1,3-benzothiazol-2-yl)-2-phenylacetamide	B				
44	1,1-dimethylethyl {2-[(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)amino]-2-oxoethyl}(phenylmethyl)carbamate	C				
45	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-N~2~(phenylmethyl)glycinamide	C				

46	N~2~-acetyl-N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-N~2~- (phenylmethyl)glycinamide	C				
47	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-3-phenylpropanamide	C				
48	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-4-phenylbutanamide	C				
49	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-N~2~-methyl-N~2~- (phenylmethyl)glycinamide	C				
50	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-[4- (methyloxy)phenyl]ethyl)ethanediamide	C		C	C	
51	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N~2~-methyl-N~2~- (phenylmethyl)glycinamide	B	A	B	B	A
52	N-{[(4-{[6,7-bis(methyloxy)quinolin-4-yl]amino}phenyl)amino]carbonothioyl}-2- phenylacetamide	A	B	B	C	A
53	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-3-phenylpropanamide	C				
54	N-{[(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)amino]carbonothioyl}-2- phenylacetamide	A	B	C		B
55	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2,3-dihydro-1H-inden-1- yl)ethanediamide	A	B	B		B
56	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2,3-dihydro-1H-inden-2- yl)ethanediamide	C				
57	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(1,2,3,4-tetrahydronaphthalen-1- yl)ethanediamide	B	B	C		C
58	N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N-(2-phenylethyl)-N- (phenylmethyl)sulfamide	1470 .06				
59	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N~2~- (trifluoroacetyl)glycinamide	B	C	B		B
60	N-{2-[(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)amino]-2-oxoethyl}benzamide	B	A	A	A	A
61	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-N'-(4-fluorophenyl)propanediamide	A	B	B		B
62	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[(2S)-1,2,3,4-tetrahydronaphthalen-2- yl]ethanediamide	C				

63	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(4-methylphenyl)ethyl]ethanediamide	C		C	C	
64	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-phenylpropyl)ethanediamide	B	A	B	B	B
65	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(4-chlorophenyl)ethyl]ethanediamide	A	C	B	C	C
66	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N,N'-bis(phenylmethyl)sulfamide	C				
67	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N,N'-bis(2-phenylethyl)sulfamide	C				
68	ethyl [(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)amino](oxo)acetate	C				
69	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-phenylethyl)ethanediamide	C				
70	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)propanediamide	A	B	B		C
71	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[(2R)-1,2,3,4-tetrahydronaphthalen-2-yl]ethanediamide	B	D	B		C
72	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(1-methylpyrrolidin-2-yl)ethyl]ethanediamide	C		C	C	
73	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(phenyloxy)ethyl]ethanediamide	B	B	B		C
74	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-hydroxy-1-(phenylmethyl)ethyl]urea	B				
75	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-[(4-methylphenyl)sulfonyl]-4-(phenylmethyl)imidazolidin-2-one	B	B	B	B	B
76	N'-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N-methyl-N-(2-phenylethyl)ethanediamide	A	B	B	B	B
77	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-{[3-(trifluoromethyl)phenyl]methyl}ethanediamide	B		B	B	
78	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-{2-[3-(trifluoromethyl)phenyl]ethyl}ethanediamide	C		A	B	
79	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-3-oxo-4-phenylbutanamide	C				

80	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl})-2-[3-(trifluoromethyl)phenyl]acetamide	C				
81	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-[2-(phenyloxy)ethyl]-1,3-benzothiazol-2-amine	B				
82	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-(2-piperidin-1-ylethyl)-1,3-benzothiazol-2-amine	C				
83	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-methyl-N-(2-phenylethyl)-1,3-benzothiazol-2-amine	C				
84	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-(2-pyrrolidin-1-ylethyl)-1,3-benzothiazol-2-amine	C				
85	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-{{[3-(trifluoromethyl)phenyl]methyl}}-1,3-benzothiazol-2-amine	C				
86	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-{2-[3-(trifluoromethyl)phenyl]ethyl}}-1,3-benzothiazol-2-amine	C				
87	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl})-N'-[3-(trifluoromethyl)phenyl]propanediamide	C				
88	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl})-2-[3-(trifluoromethyl)phenyl]acetamide	C	A	B	B	B
89	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl})-N~2~-{{[3-(trifluoromethyl)phenyl]methyl}}glycinamide	B	A	B	B	A
90	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl})-N~2~-(2-phenylethyl)glycinamide	B				
91	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl})-N~2~-{2-[3-(trifluoromethyl)phenyl]ethyl}glycinamide	B	B	B	B	A
92	1,1-dimethylethyl {2-[[6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl}amino]-2-oxoethyl}(phenylmethyl)carbamate	C				
93	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl})-N~2~-(phenylmethyl)glycinamide	C				
94	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl})-2-[3,5-bis(trifluoromethyl)phenyl]acetamide	C	B	B	D	C
95	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl})-2-[2-chloro-5-(trifluoromethyl)phenyl]acetamide	A	A	B	B	B
96	N-{3-fluoro-4-[[6-(methyloxy)-7-{{[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl]oxy]phenyl}-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A

97	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(1,2,3,4-tetrahydroisoquinolin-1-ylmethyl)ethanediamide	C				
98	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[(2-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)methyl]ethanediamide	B	B	B	A	B
99	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N~2~-methyl-N~2~-{[3-(trifluoromethyl)phenyl]methyl}glycinamide	C				
100	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N~2~-methyl-N~2~-{2-[3-(trifluoromethyl)phenyl]ethyl}glycinamide	C				
101	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N~2~-methyl-N~2~-{(2-phenylethyl)glycinamide	C				
102	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-4-(phenylmethyl)imidazolidin-2-one	B	B	B	B	C
103	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridazin-3-yl)-N'-(4-fluorophenyl)propanediamide	A	C	B	A	C
104	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-chlorophenyl)propanediamide	B				
105	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(3-chlorophenyl)propanediamide	C				
106	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N~2~-methyl-N~2~-{(phenylmethyl)glycinamide	C				
107	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-chlorophenyl)propanediamide	B				
108	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[(methyloxy)imino]propanamide	B	B			
109	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[(ethyloxy)imino]propanamide	B	A			
110	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[[{(phenylmethyl)oxy]imino}propanamide	B	B			
111	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-1-(phenylmethyl)prolinamide	C	C			
112	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-3-[(4-methylphenyl)sulfonyl]-4-(phenylmethyl)imidazolidin-2-one	B	C	B	C	C
113	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-4-(phenylmethyl)imidazolidin-2-one	C	C			

114	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-4-(phenylmethyl)-4,5-dihydro-1,3-oxazol-2-amine	C	B			
115	6,7-bis(methyloxy)-4-({4-[4-(phenylmethyl)piperazin-1-yl]phenyl}oxy)quinoline	C	C			
116	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-4-(phenylmethyl)piperazin-2-one	C	C			
117	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N~2~-(phenylmethyl)alaninamide	C	C			
118	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N~2~-methyl-N~2~-(phenylmethyl)alaninamide	C	C			
119	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N~2~-(phenylmethyl)leucinamide	C	C			
120	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N~2~-methyl-N~2~-(phenylmethyl)leucinamide	C	C			
121	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N~2~-(phenylmethyl)valinamide	C	C			
122	N-[5-chloro-6-({6-(methyloxy)-4-[(piperidin-4-ylmethyl)oxy]quinolin-7-yl}oxy)pyridin-3-yl]-N'-(4-fluorophenyl)propanediamide	C	C			
123	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-4-(phenylmethyl)tetrahydropyrimidin-2(1H)-one	C	C			
124	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
125	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	A	B
126	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	C	C			
127	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-N~2~-methyl-N~2~-(phenylmethyl)valinamide	C	C			
128	(2E)-N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-2-[(phenyloxy)imino]propanamide	C	A			
129	(2E)-N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl}-2-phenyl-2-[[[(phenylmethyl)oxy]imino]ethanamide	B	C			
130	6,7-bis(methyloxy)-4-({4-[4-(phenylmethyl)piperidin-1-yl]phenyl}oxy)quinoline	C	C			
131	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-{[2-(1-methylethyl)-1,2,3,4-tetrahydroisoquinolin-1-yl]methyl}ethanediamide	B	B			
132	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[(2-ethyl-1,2,3,4-tetrahydroisoquinolin-1-yl)methyl]ethanediamide	B	C			

133	1,1-dimethylethyl 4-({[4-{{[3-chloro-5-{{[3-{{[4-fluorophenyl]amino}-3-oxopropanoyl}amino]pyridin-2-yl]oxy}-6-(methyloxy)quinolin-7-yl]oxy}methyl]piperidine-1-carboxylate	B	C			
134	N-[5-chloro-6-{{6-(methyloxy)-7-{{[piperidin-4-ylmethyl]oxy}quinolin-4-yl}oxy]pyridin-3-yl]-N'-(4-fluorophenyl)propanediamide	A	B	B	A	
135	N-{5-chloro-6-{{6-(methyloxy)-7-{{[1-methylpiperidin-4-yl]methyl]oxy}quinolin-4-yl]oxy]pyridin-3-yl}-N'-(4-fluorophenyl)propanediamide	A	B	B	A	
136	N-(4-{{[7-{{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	B	A
137	N-[3-fluoro-4-{{6-(methyloxy)-7-{{[3-morpholin-4-ylpropyl]oxy}quinolin-4-yl}oxy]phenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	B	A
138	N-[3-fluoro-4-{{6-(methyloxy)-7-{{[3-piperidin-1-ylpropyl]oxy}quinolin-4-yl}oxy]phenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	A	
139	N-(4-{{[7-{{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
140	N'-(3-fluoro-4-{{6-(methyloxy)-7-{{[1-methylpiperidin-4-yl]methyl]oxy}quinolin-4-yl]oxy]phenyl)-N-methyl-N-(2-phenylethyl)ethanediamide	A	A			B
141	N-(3-fluoro-4-{{[7-{{[[(3aR,5r,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}phenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
142	N-(3-fluoro-4-{{[7-{{[[(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}phenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
143	2-(3,4-dihydroisoquinolin-2(1H)-yl)-N-{3-fluoro-4-{{6-(methyloxy)-7-{{[1-methylpiperidin-4-yl]methyl]oxy}quinolin-4-yl]oxy]phenyl}-2-oxoacetamide	A	A			
144	N-[3-fluoro-4-{{6-(methyloxy)-7-{{[piperidin-4-ylmethyl]oxy}quinolin-4-yl}oxy]phenyl]-2-oxo-2-(3-phenylpyrrolidin-1-yl)acetamide	A	A	A	A	A
145	N-[3-fluoro-4-{{6-(methyloxy)-7-{{[piperidin-4-ylmethyl]oxy}quinolin-4-yl}oxy]phenyl]-2-oxo-2-(2-phenylmorpholin-4-yl)acetamide	A	B	B	B	

146	N-[2-(dimethylamino)-2-phenylethyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	B	A	
147	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(2-oxo-2-phenylethyl)ethanediamide	A	B	B	B	B
148	N-(6-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-5-chloropyridin-3-yl)-2,2-difluoro-N'-(4-fluorophenyl)propanediamide	C	C			
149	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy]quinolin-4-yl)oxy]phenyl}-N'-(phenylmethyl)ethanediamide	A	A	A	B	A
150	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-[2-(2-fluorophenyl)ethyl]ethanediamide	A	A	A	A	A
151	N-[2-(3-chlorophenyl)ethyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	A	A	A
152	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-{2-[2-(methyloxy)phenyl]ethyl}ethanediamide	A	A	A	A	A
153	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(2-pyridin-3-ylethyl)ethanediamide	A	B	B	B	
154	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(phenylmethyl)ethanediamide	A	A	A	B	A
155	N-{2-[2,5-bis(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	A	A	A
156	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-{2-[2-(trifluoromethyl)phenyl]ethyl}ethanediamide	A	A	A	A	C
157	N-{2-[2-(ethyloxy)phenyl]ethyl}-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	A	A	A
158	N-[2-(2,4-dimethylphenyl)ethyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	B	B	A	B
159	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-[(1S)-2-(4-methylphenyl)-1-phenylethyl]ethanediamide	B	C			
160	N-[2-(4-chlorophenyl)ethyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	A	A	B

161	((3-fluoro-4-((6-(methyloxy)-7-((1-methylpiperidin-4-yl)methyl)oxy)quinolin-4-yl)oxy)phenyl)amino(oxo)acetic acid	B	C			
162	N-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]-N'-[2-(3-fluorophenyl)ethyl]ethanediamide	A	A			A
163	N-[2-(2-chlorophenyl)ethyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	A	B	A
164	N-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]-N'-{2-[3-(methyloxy)phenyl]ethyl}ethanediamide	A	A	A	A	A
165	N-(1,2-diphenylethyl)-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	B	B	C
166	N-[2-(2,4-dichlorophenyl)ethyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	A	B	B	A	B
167	N-{2-[3,4-bis(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	A	B	B	B	
168	N-[2-(4-ethylphenyl)ethyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	A	B	B	B	C
169	N-{2-[4-(ethyloxy)phenyl]ethyl}-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	B	C	B	B	
170	N-{2-[4-(ethyloxy)-3-(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	A	B	C	B	
171	N-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]-N'-{2-[4-(phenyloxy)phenyl]ethyl}ethanediamide	B	C	C	C	
172	N-{2-[3-(ethyloxy)-4-(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]ethanediamide	A	C	B	B	
173	N-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]-N'-(2-pyridin-2-ylethyl)ethanediamide	A	A	A	B	B
174	N-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]-N'-(2-pyridin-4-ylethyl)ethanediamide	A	B	B	B	C
175	N-[3-fluoro-4-((6-(methyloxy)-7-((piperidin-4-ylmethyl)oxy)quinolin-4-yl)oxy)phenyl]-N'-[2-(4-fluorophenyl)ethyl]ethanediamide	A	A	A	A	A

176	N-[2-(2-bromophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediarnide	A	A	A	A	A
177	N-[2-(2-chloro-6-fluorophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediarnide	A	A	A	B	A
178	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(2R)-2-phenylpropyl]ethanediarnide	A	A	A	B	A
179	N-(2,3-dihydro-1H-inden-1-yl)-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediarnide	A	A	A	A	A
180	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(2-methylpropyl)ethanediarnide	A	B	B	B	
181	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(3-methylbutyl)ethanediarnide	A	B	B	B	B
182	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-[(2R)-2-phenylpropyl]ethanediarnide	A	A	A	A	A
183	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(2-phenylpropyl)ethanediarnide	A	A	A	A	A
184	N-(2,3-dihydro-1H-inden-2-yl)-N'-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}ethanediarnide	A	A	A	B	A
185	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1R)-1-phenylethyl]ethanediarnide	A	B	B	B	
186	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1S)-1-phenylethyl]ethanediarnide	A	B	A	B	C
187	N-[2-(3-bromophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediarnide	A	A	A	A	A
188	N-[2-(2,6-dichlorophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediarnide	A	A	A	A	A
189	N-[2-(1,3-benzodioxol-5-yl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediarnide	A	A	B	A	B

190	N-{5-chloro-6-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy)quinolin-4-yl]oxy}pyridin-3-yl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	A	A
191	N-{2-[3-bromo-4-(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	B	B	B
192	N-{2-[3,5-bis(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	B	B
193	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]-N'-[2-(2-methylphenyl)ethyl]ethanediamide	A	A	A	B	A
194	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]-N'-[2-(3-methylphenyl)ethyl]ethanediamide	A	A	A	A	A
195	N-{2-[3-(ethyloxy)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	B	B	B
196	N-[2-(3,4-dimethylphenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	B	B	B
197	N-[2-(2,5-dimethylphenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	A
198	N-{2-[3-chloro-4-(propyloxy)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	B	C			
199	N-{2-[4-(butyloxy)-3-chlorophenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	B	C			
200	N-{2-[4-(1,1-dimethylethyl)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	B	C			
201	N-{2-[4-(aminosulfonyl)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B
202	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]-N'-{2-[4-hydroxy-3-(methyloxy)phenyl]ethyl}ethanediamide	A	B	B	A	B
203	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]-N'-{2-[3-hydroxy-4-(methyloxy)phenyl]ethyl}ethanediamide	A	B	B	B	B
204	N-[(2,4-dichlorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl]oxy)quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	B

205	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[4-fluoro-2-(trifluoromethyl)phenyl]methyl}ethanediamide	A	A	B	A	A
206	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1R)-1-(4-methylphenyl)ethyl]ethanediamide	A	B	B	B	B
207	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[3-fluoro-4-(trifluoromethyl)phenyl]methyl}ethanediamide	A	B	B	B	B
208	N-[(3-chloro-4-fluorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	B
209	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{(1S)-1-[3-(methyloxy)phenyl]ethyl}ethanediamide	A	B	B	B	B
210	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1R)-1-naphthalen-2-ylethyl]ethanediamide	A	B	B	B	
211	N-{[4-chloro-3-(trifluoromethyl)phenyl]methyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	A
212	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1S)-1-(4-methylphenyl)ethyl]ethanediamide	A	B	C	B	
213	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[6-(trifluoromethyl)pyridin-3-yl]methyl}ethanediamide	A	B	C	B	B
214	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(2-methylphenyl)methyl]ethanediamide	A	A	A	A	B
215	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(3-methylphenyl)methyl]ethanediamide	A	A	A	B	A
216	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[4-fluoro-3-(trifluoromethyl)phenyl]methyl}ethanediamide	A	A	B	A	A
217	N-[(3,5-dichlorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	A
218	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1R)-1,2,3,4-tetrahydronaphthalen-1-yl]ethanediamide	A	B	B	B	A

219	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1S)-1,2,3,4-tetrahydronaphthalen-1-yl]ethanediamide	A	A	A	A	A
220	N-cyclopentyl-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B
221	N-[1-(4-bromophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B
222	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(2-fluorophenyl)methyl]ethanediamide	A	A	B	B	A
223	N-[2-(3,4-dichlorophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	
224	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(4-fluorophenyl)methyl]ethanediamide	A	A	A	A	A
225	N-[(2,3-difluorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	B	B	A
226	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[2-(phenyloxy)ethyl]ethanediamide	A	A	A	A	A
227	N-(2,2-diphenylethyl)-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	A	B
228	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[2-[4-(methyloxy)phenyl]ethyl]ethanediamide	A	B	B	B	B
229	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(2-phenylpropyl)ethanediamide	A	A	A	A	A
230	N-[2-(4-bromophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	B	B	B
231	N-(4-{[7-{{(1-ethylpiperidin-4-yl)methyl}oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-2-oxo-2-(2-phenylmorpholin-4-yl)acetamide	A	B	B	B	B
232	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[3-fluoro-5-(trifluoromethyl)phenyl]methyl}ethanediamide	A	A	B	A	B
233	N-[(3,5-difluorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	B	A

234	N-{[2-chloro-5-(trifluoromethyl)phenyl]methyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	B	A	B
235	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(dimethylamino)-2-phenylethyl]ethanediamide	B	B	B	A	
236	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[4-(methyloxy)phenyl]methyl}ethanediamide	A	A	A	B	B
237	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[4-(trifluoromethyl)phenyl]methyl}ethanediamide	A	B	B	B	A
238	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[3-(methyloxy)phenyl]methyl}ethanediamide	A	A	A	A	B
239	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[3-(trifluoromethyl)phenyl]methyl}ethanediamide	A	A	A	A	B
240	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[3-[(trifluoromethyl)oxy]phenyl]methyl}ethanediamide	A	A	A	A	A
241	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[2-(methyloxy)phenyl]methyl}ethanediamide	A	A	A	A	A
242	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[2-(trifluoromethyl)phenyl]methyl}ethanediamide	A	A	A	A	A
243	N-[(3-chlorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	B	B
244	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[2-[(trifluoromethyl)oxy]phenyl]methyl}ethanediamide	A	A	A	A	A
245	N-[(2-chlorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	A
246	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[4-[(trifluoromethyl)oxy]phenyl]methyl}ethanediamide	A	B	B	B	B
247	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy]quinolin-4-yl}oxy]phenyl}-N'-{[4-(methyloxy)phenyl]methyl}ethanediamide	A	B	A	B	
248	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy]quinolin-4-yl}oxy]phenyl}-N'-{[4-(trifluoromethyl)phenyl]methyl}ethanediamide	A	B	B	B	B

249	N-(4-{[7-[(azetidin-3-ylmethyl)oxy]-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-phenylethyl)ethanediamide	A	A			
250	N-(3-fluoro-4-{[7-[(1-methylazetidin-3-yl)methyl]oxy]-6-(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
251	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(2-hydroxy-2-phenylethyl)ethanediamide	B	B	B	B	
252	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2,4-difluorophenyl)propanediamide	A	C			
253	N'-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N-(4-fluorophenyl)-N-methylpropanediamide	B	C			
254	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-[(1R)-1-phenylpropyl]ethanediamide	A	B	B	B	B
255	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-[(1S)-1-phenylpropyl]ethanediamide	A	B	C	B	
256	N-[(3,4-difluorophenyl)methyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	A	B	A
257	N-[(2,6-difluorophenyl)methyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	A	A	B	A
258	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy]quinolin-4-yl]oxy}phenyl}-N'-[2-(4-fluorophenyl)ethyl]ethanediamide	A	A	A	A	A
259	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy]quinolin-4-yl]oxy}phenyl}-N'-phenylethanediamide	A	B	C	C	B
260	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(3-fluorophenyl)ethanediamide	A	C	B	C	C
261	N-(3-chloro-4-fluorophenyl)-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	C	C	B	B
262	N-[3,4-bis(methyloxy)phenyl]-N'-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]ethanediamide	A	C	C	B	B
263	N-[3-fluoro-4-((6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(3-methylbutyl)ethanediamide	A	B	A	B	B

264	N-(3,3-dimethylbutyl)-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	B
265	N-[5-chloro-6-({6-(methyloxy)-7-[(3-piperidin-1-ylpropyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]-N'-(4-fluorophenyl)propanediamide	A	B	B	A	B
266	N-[5-chloro-6-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]-N'-(4-fluorophenyl)propanediamide	A	B	B	A	B
267	N-(5-chloro-6-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl)-N'-(4-fluorophenyl)propanediamide	A	B	B	A	B
268	N-[(4-chlorophenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A			
269	N-{[3,5-bis(methyloxy)phenyl]methyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A		A	
270	N-[(4-butylphenyl)methyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B
271	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[2-(4-methylphenyl)ethyl]ethanediamide	A	B	B	A	B
272	N-{[3,5-bis(trifluoromethyl)phenyl]methyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	C	B	
273	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(pyrazin-2-ylmethyl)ethanediamide	B	C	C	B	
274	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(pyridin-2-ylmethyl)ethanediamide	A	B	B	B	B
275	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
276	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinazolin-4-yl]oxy}phenyl}-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
277	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{[2-fluoro-3-(trifluoromethyl)phenyl]methyl}ethanediamide	A	A	A	A	A
278	N-{2-[2-bromo-6-(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	A

279	N-{2-[3,4-bis(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N-methylethanediamide	B	B	B	A	B
280	N-{2-[5-bromo-2-(methyloxy)phenyl]ethyl}-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	A	A	A	A
281	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-{2-fluoro-5-(trifluoromethyl)phenyl}methyl}ethanediamide	A	A	B	A	B
282	N-[5-chloro-6-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
283	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[1-(4-fluorophenyl)ethyl]ethanediamide	A	A	B	A	B
284	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[(1S)-2-oxo-1-(phenylmethyl)-2-pyrrolidin-1-ylethyl]ethanediamide	A	C	B	B	C
285	N-{3-fluoro-4-[(6-(methyloxy)-7-[(3aR,6aS)-octahydrocyclopenta[c]pyrrol-5-ylmethyl]oxy}quinazolin-4-yl]oxy}phenyl}-N'-(2-phenylethyl)ethanediamide	A	A	A	A	A
286	N-[2-(4-aminophenyl)ethyl]-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B
287	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-2-oxo-2-[4-(phenylmethyl)piperidin-1-yl]acetamide	A	A	B	A	B
288	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)propanediamide	A	A	A	A	A
289	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	5.581
290	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloropyridin-3-yl)-N'-(3-fluorophenyl)propanediamide	B	C			
291	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloropyridin-3-yl)-N'-phenylpropanediamide	A	C	B	A	C
292	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)-2,2-dimethylpropanediamide	B	B	B	A	B
293	N-ethyl-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B

294	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(1-methylethyl)ethanediamide	A	B	B	B	C
295	N-butyl-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B
296	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-[2-(methyloxy)ethyl]ethanediamide	B	B	B	B	C
297	N-(cyclopropylmethyl)-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]ethanediamide	A	B	B	B	B
298	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(2-morpholin-4-ylethyl)ethanediamide	B	A	B	A	B
299	N-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-2-oxo-2-pyrrolidin-1-ylacetamide	A	B	B	B	B
300	N-ethyl-N'-[3-fluoro-4-({6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl}oxy)phenyl]-N-methylethanediamide	A	B	C	B	B
301	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(phenylmethyl)cyclopropane-1,1-dicarboxamide	A	C	B	B	B
302	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-phenylethyl)cyclopropane-1,1-dicarboxamide	C	C			
303	N-{4-[(7-chloroquinolin-4-yl)oxy]-3-fluorophenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	C	C	B
304	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
305	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-phenylcyclopropane-1,1-dicarboxamide	A	B	B	A	
306	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-methylpyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	B	B	C	B
307	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A	A	B	A
308	N-{4-[(7-chloroquinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	C	B
309	N-[5-chloro-6-({6-(methyloxy)-7-[(phenylmethyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	C	C	B	C

310	N-(4-{[6,7-bis(methyloxy)quinazolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
311	N-(4-{[6,7-bis(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	A	A
312	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
313	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperidin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
314	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperidin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A	A	B	A
315	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
316	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy)quinazolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
317	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	C	A
318	N-(4-fluorophenyl)-N'-[2-methyl-6-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]cyclopropane-1,1-dicarboxamide	A	A	B	B	B
319	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
320	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloro-2-methylpyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	C	C	B	C
321	N-[3-fluoro-4-({7-(methyloxy)-6-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	A	A
322	N-[3-fluoro-4-({7-(methyloxy)-6-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
323	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3,5-difluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	A	A
324	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2,5-difluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A

325	N-{3-fluoro-4-[(7-(methyloxy)-6-[(1-methylpiperidin-4-yl)methyl]oxy)quinazolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	A	A
326	N-[5-fluoro-2-methyl-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	B	B
327	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2,3,5-trifluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	B	B	B
328	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-2-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	B	B	C	C
329	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-chloro-5-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	B	B	C	B
330	N-(3-fluoro-4-{[6-hydroxy-7-(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
331	N-(4-fluorophenyl)-N'-[2-methyl-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]cyclopropane-1,1-dicarboxamide	A	A	A	B	A
332	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
333	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
334	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy)quinolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
335	N-(4-fluorophenyl)-N'-[4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]cyclopropane-1,1-dicarboxamide	A	A	A	A	A
336	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A	A	A	A
337	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-chloro-5-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	B	B	C	C
338	N-(4-{[6,7-bis(methyloxy)-2-(methylthio)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	C	C			

339	N-(4-fluorophenyl)-N'-(4-{[2-methyl-6,7-bis(methyloxy)quinazolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide	C	C			
340	N-(4-{[2-amino-6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	C	C			
341	N-(3-fluoro-4-{[2-(methylamino)-6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	C	C			
342	(1S,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A	B	A	A
343	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A	A	A	A
344	N-(4-{[6-{[3-(diethylamino)propyl]oxy}-7-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A		A	
345	N-(4-{[6-{[2-(diethylamino)ethyl]oxy}-7-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A		A	
346	1,1-dimethylethyl 4-(3-{[4-[(2-fluoro-4-{[(1-{[(4-fluorophenyl)amino]carbonyl}cyclopropyl)carbonyl]amino}phenyl)oxy]-6-(methyloxy)quinolin-7-yl]oxy}propyl)piperazine-1-carboxylate	A	A		B	
347	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A			
348	(1R,2R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		B	
349	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A		A	
350	N-(4-{[7-{[3-(4-acetylpiperazin-1-yl)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A		A	
351	1,1-dimethylethyl 4-(3-{[4-[(2-fluoro-4-{[(1R,2R)-1-{[(4-fluorophenyl)amino]carbonyl}-2-methylcyclopropyl)carbonyl]amino}phenyl)oxy]-6-(methyloxy)quinolin-7-yl]oxy}propyl)piperazine-1-carboxylate	A	A		B	

352	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)-1-(phenylmethyl)azetidine-3,3-dicarboxamide	A	C		C	
353	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)azetidine-3,3-dicarboxamide	B	C		C	
354	(1R,2S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
355	(1R,2R)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
356	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
357	N-(3-fluoro-4-{[7-({3-[4-(1-methylethyl)piperazin-1-yl]propyl}oxy)-6-(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A		A	
358	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	A	A		A	
359	(1R,2R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
360	(1R,2R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
361	(1R,2S)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
362	(1R,2S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
363	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A			
364	(1R,2S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	


365	(1r,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	B		B	
366	(1r,2R,3S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	B		B	
367	(1r,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	B	A		B	
368	(1r,2R,3S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	A		B	
369	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	B	A		B	
370	(2R,3R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	A		B	
371	(2R,3R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	A		B	
372	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	A	A		B	
373	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	A	A		C	
374	(1r,2R,3S)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	B		B	
375	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	A	A		C	
376	(1r,2R,3S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	B		B	

377	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	A	A		B	
378	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	A	B		C	
379	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	A	A		C	
380	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A		B	
381	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A		B	
382	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A		B	
383	(2R,3R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	A		C	
384	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A		C	
385	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	A	A		C	
386	(1R,2R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
387	(1R,2R)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
388	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	B	A		C	

389	(2R,3R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	A		B	
390	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	A	A		A	
391	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	A	A		B	
392	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[(4-fluorophenyl)methyl]cyclopropane-1,1-dicarboxamide	B	B		A	
393	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(2-morpholin-4-ylethyl)cyclopropane-1,1-dicarboxamide	C	D		B	
394	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	B	D		B	
395	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	B	C		B	
396	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	B	A		A	
397	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	B	C		B	
398	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-phenylcyclopropane-1,1-dicarboxamide	A	A		A	
399	N-[3-(aminomethyl)phenyl]-N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide	B	A		B	
400	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	B	A			
401	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	A	A			

What is claimed is:

1. A compound for modulating kinase activity according to formula I,





or a pharmaceutically acceptable salt, hydrate, or prodrug thereof, wherein,

R^1 is selected from -H, halogen, $-OR^3$, $-NO_2$, $-NH_2$, $-NR^3R^4$, and optionally substituted lower alkyl;

A^1 is selected from $=N-$, $=C(H)-$, and $=C(CN)-$;

Z is selected from $-S(O)_{0-2}-$, $-O-$, and $-NR^5-$;

Ar is either a group of formula II, or of formula III,



wherein,

R^2 is selected from -H, halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted lower alkyl;

q is 0 to 4;

G is a group -B-L-T, wherein

B is selected from absent, $-N(R^{13})-$, $-N(SO_2R^{13})-$, $-O-$, $-S(O)_{0-2}-$, and $-C(=O)-$;

L is selected from absent, $-C(=S)N(R^{13})-$, $-C(=NR^{14})N(R^{13})-$, $-SO_2N(R^{13})-$, $-SO_2-$, $-C(=O)N(R^{13})-$, $-N(R^{13})-$, $-C(=O)C_{1-2}alkylN(R^{13})-$, $-N(R^{13})C_{1-2}alkylC(=O)-$, $-C(=O)C_{0-1}alkylC(=O)N(R^{13})-$, $-C_{0-4}alkylene-$, $-C(=O)C_{0-1}alkylC(=O)OR^3-$, $-C(=NR^{14})C_{0-1}alkylC(=O)-$, $-C(=O)-$, $-C(=O)C_{0-1}alkylC(=O)-$, and an optionally substituted four to six-membered heterocyclyl containing

between one and three annular heteroatoms including at least one nitrogen;
and

T is selected from -H, -R¹³, -C₀₋₄alkyl, -C₀₋₄alkylQ, -OC₀₋₄alkylQ, -C₀₋₄alkylOQ, -N(R¹³)C₀₋₄alkylQ, -SO₂C₀₋₄alkylQ, -C(=O)C₀₋₄alkylQ, -C₀₋₄alkylN(R¹³)Q, and -C(=O)N(R¹³)C₀₋₄alkylQ, wherein each of the aforementioned C₀₋₄alkyl is optionally substituted;

J is selected from -S(O)₀₋₂-, -O-, and -NR¹⁵-;

R³ is -H or R⁴;

R⁴ is selected from optionally substituted lower alkyl, optionally substituted aryl, optionally substituted lower arylalkyl, optionally substituted heterocyclyl, and optionally substituted lower heterocyclylalkyl; or


R³ and R⁴, when taken together with a common nitrogen to which they are attached, form an optionally substituted five- to seven-membered heterocyclyl, said optionally substituted five- to seven-membered heterocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

A² and A³ are each independently selected from =N-, =C(R²)-;

R⁵ is -H or optionally substituted lower alkyl;

D is selected from -O-, -S(O)₀₋₂-, and -NR¹⁵-;

R⁵⁰ is either R³, or according to formula IV;



IV

wherein X¹, X², and optionally X³, represent the atoms of a saturated bridged ring system, said saturated bridged ring system comprising up to four annular heteroatoms represented by any of X¹, X², and X³; wherein,

each X¹ is independently selected from -C(R⁶)R⁷-, -O-, -S(O)₀₋₂-, and -NR⁸-;

each X² is independently an optionally substituted bridgehead methine or a bridgehead nitrogen;

each X³ is independently selected from -C(R⁶)R⁷-, -O-, -S(O)₀₋₂-, and -NR⁸-;

Y is either:

an optionally substituted lower alkylene linker, between D and either 1) any annular atom of the saturated bridged ring system, except X^2 when X^2 is a bridgehead nitrogen, or 2) any heteroatom, represented by any of R^6 or R^7 ; provided there are at least two carbon atoms between D and any annular heteroatom of the saturated bridged ring system or any heteroatom represented by any of R^6 or R^7 ;

or Y is absent, when Y is absent, said saturated bridged ring system, is directly attached to D via an annular carbon of said saturated bridged ring system, unless D is $-SO_2-$, in which case said saturated bridged ring system, is directly attached to D via an any annular atom of said saturated bridged ring system;

m and p are each independently 1-4;

n is 0-2, when n = 0, then there is a single bond between the two bridgehead X^2 's;

R^6 and R^7 are each independently selected from -H, halogen, trihalomethyl, -CN, $-NH_2$, $-NO_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^4$, $-SO_2NR^3R^4$, $-CO_2R^3$, $-C(O)NR^3R^4$, $-N(R^3)SO_2R^4$, $-N(R^3)C(O)R^3$, $-NCO_2R^3$, $-C(O)R^3$, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted lower arylalkyl, optionally substituted heterocyclyl, optionally substituted lower heterocyclylalkyl, and a bond to either Y or D; or

R^6 and R^7 , when taken together are oxo; or

R^6 and R^7 , when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

R^8 is selected from $-R^3$, Y, $-SO_2NR^3R^4$, $-CO_2R^4$, $-C(O)NR^3R^3$, $-SO_2R^4$, and $-C(O)R^3$;

R^{13} is selected from -H, $-C(=O)R^3$, $-C(=O)OR^3$, $-C(=O)SR^3$, $-SO_2R^4$, $-C(=O)N(R^3)R^3$, and optionally substituted lower alkyl,

two R^{13} , together with the atom or atoms to which they are attached, can combine to form a heteroalicyclic optionally substituted with between one and four of R^{60} , said heteroalicyclic can have up to four annular heteroatoms, and said heteroalicyclic can have

an aryl or heteroaryl fused thereto, in which case said aryl or heteroaryl is optionally substituted with an additional one to four of R^{60} ;

R^{14} is selected from -H, -NO₂, -NH₂, -N(R³)R⁴, -CN, -OR³, optionally substituted lower alkyl, optionally substituted heteroalicycylalkyl, optionally substituted aryl, optionally substituted arylalkyl and optionally substituted heteroalicyclic;

R^{15} is a group -M¹-M², wherein M¹ is selected from absent, -C(=S)N(R¹³)-, -C(=NR¹⁴)N(R¹³)-, -SO₂N(R¹³)-, -SO₂-, -C(=O)N(R¹³)-, -C(=O)C(=O)N(R¹³)-, -C₀₋₄alkylene-, -C(=O)-, and an optionally substituted four to six-membered heterocycl annular containing between one and three heteratoms including at least one nitrogen; and M² is selected from -H, -C₀₋₆alkyl, alkoxy, -C(=O)C₀₋₄alkylQ, -C₀₋₄alkylQ, -OC₀₋₄alkylQ-, -N(R¹³)C₀₋₄alkylQ-, and -C(=O)N(R¹³)C₀₋₄alkylQ; and

Q is a five- to ten-membered ring system, optionally substituted with between zero and four of R^{20} ;

R^{20} is selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl;

R^{60} is selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted heteroarylalkyl, and optionally substituted arylalkyl;

two of R^{60} , when attached to a non-aromatic carbon, can be oxo;

with the proviso, only when Ar is according to formula II, if Y is a C₁₋₆ alkylene; Z is -NH- or -N(CH₃)-; R¹ is a C₁₋₆alkyl optionally substituted in the 2-position by -OH or a C₁₋₄alkoxy group; R² is -H or halogen; n = 0; and the atoms, X¹, of one bridge of the saturated bridged ring system, when combined with both bridgehead atoms, X², of the saturated bridged ring system, represent:


- 1) either a pyrrolidine or a piperidine, and any atom, X¹ or X², of either of said pyrrolidine or said piperidine is attached to Y, then the other bridge of said saturated bridged ring system cannot be any one of -OC(O)CH₂-, -CH₂OC(O)-, -OC(O)CH₂CH₂-, -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, -OC(O)CH₂NH-, -OC(O)CH₂N(C₁₋₄alkyl)-, and -OC(O)CH₂O-; or

- 2) either a piperazine or a 4-(C₁₋₄alkyl)-piperazine, and any atom, X¹ or X², of either of said piperazine or said 4-(C₁₋₄alkyl)-piperazine is attached to Y, then the other bridge of said saturated bridged ring system, only when attached via the 2- and the 3-position of either of said piperazine or said 4-(C₁₋₄alkyl)-piperazine, cannot be one of -CH₂OC(O)CH₂-, -CH₂CH₂OC(O)-, and either of the two aforementioned bridges optionally substituted by one or two C₁₋₂alkyl groups; or
- 3) a piperazine, and any atom, X¹ or X², of said piperazine is attached to Y, then the other bridge of said saturated bridged ring system, only when attached via the 3- and the 4-position of said piperazine, cannot be one of -C(O)OCH₂CH₂-, -CH₂OC(O)CH₂-, and either of the two aforementioned bridges optionally substituted by one or two C₁₋₂alkyl groups, and only when either of the two aforementioned bridges are attached to the 3-position of said piperazine via their left-hand end as depicted above; or
- 4) a 2-oxomorpholine, said 2-oxomorpholine attached to Y via its 4-position, then the other bridge of said saturated bridged ring system, only when attached via the 5- and the 6-position of said 2-oxomorpholine, cannot be one of -(CH₂)_g-, -CH₂WCH₂-, -CH₂WCH₂CH₂-, and -CH₂CH₂WCH₂-, wherein W is -O-, -S(O)₀₋₂-, -NH-, or -N(C₁₋₄alkyl)- wherein g is 2, 3, or 4;

and with the proviso that when Z is -O-, Ar is according to formula II, and the portion of G directly attached to Ar is selected from:

then R⁵⁰ must be of formula IV;

and with the proviso that when Ar is phenylene or substituted phenylene, Z is $-S(O)_{0-2}-$ or





$-O-$, then the portion of G directly attached to Ar cannot contain

when R^{70} is selected from $-H$, C_{1-4} alkyl, and C_{1-4} alkoxyl.

2. The compound according to claim 1, wherein In one example, the compound is according to paragraph [0033], wherein Z is either $-O-$ or $-NR^5-$.




3. The compound according to claim 2, wherein G is selected from the following:





wherein wherein Q, R^{20} , and R^{13} are as defined above; each E is selected from -O-, -N(R^{13})-, -CH₂-, and -S(O)₀₋₂-; M is selected from -O-, -N(R^{13})-, -CH₂-, and -C(=O)N(R^{13})-; each V is independently either =N- or =C(H)-; each methylene in any of the above formulae is independently optionally substituted with R^{25} ; and R^{25} is selected from halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic, two of R^{25} on a single carbon can be oxo.

4. The compound according to claim 3, wherein Ar is according to one of formula **IIa**, **IIb**, and **IIIa**.

**IIa****IIb****IIIa**


5. The compound according to claim 4, wherein D is -O- and R^1 is -OR³.

6. The compound according to claim 5, wherein -O-R⁵⁰ and R^1 are interchangeably located at the 6-position and 7-position of the quinazoline or quinoline according to formula **I**.

7. The compound according to claim 6, wherein R^1 is -OH or -OC₁₋₆alkyl.


8. The compound according to claim 7, wherein A^1 is =N- or =C(H)-.

9. The compound according to claim 8, wherein G is selected from:




wherein Q, R²⁰, R¹³, E, and R⁶⁰ are as defined above; each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R²⁵, and R²⁵ is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic.

10. The compound according to claim 9, wherein Q is selected from:




wherein R²⁰ is defined as above, and P is a five- to seven-membered ring, including the two shared carbons of the aromatic ring to which P is fused, P optionally containing between one and three heteroatoms.

11. The compound according to claim 10, wherein Ar is according to formula **IIa**, and G is selected from:



wherein Q, R²⁰, R¹³, E, and R⁶⁰ are as defined above, and each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R²⁵; and R²⁵ is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic.

12. The compound according to claim 10, wherein Ar is according to formula **IIb**, and G is selected from:



wherein Q, R²⁰, R¹³, E, and R⁶⁰ are as defined above, and each methylene in any of the above formulae, other than those depicted in a ring, is independently optionally substituted with R²⁵; and R²⁵ is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic.

13. The compound according to claim 12, wherein the methylene between the two carbonyls of the depicted formulae is di-substituted with either optionally substituted lower alkyl, or an optionally substituted spirocycle.

14. The compound according to claim 11 or claim 12, wherein R⁵⁰ is a heteroalicyclic or a C₁₋₆alkyl-heteroalicyclic.

15. The compound according to claim 14, wherein at least one of R² is halogen.

16. The compound according to claim 14, wherein R⁵⁰ is according to formula IV.

17. The compound according to claim 16, wherein the saturated bridged ring system according to formula IV has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], [3.1.0], [3.3.3], [3.3.2], [3.3.1], [3.2.2], [3.2.1], [2.2.2], and [2.2.1].


18. The compound according to claim 17, wherein Y is selected from -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂-, -CH₂-, and absent.

19. The compound according to claim 18, wherein n is 0 and the saturated bridged ring system according to formula IV has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], and [3.1.0].

20. The compound according to claim 19, wherein said saturated bridged ring system contains at least one annular nitrogen or at least one annular oxygen.

21. The compound according to claim 20, wherein said saturated bridged ring system contains -NR⁸-, wherein R⁸ is selected from -H, optionally substituted lower alkyl, -CO₂R³, -C(O)NR³R³, -SO₂R³, and -C(O)R³.


22. The compound according to claim 20, wherein said saturated bridged ring system is of formula V,



V

wherein U^1 is selected from -O-, $-S(O)_{0-2}-$, $-NR^8-$, $-CR^6R^7-$, and absent; and e is 0 or 1.

23. The compound according to claim 22, wherein Y is $-CH_2-$.
24. The compound according to claim 23, wherein U^1 is $-NR^8-$, wherein R^8 is selected from -H, optionally substituted lower alkyl, $-CO_2R^3$, $-C(O)NR^3R^3$, $-SO_2R^3$, and $-C(O)R^3$.
25. The compound according to claim 23, wherein U^1 is -O-.
26. The compound according to claim 23, wherein U^1 is absent.
27. The compound according to claim 20, wherein Y is selected from $-CH_2CH_2-$, $-CH_2-$, and absent.
28. The compound according to claim 27, wherein said saturated bridged ring system is of formula VI,



VI

wherein R^9 , R^{10} , and R^{11} are each independently selected from -H, and $-OR^{12}$; or

R^9 is selected from -H, and $-OR^{12}$, and R^{10} and R^{11} , when taken together, are either an optionally substituted alkylidene or an oxo;

R^{12} is selected from -H, $-C(O)R^3$, optionally substituted lower alkylidyne, optionally substituted lower arylalkylidyne, optionally substituted lower heterocyclylalkylidyne, optionally substituted lower alkylidene, optionally

substituted lower alkylidenearyl, optionally substituted lower alkylideneheterocyclyl, optionally substituted lower alkyl, optionally substituted lower alkylaryl, optionally substituted aryl, optionally substituted lower heterocyclalkyl, and optionally substituted heterocyclyl;


or two R^{12} 's, when taken together, form 1) a corresponding spirocyclic ketal when said two R^{12} 's stem from R^{10} and R^{11} , or 2) a corresponding cyclic ketal when said two R^{12} 's stem from R^9 and one of R^{10} and R^{11} .

29. The compound according to claim 28, wherein one of R^{10} and R^{11} is $-OR^{12}$, wherein R^{12} is selected from $-H$, $-C(O)R^3$, and optionally substituted lower alkyl; and R^9 and the other of R^{10} and R^{11} are both $-H$.

30. The compound according to claim 29, wherein Y is either $-CH_2-$ or absent.

31. The compound according to claim 30, wherein R^9 is an alkyl group containing at least one fluorine substitution thereon.

32. The compound according to claim 21, wherein said saturated bridged ring system is of formula **VII**.



VII

33. The compound according to claim 32, wherein Y is either $-CH_2-$ or absent.

34. The compound according to claim 33, wherein R^8 is methyl or ethyl.

35. The compound according to claim 21, wherein said saturated bridged ring system is of formula **VIII**.




VIII

36. The compound according to claim 35, wherein Y is $-CH_2-$.

37. The compound according to claim 36, wherein R^8 is methyl or ethyl.

38. The compound according to claim 20, wherein said saturated bridged ring system is of formula **IX**

**IX**

wherein U² is selected from -O-, -S(O)₀₋₂-, -NR⁸-, -CR⁶R⁷-, and absent.

39. The compound according to claim 38, wherein R³ of formula **IX** is selected from -H and optionally substituted alkyl.

40. The compound according to claim 39, wherein U² is either -CR⁶R⁷- or absent.

41. The compound according to claim 40, wherein U² is either -CH₂- or absent.

42. The compound according to claim 41, wherein Y is -CH₂-.

43. The compound according to claim 21, wherein said saturated bridged ring system is according to formula **X**.

**X**

44. The compound according to claim 43, wherein R⁸ is methyl or ethyl.

45. The compound according to claim 1, selected from Table 1.

Table 1

Entry	Name	Structure
-------	------	-----------

Table 1

Entry	Name	Structure
1	N-[(3-fluoro-4-[(6-(methyloxy)-7-[[[(3aR,6aS)-octahydrocyclopenta[c]pyrrol-5-ylmethyl]oxy}quinazolin-4-yl]oxy]phenyl]amino)carbonylthioyl]-2-phenylacetamide	
2	N-{[(3-fluoro-4-{[7-(([(3aR,6aS)-2-methyloctahydrocyclopenta[c]pyrrol-5-yl]methyl]oxy)-6-(methyloxy)quinazolin-4-yl]oxy}phenyl]amino]carbonylthioyl}-2-phenylacetamide	
3	N-{[(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)(methyl)amino]carbonylthioyl}-2-phenylacetamide	
4	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)imidazolidin-2-one	
5	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-(phenylmethyl)imidazolidin-2-one	
6	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-(phenylacetyl)imidazolidin-2-one	

Table 1







Entry	Name	Structure
7	ethyl [(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)amino](oxo)acetate	
8	N-{[(4-{[6,7-bis(methyloxy)quinazolin-4-yl]amino}-3-fluorophenyl)amino]carbonothioyl}-2-phenylacetamide	
9	N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N-methyl-N-(2-phenylethyl)sulfamide	
10	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-3-(phenylmethyl)-1,2,4-oxadiazol-5-amine	
11	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)piperidin-2-one	
12	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(phenylmethyl)ethanediamide	

Table 1

Entry	Name	Structure
13	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-4-phenyl-1,3-thiazol-2-amine	
14	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-phenylethyl)ethanediamide	
15	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-1-phenylmethanesulfonamide	
16	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-2-phenylethanesulfonamide	
17	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(phenylmethyl)benzenesulfonamide	
18	4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(phenylmethyl)benzenesulfonamide	

Table 1

Entry	Name	Structure
19	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(2-phenylethyl)benzenesulfonamide	
20	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(2-phenylethyl)benzenesulfonamide	
21	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-(3-phenylpropyl)benzenesulfonamide	
22	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}pyrrolidin-2-one	
23	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl (phenylmethyl)carbamate	
24	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl (2-phenylethyl)carbamate	

Table 1

Entry	Name	Structure
25	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-methyl-N-(3-phenylpropyl)benzenesulfonamide}	
26	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-phenylethanediamide}	
27	N-{{[(3-fluoro-4-{{[7-{{[(2-methyloctahydrocyclopenta[c]pyrrol-5-yl)methyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}phenyl]amino]carbonylthioyl]-2-phenylacetamide}	
28	N-[(Z)-[4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl]amino](imino)methyl]-2-phenylacetamide}	
29	4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluoro-N-[2-(phenyloxy)ethyl]benzenesulfonamide}	
30	N,N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-bis(3-phenylpropane-1-sulfonamide)	

Table 1







Entry	Name	Structure
31	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-3-phenylpropane-1-sulfonamide	
32	N2-[(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)sulfonyl]-N1-phenylglycinamide	
33	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl}-2-phenylacetamide	
34	N-{{[6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridin-3-yl}amino]carbonothioyl}-2-phenylacetamide	
35	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-1,3-benzothiazol-2-amine	
36	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-amine	

Table 1

Entry	Name	Structure
37	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-2-phenylacetamide	
38	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2-morpholin-4-ylethyl)ethanediamide	
39	benzyl-{[4-(6,7-dimethoxyquinolin-4-yloxy)-3-fluorophenylcarbamoyl]-methyl}-carbamic acid tert-butyl ester	
40	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-(phenylmethyl)glycinamide	
41	N2-acetyl-N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-(phenylmethyl)glycinamide	
42	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-1,3-benzothiazol-2-yl)-2-phenylacetamide	

Table 1







Entry	Name	Structure
43	benzyl-{{6-(6,7-dimethoxyquinolin-4-yloxy)-pyridin-3-ylcarbamoyl}-methyl}-carbamic acid tert-butyl ester	
44	N1-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}pyridin-3-yl)-N2-(phenylmethyl)glycinamide	
45	N2-acetyl-N1-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}pyridin-3-yl)-N2-(phenylmethyl)glycinamide	
46	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}pyridin-3-yl)-3-phenylpropanamide	
47	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}pyridin-3-yl)-4-phenylbutanamide	
48	N1-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}pyridin-3-yl)-N2-methyl-N2-(phenylmethyl)glycinamide	

Table 1

Entry	Name	Structure
49	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-{2-[4-(methyloxy)phenyl]ethyl}ethanediamide	
50	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-methyl-N2-(phenylmethyl)glycinamide	
51	4-[(2-amino-1,3-benzothiazol-6-yl)oxy]-6,7-bis(methyloxy)-1-(2-oxo-2-phenylethyl)quinolinium	
52	N-[[4-{[6,7-bis(methyloxy)quinolin-4-yl]amino}phenyl]amino]carbonothioyl]-2-phenylacetamide	
53	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-3-phenylpropanamide	
54	N-[[6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl]amino]carbonothioyl]-2-phenylacetamide	

Table 1

Entry	Name	Structure
55	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2,3-dihydro-1H-inden-1-yl)ethanediamide	
56	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(2,3-dihydro-1H-inden-2-yl)ethanediamide	
57	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(1,2,3,4-tetrahydronaphthalen-1-yl)ethanediamide	
58	N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N-(2-phenylethyl)-N-(phenylmethyl)sulfamide	
59	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-(trifluoroacetyl)glycinamide	
60	N-{[4-(6,7-dimethoxyquinolin-4-yloxy)-3-fluorophenylcarbamoyl]-methyl}-benzamide	

Table 1

Entry	Name	Structure
61	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}pyridin-3-yl)-N'-(4-fluorophenyl)propanediamide	
62	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-[(2S)-1,2,3,4-tetrahydronaphthalen-2-yl]ethanediamide	
63	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-[2-(4-methylphenyl)ethyl]ethanediamide	
64	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-(2-phenylpropyl)ethanediamide	
65	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N'-[2-(4-chlorophenyl)ethyl]ethanediamide	
66	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-3-fluorophenyl)-N,N'-bis(phenylmethyl)sulfamide	

Table 1

Entry	Name	Structure
67	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N,N'-bis(2-phenylethyl)sulfamide	
68	ethyl [(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)amino](oxo)acetate	
69	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-phenylethyl)ethanediamide	
70	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)propanediamide	
71	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(1,2,3,4-tetrahydronaphthalen-2-yl)ethanediamide	
72	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[2-(1-methylpyrrolidin-2-yl)ethyl]ethanediamide	

Table 1

Entry	Name	Structure
73	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-[2-(phenyloxy)ethyl]ethanediamide	
74	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-[2-hydroxy-1-(phenylmethyl)ethyl]urea	
75	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-3-[(4-methylphenyl)sulfonyl]-4-(phenylmethyl)imidazolidin-2-one	
76	N'-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N-methyl-N-(2-phenylethyl)ethanediamide	
77	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-{{[3-(trifluoromethyl)phenyl]methyl}ethanediamide	
78	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N'-{2-[3-(trifluoromethyl)phenyl]ethyl}ethanediamide	

Table 1

Entry	Name	Structure
79	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl}-3-oxo-4-phenylbutanamide	
80	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl}-2-[3-(trifluoromethyl)phenyl]acetamide	
81	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-[2-(phenyloxy)ethyl]-1,3-benzothiazol-2-amine	
82	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-(2-piperidin-1-ylethyl)-1,3-benzothiazol-2-amine	
83	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-methyl-N-(2-phenylethyl)-1,3-benzothiazol-2-amine	
84	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-(2-pyrrolidin-1-ylethyl)-1,3-benzothiazol-2-amine	

Table 1

Entry	Name	Structure
85	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-{{[3-(trifluoromethyl)phenyl]methyl}}-1,3-benzothiazol-2-amine	
86	6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-N-{{2-[[3-(trifluoromethyl)phenyl]ethyl}}-1,3-benzothiazol-2-amine	
87	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl}-N'-[[3-(trifluoromethyl)phenyl]propyl]diacetamide	
88	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl}-2-[[3-(trifluoromethyl)phenyl]acetyl]acetamide	
89	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N2-{{[3-(trifluoromethyl)phenyl]methyl}}glycinamide	
90	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl}-N2-(2-phenylethyl)glycinamide	

Table 1

Entry	Name	Structure
91	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-{2-[3-(trifluoromethyl)phenyl]ethyl}glycinamide	
92	benzyl-[[5-chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]carbonyl]-methyl}-carbamic acid tert-butyl ester	
93	N1-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N2-(phenylmethyl)glycinamide	
94	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-2-[3,5-bis(trifluoromethyl)phenyl]acetamide	
95	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-fluoro-1,3-benzothiazol-2-yl)-2-[2-chloro-5-(trifluoromethyl)phenyl]acetamide	
96	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy)quinolin-4-yl]oxy}phenyl}-N'-(2-phenylethyl)ethanediamide	

Table 1

Entry	Name	Structure
97	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(1,2,3,4-tetrahydroisoquinolin-1-ylmethyl)ethanediamide	
98	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-[(2-methyl-1,2,3,4-tetrahydroisoquinolin-1-yl)methyl]ethanediamide	
99	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-methyl-N2-{[3-(trifluoromethyl)phenyl]methyl}glycinamide	
100	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-methyl-N2-{2-[3-(trifluoromethyl)phenyl]ethyl}glycinamide	
101	N1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N2-methyl-N2-(2-phenylethyl)glycinamide	
102	1-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-4-(phenylmethyl)imidazolidin-2-one	
103	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}pyridazin-3-yl)-N'-(4-fluorophenyl)propanediamide	

Table 1

Entry	Name	Structure
104	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(2-chlorophenyl)propanediamide	
105	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(3-chlorophenyl)propanediamide	
106	N1-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N2-methyl-N2-(phenylmethyl)glycinamide	
107	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-chlorophenyl)propanediamide	
108	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[(methyloxy)imino]propanamide	
109	(2E)-N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-2-[(ethyloxy)imino]propanamide	

Table 1

Entry	Name	Structure
110	(2E)-N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-2-{{[(phenylmethyl)oxy]imino}propanamide	
111	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-1-(phenylmethyl)prolinamide	
112	1-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-3-[(4-methylphenyl)sulfonyl]-4-(phenylmethyl)imidazolidin-2-one	
113	1-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-4-(phenylmethyl)imidazolidin-2-one	
114	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-4-(phenylmethyl)-4,5-dihydro-1,3-oxazol-2-amine	
115	6,7-bis(methyloxy)-4-({4-[4-(phenylmethyl)piperazin-1-yl]phenyl}oxy)quinoline	

Table 1

Entry	Name	Structure
116	1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-4-(phenylmethyl)piperazin-2-one	
117	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N2-(phenylmethyl)alaninamide	
118	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N2-methyl-N2-(phenylmethyl)alaninamide	
119	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N2-(phenylmethyl)leucinamide	
120	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N2-methyl-N2-(phenylmethyl)leucinamide	
121	N1-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N2-(phenylmethyl)valinamide	

Table 1

Entry	Name	Structure
122	4-(6,7-dimethoxy-quinolin-4-ylamino)-N-(3-phenyl-propyl)-benzamide	
123	4-benzyl-1-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenyl]-tetrahydro-pyrimidin-2-one	
124	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
125	2-(Benzyl-methyl-amino)-N-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenyl]-3-methyl-butyramide (note: Alphabetic order of prefixes ignored while selecting parent chain)	
126	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-phenyl]-2-phenoxylimino-propionamide	

Table 1





Entry	Name	Structure
127	2-Benzyloxyimino-N-[4-(6,7-dimethoxy-quinolin-4-yloxy)-phenyl]-2-phenyl-acetamide	
128	4-[4-(4-Benzyl-piperidin-1-yl)-phenoxy]-6,7-dimethoxy-quinoline	
129	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-3-fluorophenyl]-N'-(2-isopropyl-1,2,3,4-tetrahydro-isoquinolin-1-ylmethyl)-oxalamide	
130	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-3-fluorophenyl]-N'-(2-ethyl-1,2,3,4-tetrahydro-isoquinolin-1-ylmethyl)-oxalamide	

Table 1

Entry	Name	Structure
131	4-(4-{3-Chloro-5-[2-(4-fluoro-phenylcarbamoyl)-acetylamino]-pyridin-2-yloxy}-6-methoxy-quinolin-7-yloxymethyl)-piperidine-1-carboxylic acid tert-butyl ester	
132	N-{5-Chloro-6-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	
133	N-{5-Chloro-6-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	
134	N-{4-[7-(3-Diethylamino-propoxy)-6-methoxy-quinolin-4-yloxy]-3-fluoro-phenyl}-N'-phenethyl-oxalamide	

Table 1





Entry	Name	Structure
135	N-{3-Fluoro-4-[6-methoxy-7-(3-morpholin-4-yl-propoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
136	N-{3-Fluoro-4-[6-methoxy-7-(3-piperidin-1-yl-propoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
137	N-{4-[7-(2-Diethylamino-ethoxy)-6-methoxy-quinolin-4-yloxy]-3-fluoro-phenyl}-N'-phenethyl-oxalamide	
138	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-methyl-N'-phenethyl-oxalamide	

Table 1

Entry	Name	Structure
139	N-{3-Fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
140	N-{3-Fluoro-4-[6-methoxy-7-(2-methyl-octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
141	2-(3,4-Dihydro-1H-isoquinolin-2-yl)-N-{3-fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-acetamide	
142	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-2-(3-phenyl-pyrrolidin-1-yl)-acetamide	

Table 1





Entry	Name	Structure
143	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-2-(2-phenyl-morpholin-4-yl)-acetamide	
144	N-(2-Dimethylamino-2-phenyl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
145	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-oxo-2-phenyl-ethyl)-oxalamide	
146	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-2,2-difluoro-N'-(4-fluoro-phenyl)-malonamide	

Table 1

Entry	Name	Structure
147	N-Benzyl-N'-{3-fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
148	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(2-fluoro-phenyl)-ethyl]-oxalamide	
149	N-[2-(3-Chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
150	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(2-methoxy-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
151	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-pyridin-3-yl-ethyl)-oxalamide	
152	N-Benzyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
153	N-[2-(2,5-Dimethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
154	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(2-trifluoromethyl-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
155	N-[2-(2-Ethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
156	N-[2-(2,4-Dimethyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
157	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1S-phenyl-2-p-tolyl-ethyl)-oxalamide	
158	N-[2-(4-Chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
159	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamic acid	
160	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(3-fluoro-phenyl)-ethyl]-oxalamide	
161	N-[2-(2-Chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
162	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(3-methoxy-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
163	N-(1,2-Diphenyl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
164	N-[2-(2,4-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
165	N-[2-(3,4-Dimethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
166	N-[2-(4-Ethyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
167	N-[2-(4-Ethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
168	N-[2-(4-Ethoxy-3-methoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
169	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-phenoxy-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
170	N-[2-(3-Ethoxy-4-methoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
171	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-pyridin-2-yl-ethyl)-oxalamide	
172	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-pyridin-4-yl-ethyl)-oxalamide	
173	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-fluoro-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
174	N-[2-(2-Bromo-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
175	N-[2-(2-Chloro-6-fluoro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
176	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2 <i>R</i> -phenyl-propyl)-oxalamide	
177	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-indan-1-yl-oxalamide	

Table 1

Entry	Name	Structure
178	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-isobutyl-oxalamide	
179	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methyl-butyl)-oxalamide	
180	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2 <i>R</i> -phenyl-propyl)-oxalamide	
181	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-phenyl-propyl)-oxalamide	

Table 1

Entry	Name	Structure
182	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-indan-2-yl-oxalamide	
183	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R-phenyl-ethyl)-oxalamide	
184	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1S-phenyl-ethyl)-oxalamide	
185	N-[2-(3-Bromo-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
186	N-[2-(2,6-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
187	N-[2-(2,4-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
188	N-(2-Benzo[1,3]dioxol-5-yl-ethyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
189	N-[2-(3-Bromo-4-methoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
190	N-[2-(3,5-Dimethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
191	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-o-tolyl-ethyl)-oxalamide	
192	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-m-tolyl-ethyl)-oxalamide	

Table 1

Entry	Name	Structure
193	N-[2-(3-Ethoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
194	N-[2-(3,4-Dimethyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
195	N-[2-(2,5-Dimethyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
196	N-[2-(3-Chloro-4-propoxy-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
197	N-[2-(4-Butoxy-3-chloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
198	N-[2-(4-tert-Butyl-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
199	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-sulfamoyl-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
200	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-hydroxy-3-methoxy-phenyl)-ethyl]-oxalamide	
201	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(3-hydroxy-4-methoxy-phenyl)-ethyl]-oxalamide	
202	N-(2,4-Dichloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
203	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-fluoro-2-trifluoromethyl-benzyl)-oxalamide	
204	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1-p-tolyl-ethyl)-oxalamide	
205	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-fluoro-4-trifluoromethyl-benzyl)-oxalamide	

Table 1




Entry	Name	Structure
206	N-(3-Chloro-4-fluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
207	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[1-(3-methoxy-phenyl)-ethyl]-oxalamide	
208	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1-naphthalen-2-yl-ethyl)-oxalamide	

Table 1




Entry	Name	Structure
209	N-(4-Chloro-3-trifluoromethyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
210	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1-p-tolyl-ethyl)-oxalamide	
211	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(6-trifluoromethyl-pyridin-3-ylmethyl)-oxalamide	

Table 1





Entry	Name	Structure
212	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-methyl-benzyl)-oxalamide	
213	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methyl-benzyl)-oxalamide	
214	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-fluoro-3-trifluoromethyl-benzyl)-oxalamide	
215	N-(3,5-Dichloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1





Entry	Name	Structure
216	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R,2,3,4-tetrahydronaphthalen-1-yl)-oxalamide	
217	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1S,2,3,4-tetrahydronaphthalen-1-yl)-oxalamide	
218	N-Cyclopentyl-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	
219	N-[1-(4-Bromo-phenyl)-ethyl]-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	

Table 1

Entry	Name	Structure
220	N-(2-Fluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
221	N-[2-(3,4-Dichloro-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
222	N-(4-Fluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
223	N-(2,3-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
224	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-phenoxy-ethyl)-oxalamide	
225	N-(2,2-Diphenyl-ethyl)-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	
226	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-methoxy-phenyl)-ethyl]-oxalamide	
227	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-phenyl-propyl)-oxalamide	

Table 1





Entry	Name	Structure
228	N-[2-(4-Bromo-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
229	N-{4-[7-(1-Ethyl-piperidin-4-ylmethoxy)-6-methoxy-quinolin-4-yloxy]-3-fluorophenyl}-2-oxo-2-(2-phenyl-morpholin-4-yl)-acetamide	
230	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-fluoro-5-trifluoromethylbenzyl)-oxalamide	
231	N-(3,5-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
232	N-(2-Chloro-5-trifluoromethyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
233	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-3-fluoro-phenyl]-N'-(2-dimethylamino-2-phenyl-ethyl)-oxalamide	
234	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-methoxy-benzyl)-oxalamide	

Table 1





Entry	Name	Structure
235	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-trifluoromethyl-benzyl)-oxalamide	
236	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methoxy-benzyl)-oxalamide	
237	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-trifluoromethyl-benzyl)-oxalamide	
238	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-trifluoromethoxy-benzyl)-oxalamide	

Table 1





Entry	Name	Structure
239	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-methoxy-benzyl)-oxalamide	
240	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-trifluoromethyl-benzyl)-oxalamide	
241	N-(3-Chloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
242	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-trifluoromethoxy-benzyl)-oxalamide	

Table 1

Entry	Name	Structure
243	N-(2-Chloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
244	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-trifluoromethoxy-benzyl)-oxalamide	
245	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-methoxy-benzyl)-oxalamide	

Table 1

Entry	Name	Structure
246	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(4-trifluoromethyl-benzyl)-oxalamide	
247	N-{4-[7-(Azetidin-3-ylmethoxy)-6-methoxy-quinolin-4-yloxy]-3-fluorophenyl}-N'-phenethyl-oxalamide	
248	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-azetidin-3-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
249	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-hydroxy-2-phenyl-ethyl)-oxalamide	

Table 1

Entry	Name	Structure
250	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(2,4-difluorophenyl)-malonamide	
251	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(4-fluorophenyl)-N'-methyl-malonamide	
252	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R-phenyl-propyl)-oxalamide	

Table 1





Entry	Name	Structure
253	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(1R-phenyl-propyl)-oxalamide	
254	N-(3,4-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
255	N-(2,6-Difluoro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
256	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[2-(4-fluoro-phenyl)-ethyl]-oxalamide	

Table 1

Entry	Name	Structure
257	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-phenyl-oxalamide	
258	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-fluoro-phenyl)-oxalamide	
259	N-(4-Chloro-3-fluoro-phenyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
260	N-(3,4-Dimethoxy-phenyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
261	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(3-methyl-butyl)-oxalamide	
262	N-(3,3-Dimethyl-butyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
263	N-{5-Chloro-6-[6-methoxy-7-(3-piperidin-1-yl-propoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	
264	N-{5-Chloro-6-[6-methoxy-7-(3-morpholin-4-yl-propoxy)-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluoro-phenyl)-malonamide	

Table 1




Entry	Name	Structure
265	N-{5-Chloro-6-[7-(3-diethylamino-propoxy)-6-methoxy-quinolin-4-yloxy]-pyridin-3-yl}-N'-(4-fluorophenyl)-malonamide	
266	N-(4-Chloro-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
267	N-(3,5-Dimethoxy-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1




Entry	Name	Structure
268	N-(4-Butyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
269	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-p-tolyl-ethyl)-oxalamide	
270	N-(3,5-Bis-trifluoromethyl-benzyl)-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1





Entry	Name	Structure
271	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-pyrazin-2-ylmethyl-oxalamide	
272	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-pyridin-2-ylmethyl-oxalamide	
273	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	
274	N-{3-Fluoro-4-[6-methoxy-7-(1-methyl-piperidin-4-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	

Table 1

Entry	Name	Structure
275	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-fluoro-3-trifluoromethylbenzyl)-oxalamide	
276	N-[2-(2-Bromo-6-methoxyphenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
277	N-[2-(3,4-Dimethoxyphenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N-methyl-oxalamide	
278	N-[2-(5-Bromo-2-methoxyphenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1

Entry	Name	Structure
279	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-fluoro-5-trifluoromethyl-benzyl)-oxalamide	
280	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-[1-(4-fluorophenyl)-ethyl]-oxalamide	
281	N-(1S-Benzyl-2-oxo-2-pyrrolidin-1-yl-ethyl)-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-oxalamide	
282	N-{3-Fluoro-4-[6-methoxy-7-(octahydro-cyclopenta[c]pyrrol-5-ylmethoxy)-quinazolin-4-yloxy]-phenyl}-N'-phenethyl-oxalamide	

Table 1




Entry	Name	Structure
283	N-[2-(4-Amino-phenyl)-ethyl]-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
284	2-(4-Benzyl-piperidin-1-yl)-N-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-acetamide	
285	N-[4-(6,7-Dimethoxy-quinolin-4-yloxy)-phenyl]-N'-(4-fluoro-phenyl)-malonamide	

Table 1





Entry	Name	Structure
286	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(3-fluorophenyl)-malonamide	
287	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-phenyl-malonamide	
288	N-[5-Chloro-6-(6,7-dimethoxy-quinolin-4-yloxy)-pyridin-3-yl]-N'-(4-fluorophenyl)-2,2-dimethyl-malonamide	
289	N-Ethyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	

Table 1


Entry	Name	Structure
290	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-isopropyl-oxalamide	
291	N-Butyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
292	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-methoxy-ethyl)-oxalamide	
293	N-Cyclopropylmethyl-N'-{3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-oxalamide	
294	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-N'-(2-morpholin-4-yl-ethyl)-oxalamide	

Table 1


Entry	Name	Structure
295	N-{3-Fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl}-2-oxo-2-pyrrolidin-1-yl-acetamide	
296	N-Ethyl-N'-(3-fluoro-4-[6-methoxy-7-(piperidin-4-ylmethoxy)-quinolin-4-yloxy]-phenyl)-N-methyl-oxalamide	

46. A compound for modulating kinase activity of formula A-B-C, or a pharmaceutically acceptable salt, hydrate, or prodrug thereof, wherein, A is selected from:


$-R^3$		




B is selected from:



and, C is selected from:





wherein R^2 is selected from -H, halogen, trihalomethyl, -CN, -NH₂, -NO₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl;

q is 0 to 2;

each R^3 is independently selected from -H, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted arylalkyl, and optionally substituted heteroarylalkyl;

two R^3 , together with the nitrogen to which they are attached, form a four- to seven-membered heteroalicyclic, said four- to seven-membered heteroalicyclic optionally containing one additional heteroatom; when one said additional heteroatom is a nitrogen, then said nitrogen is optionally substituted with a group selected from -H, trihalomethyl, -SO₂R⁵, -SO₂NR⁵R⁵, -CO₂R⁵, -C(O)NR⁵R⁵, -C(O)R⁵, and optionally substituted lower alkyl;

each R^{35} is independently selected from -H, -C(=O)R³, -C(=O)OR³, -C(=O)SR³, -SO₂R³, -C(=O)N(R³)R³, and optionally substituted lower alkyl;

two R^{35} , together with the nitrogen to which they are attached, can combine to form a heteroalicyclic optionally substituted with between one and four of R^{60} , said heteroalicyclic may have an additional annular heteroatom, and said heteroalicyclic may have an aryl fused thereto, said aryl optionally substituted with an additional one to four of R^{60} ;

A^1 is selected from $=N-$, $=C(H)-$, and $=C(CN)-$;

A^2 is either $=N-$ or $=C(H)-$;

R^5 is $-H$ or optionally substituted lower alkyl;

R^8 is selected from R^3 , $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-SO_2R^3$, and $-C(O)R^3$;

R^9 , R^{10} , and R^{11} are each independently selected from $-H$, and $-OR^{12}$; or

R^9 is selected from $-H$, and $-OR^{12}$, and R^{10} and R^{11} , when taken together, are either an optionally substituted alkylidene or an oxo; and

R^{12} is selected from $-H$, $-C(O)R^3$, optionally substituted lower alkylidyne, optionally substituted lower arylalkylidyne, optionally substituted lower heterocyclalkylidyne, optionally substituted lower alkylidene, optionally substituted lower alkylidenearyl, optionally substituted lower alkylideneheterocycl, optionally substituted lower alkyl, optionally substituted lower alkylaryl, optionally substituted aryl, optionally substituted lower heterocyclalkyl, and optionally substituted heterocycl;

or two R^{12} 's, when taken together, form 1) a corresponding spirocyclic ketal when said two R^{12} 's stem from R^{10} and R^{11} , or 2) a corresponding cyclic ketal when said two R^{12} 's stem from R^9 and one of R^{10} and R^{11} ;

E^1 is selected from $-O-$, $-CH_2-$, $-N(R^5)-$, and $-S(O)_{0-2}-$;

Q is a five- to ten-membered ring system, optionally substituted with between zero and four of R^{20} ;

R^{20} is selected from $-H$, halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted lower alkyl;

R^{60} is selected from $-H$, halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$,



$-N(R^3)CO_2R^3$, $-C(O)R^3$, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted heteroarylalkyl, and optionally substituted arylalkyl;

two of R^{60} , when attached to a non-aromatic carbon, can be oxo;

each methylene in any of the above formulae is independently optionally substituted with R^{25} ;


each R^{25} is independently selected from halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^3$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, optionally substituted aryl, optionally substituted arylalkyl, heteroarylalkyl, and optionally substituted lower alkyl; two of R^{25} , together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic, two of R^{25} on a single carbon can be oxo;


with the proviso that when B is selected from:














and C contains

$(R^2)_q$, and the remaining portion of C contains one of:










 directly attached to $(R^2)_q$, then A must be one of:


and with the proviso that when C contains




 $(R^2)_q$, and B is selected from:

then the portion of C directly attached to





cannot contain



, when R^{70} is selected from -H, C_{1-4} alkyl, and C_{1-4} alkoxyl.


47. The compound according to claim 46, wherein Q is selected from phenyl, naphthyl, 1,2,3,4-tetrahydronaphthyl, indanyl, benzodioxanyl, benzofuranyl, phenazinyl, phenothiazinyl, phenoxazinyl, tetrahydroisoquinolyl, pyrrolyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolyl, imidazolidinyl, tetrahydropyridinyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolyl, oxazolidinyl, triazolyl, isoxazolyl, isoxazolidinyl, thiazolyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, isoindolyl, indolyl, isoindolyl, octahydroindolyl, octahydroisoindolyl, quinolyl, isoquinolyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, thienyl, benzothioliyl, and oxadiazolyl; each optionally substituted with between one and four of R^{20} ; wherein each R^{20} is independently selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl.

48. The compound according to claim 47, wherein B is either of the following:





wherein A^1 is either =N- or =C(H)-.

49. The compound according to claim 48, wherein B is




50. The compound according to claim 49, wherein C is selected from:





wherein R^2 , R^3 , R^5 , R^{20} , R^{25} and R^{60} are as defined above.

51. The compound according to claim 50, R^2 is selected from halogen, trihalomethyl, -CN, -NO₂, -OR³, -NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted lower alkyl
52. The compound according to claim 51, wherein R^2 is halogen.
53. The compound according to claim 52, wherein R^2 is either fluorine or chlorine.
54. A compound for modulating kinase activity according to Formula XI,



XI

or a pharmaceutically acceptable salt, hydrate, or prodrug thereof, wherein,

each R^1 is independently selected from halogen, -OR³, -NO₂, -NH₂, -NR³R⁴, -D-R⁵⁰ and optionally substituted C₁₋₆alkyl;

R^{70} is selected from -H, halogen, -OR³, -S(O)₀₋₂R³, -NO₂, -NH₂, -NR³R⁴, and optionally substituted C₁₋₆alkyl;

Q is selected from =N-, =C(H)-, and =C(CN)-;

Z is selected from $-S(O)_{0-2}-$, $-O-$, and $-NR^5-$;

Ar is either a five- or six-membered arylene or a five- or six-membered heteroarylene containing between one and three heteroatoms;

G is either an optionally substituted cycloalkyl or an optionally substituted heteroalicyclic;

each R^2 is independently selected from halogen, trihalomethyl, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted C_{1-6} alkyl;

each R^3 is independently $-H$ or R^4 ;


each R^4 is independently selected from optionally substituted C_{1-6} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C_{1-6} alkyl; or

R^3 and R^4 , when taken together with a common nitrogen to which they are attached, form an optionally substituted five- to seven-membered heterocyclyl, said optionally substituted five- to seven-membered heterocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

R^5 is $-H$ or optionally substituted C_{1-6} alkyl;

each D is independently selected from $-O-$, $-S(O)_{0-2}-$, and $-NR^5-$;

each R^{50} is independently either R^3 , or according to formula XII;



wherein X^1 , X^2 , and optionally X^3 , represent the atoms of a saturated bridged ring system, said saturated bridged ring system comprising up to four annular heteroatoms represented by any of X^1 , X^2 , and X^3 ; wherein,

each X^1 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

each X^2 is independently an optionally substituted bridgehead methine or a bridgehead nitrogen;

each X^3 is independently selected from $-C(R^6)R^7-$, $-O-$, $-S(O)_{0-2}-$, and $-NR^8-$;

Y is either:

an optionally substituted lower alkylene linker, between D and either 1) any annular atom of the saturated bridged ring system, except X^2 when X^2 is a bridgehead nitrogen, or 2) any heteroatom, represented by any of R^6 or R^7 ; provided there are at least two carbon atoms between D and any annular heteroatom of the saturated bridged ring system or any heteroatom represented by any of R^6 or R^7 ;

or Y is absent, when Y is absent, said saturated bridged ring system, is directly attached to D via an annular carbon of said saturated bridged ring system, unless D is $-SO_2-$, in which case said saturated bridged ring system, is directly attached to D via an any annular atom of said saturated bridged ring system;

m and p are each independently one to four;

n is zero to two, when n equals zero, then there is a single bond between the two bridgehead X^2 's;

R^6 and R^7 are each independently selected from -H, halogen, trihalomethyl, -CN, $-NH_2$, $-NO_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^4$, $-SO_2NR^3R^4$, $-CO_2R^3$, $-C(O)NR^3R^4$, $-N(R^3)SO_2R^4$, $-N(R^3)C(O)R^3$, $-NCO_2R^3$, $-C(O)R^3$, optionally substituted C_{1-6} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-6} alkyl, optionally substituted heterocyclyl, optionally substituted heterocyclyl C_{1-6} alkyl, and a bond to either Y or D; or

R^6 and R^7 , when taken together are oxo; or

R^6 and R^7 , when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;


R^8 is selected from $-R^3$, Y, $-SO_2NR^3R^4$, $-CO_2R^4$, $-C(O)NR^3R^3$, $-SO_2R^4$, and $-C(O)R^3$; and each R^{30} is independently selected from halogen, trihalomethyl, -CN, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-S(O)_{0-2}R^3$, $-SO_2NR^3R^3$, $-CO_2R^3$, $-C(O)NR^3R^3$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, $-C(O)R^3$, and optionally substituted C_{1-6} alkyl.

55. The compound according to claim 54, wherein Z is either -O- or $-NR^5-$.

56. The compound according to claim 55, wherein at least one of R^1 is $-D-R^{50}$.

57. The compound according to claim 56, wherein D is -O- and at least one other R^1 is -OR³.

58. The compound according to claim 57, of formula **XIIIa** or **XIIIb**:



wherein Q^1 is either =N- or =C(H)-.

59. The compound according to claim 58, wherein R^{50} is selected from C_{1-6} alkyl optionally substituted with at least one of optionally substituted amino, optionally substituted C_{1-6} alkyl amino, optionally substituted C_{1-6} dialkyl amino, optionally substituted heteroalicyclic, and a group of formula **XII**.

60. The compound according to claim 59, wherein R^{3a} is C_{1-6} alkyl.


61. The compound according to claim 60, wherein Z is -O-.

62. The compound according to claim 61, wherein G is selected from cyclopropyl, aziradine, cyclobutyl, and azetidine, each optionally substituted with between zero and four of R^{30} .

63. The compound according to claim 62, wherein Q is either =N- or =C(H)-.

64. The compound according to claim 63, wherein R^2 is selected from -H, halogen, C_{1-6} alkyl and perfluoro C_{1-6} alkyl.

65. The compound according to claim 64, wherein $-N(R^{3b})R^4$ is selected from the following:



wherein J, is a five- to ten-membered ring, optionally substituted with between zero and five of R^{20} ;

each R^{20} is independently selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted C₁₋₆alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₆alkyl, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C₁₋₆alkyl;

two of R^{20} , together with the atom or atoms to which they are attached, combine to form an optionally substituted three- to seven-membered heteroalicyclic, said optionally substituted three- to seven-membered heteroalicyclic either spiro- to J or fused to J;

E is selected from -O-, -N(R⁵)-, -CH₂-, and -S(O)₀₋₂-;

each R^{60} is independently selected from halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted C₁₋₆alkyl, optionally substituted aryl, optionally substituted heteroaryl C₁₋₆alkyl, and optionally substituted aryl C₁₋₆alkyl;

each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R^{25} ; and


R^{25} is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R⁴, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted aryl C₁₋₆alkyl, heteroaryl C₁₋₆alkyl, and optionally substituted C₁₋₆alkyl; or

two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered alicyclic or heteroalicyclic;

R^{3b} is equivalent to R^3 as defined above; and

R^4 and R^5 are as defined above.

66. The compound according to claim 65, of formula **XIVa** or **XIVb**:



67. The compound according to claim 66, wherein R⁵⁰ is C₁₋₆alkyl optionally substituted with a group selected from optionally substituted amino, an optionally substituted alkylamino, optionally substituted dialkylamino, and optionally substituted heteroalicyclic.

68. The compound according to claim 67, wherein the heteroalicyclic portion of said optionally substituted heteroalicyclic of R⁵⁰ is selected from the group consisting of piperidine, piperazine, morpholine, thiomorpholine, thiomorpholine 1-oxide, thiomorpholine 1,1-dioxide, 2-oxo-morpholine, pyrrolidine, and azepine.

69. The compound according to claim 67, wherein R⁵⁰ is according to formula XII.

70. The compound according to claim 69, wherein the saturated bridged ring system according to formula XII has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], [3.1.0], [3.3.3], [3.3.2], [3.3.1], [3.2.2], [3.2.1], [2.2.2], and [2.2.1].


71. The compound according to claim 70, wherein Y is selected from -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂-, -CH₂-, and absent.

72. The compound according to claim 71, wherein n is 0 and the saturated bridged ring system according to formula XII has a geometry selected from the group consisting of [4.4.0], [4.3.0], [4.2.0], [4.1.0], [3.3.0], [3.2.0], and [3.1.0].

73. The compound according to claim 72, wherein said saturated bridged ring system contains at least one annular nitrogen or at least one annular oxygen.

74. The compound according to claim 73, wherein said saturated bridged ring system contains $-NR^8$ -, wherein R^8 is selected from $-H$, optionally substituted C_{1-6} alkyl, $-CO_2R^3$, $-C(O)NR^3R^3$, $-SO_2R^3$, and $-C(O)R^3$.

75. The compound according to claim 73, wherein said saturated bridged ring system is of formula XV,



XV

wherein U^1 is selected from $-O-$, $-S(O)_{0-2}-$, $-NR^8$ -, $-CR^6R^7$ -, and absent; and e is 0 or 1.

76. The compound according to claim 75, wherein Y is $-CH_2-$.


77. The compound according to claim 76, wherein U^1 is $-NR^8$ -, wherein R^8 is selected from $-H$, optionally substituted lower alkyl, $-CO_2R^3$, $-C(O)NR^3R^3$, $-SO_2R^3$, and $-C(O)R^3$.

78. The compound according to claim 76, wherein U^1 is $-O-$.

79. The compound according to claim 76, wherein U^1 is absent.

80. The compound according to claim 71, wherein Y is selected from $-CH_2CH_2-$, $-CH_2-$, and absent.

81. The compound according to claim 80, wherein said saturated bridged ring system is of formula XVI,



XVI

wherein R^9 , R^{10} , and R^{11} are each independently selected from -H, and $-OR^{12}$; or

R^9 is selected from -H, and $-OR^{12}$, and R^{10} and R^{11} , when taken together, are either an optionally substituted alkylidene or an oxo;

R^{12} is selected from -H, $-C(O)R^3$, optionally substituted lower alkylidene, optionally substituted lower arylalkylidene, optionally substituted lower heterocyclalkylidene, optionally substituted lower alkylidene, optionally substituted lower alkylidenearyl, optionally substituted lower alkylideneheterocycl, optionally substituted lower alkyl, optionally substituted lower alkylaryl, optionally substituted aryl, optionally substituted lower heterocyclalkyl, and optionally substituted heterocycl;


or two R^{12} 's, when taken together, form 1) a corresponding spirocyclic ketal when said two R^{12} 's stem from R^{10} and R^{11} , or 2) a corresponding cyclic ketal when said two R^{12} 's stem from R^9 and one of R^{10} and R^{11} .

82. The compound according to claim 81, wherein one of R^{10} and R^{11} is $-OR^{12}$, wherein R^{12} is selected from -H, $-C(O)R^3$, and optionally substituted lower alkyl; and R^9 and the other of R^{10} and R^{11} are both -H.

83. The compound according to claim 82, wherein Y is either $-CH_2-$ or absent.

84. The compound according to claim 81, wherein R^9 is an alkyl group containing at least one fluorine substitution thereon.

85. The compound according to claim 74, wherein said saturated bridged ring system is of formula **XVII**.




XVII

86. The compound according to claim 85, wherein Y is either $-CH_2-$ or absent.

87. The compound according to claim 86, wherein R^8 is methyl or ethyl.
88. The compound according to claim 87, wherein at least one of R^2 is halogen.
89. The compound according to claim 74, wherein said saturated bridged ring system is of formula **XVIII**.

**XVIII**

90. The compound according to claim 89, wherein Y is $-\text{CH}_2-$.
91. The compound according to claim 90, wherein R^8 is methyl or ethyl.
92. The compound according to claim 73, wherein said saturated bridged ring system is of formula **XIX**.

**XIX**

wherein U^2 is selected from $-\text{O}-$, $-\text{S}(\text{O})_{0-2}-$, $-\text{NR}^8-$, $-\text{CR}^6\text{R}^7-$, and absent.

93. The compound according to claim 92, wherein R^3 of formula **XIX** is selected from $-\text{H}$ and optionally substituted alkyl.
94. The compound according to claim 93, wherein U^2 is either $-\text{CR}^6\text{R}^7-$ or absent.
95. The compound according to claim 94, wherein U^2 is either $-\text{CH}_2-$ or absent.
96. The compound according to claim 95, wherein Y is $-\text{CH}_2-$.
97. The compound according to claim 74, wherein said saturated bridged ring system is according to formula **XX**.



XX

98. The compound according to claim 97, wherein R^8 is methyl or ethyl.
99. The compound according to any of claims 67 through 98, wherein R^2 is selected from C_{1-6} alkyl, perfluoro C_{1-6} alkyl, and halogen.
100. The compound according to claim 99, wherein R^2 is selected from perfluoro C_{1-3} alkyl and halogen.
101. The compound according to any of claims 67 through 98, wherein R^{20} is selected from halogen, $-CN$, $-NO_2$, $-NH_2$, $-OR^3$, $-NR^3R^4$, $-N(R^3)SO_2R^3$, $-N(R^3)C(O)R^3$, $-N(R^3)CO_2R^3$, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C_{1-6} alkyl, and (two of R^{20}) together with the atom or atoms to which they are attached, an optionally substituted three- to six-membered heteroalicyclic, said optionally substituted three- to six-membered heteroalicyclic fused to the phenyl as in **XIVa** or **XIVb**.
102. The compound according to claim 101, wherein R^{20} is selected from halogen, $-NR^3R^4$, optionally substituted heterocyclyl, and optionally substituted heterocyclyl C_{1-6} alkyl, and (two of R^{20}) together with the atom or atoms to which they are attached, an optionally substituted five- to six-membered heteroalicyclic, said optionally substituted five- to six-membered heteroalicyclic fused to the phenyl as in **XIVa** or **XIVb**.
103. The compound according to claim 102, wherein R^2 is selected from C_{1-6} alkyl, perfluoro C_{1-6} alkyl, and halogen.
104. The compound according to claim 103, wherein R^2 is selected from perfluoro C_{1-3} alkyl and halogen.
105. The compound according to claim 54, selected from Table 2.

Table 2

Entry	Name	Structure
1	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
2	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
3	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-(phenylmethyl)cyclopropane-1,1-dicarboxamide	
4	N-(6-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloropyridin-3-yl)-N'-phenylcyclopropane-1,1-dicarboxamide	
5	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
6	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperidin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
7	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperidin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
8	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-5-chloropyridin-3-yl)-N'-(2-phenylethyl)cyclopropane-1,1-dicarboxamide	
9	N-(6-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}-2-methylpyridin-3-yl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
10	N-{4-[(7-chloroquinolin-4-yl)oxy]-3-fluorophenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
11	N-{4-[(7-chloroquinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
12	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
13	N-(4-{[6,7-bis(methyloxy)quinazolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
14	N-(4-{[6,7-bis(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
15	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
16	N-{5-chloro-6-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy]quinolin-4-yl]oxy}pyridin-3-yl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
17	N-[5-chloro-6-[(6-(methyloxy)-7-[(piperidin-4-ylmethyl)oxy]quinolin-4-yl]oxy]pyridin-3-yl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
18	N-[5-chloro-6-[(6-(methyloxy)-7-[(phenylmethyl)oxy]quinolin-4-yl]oxy]pyridin-3-yl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
19	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
20	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
21	N-{3-fluoro-4-[(6-(methyloxy)-7-{[(1-methylpiperidin-4-yl)methyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
22	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
23	N-(4-fluorophenyl)-N'-[2-methyl-6-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)pyridin-3-yl]cyclopropane-1,1-dicarboxamide	
24	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
25	N-(6-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-5-chloro-2-methylpyridin-3-yl})-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
26	N-[3-fluoro-4-({7-(methyloxy)-6-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
27	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-3,5-difluorophenyl})-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
28	N-(4-{{[6,7-bis(methyloxy)quinolin-4-yl]oxy}-2,5-difluorophenyl})-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
29	N-[3-fluoro-4-({7-(methyloxy)-6-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
30	N-{3-fluoro-4-[(6-(methyloxy)-7-(2-methyloctahydrocyclopenta[c]pyrrol-5-ylmethoxy)quinazolin-4-yl]oxy}phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
31	N-{3-fluoro-4-[(7-(methyloxy)-6-[[1-methylpiperidin-4-yl)methyl]oxy}quinazolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
32	N-[5-fluoro-2-methyl-4-((6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl)oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
33	N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-2,3,5-trifluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
34	N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-5-fluoro-2-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
35	N-(4-[[6,7-bis(methyloxy)quinolin-4-yl]oxy]-2-chloro-5-methylphenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
36	N-(3-fluoro-4-[[6-hydroxy-7-(methyloxy)quinolin-4-yl]oxy]phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
37	N-(4-fluorophenyl)-N'-[2-methyl-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]cyclopropane-1,1-dicarboxamide	
38	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
39	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
40	N-{3-fluoro-4-[(6-(methyloxy)-7-[(1-methylpiperidin-4-yl)methyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
41	N-(4-fluorophenyl)-N'-[4- ({6-(methyloxy)-7-[(3- morpholin-4- ylpropyl)oxy]quinolin-4- yl}oxy)phenyl]cyclopropan- e-1,1-dicarboxamide	
42	N-(4-{[7-{[3- (diethylamino)propyl]oxy }-6-(methyloxy)quinolin- 4-yl]oxy}-3- fluorophenyl)-N'-(4- fluorophenyl)cyclopropan- e-1,1-dicarboxamide	
43	N-(4-{[6,7- bis(methyloxy)quinolin-4- yl]oxy}-2-chloro-5- fluorophenyl)-N'-(4- fluorophenyl)cyclopropan- e-1,1-dicarboxamide	
44	N-(4-{[6,7- bis(methyloxy)-2- (methylthio)quinolin-4- yl]oxy}-3-fluorophenyl)- N'-(4- fluorophenyl)cyclopropan- e-1,1-dicarboxamide	
45	N-(4-fluorophenyl)-N'-[4- {[2-methyl-6,7- bis(methyloxy)quinazolin- 4- yl]oxy}phenyl]cyclopropan- e-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
46	N-(4-{[2-amino-6,7-bis(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
47	N-(3-fluoro-4-{[2-(methylamino)-6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
48	(1S,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
49	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
50	N-(4-{[6-{[3-(diethylamino)propyl]oxy}-7-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
51	N-(4-{[6-{[2-(diethylamino)ethyl]oxy}-7-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
52	1,1-dimethylethyl 4-(3-{[4-[(2-fluoro-4-{[(1-{[(4-fluorophenyl)amino]carbonyl}cyclopropyl)carbonyl]amino}phenyl)oxy]-6-(methyloxy)quinolin-7-yl]oxy}propyl)piperazine-1-carboxylate	
53	(1R,2R)-N-[3-fluoro-4-((6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl)oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
54	(1R,2R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
55	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
56	N-(4-{[7-{[3-(4-acetylpiperazin-1-yl)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
57	1,1-dimethylethyl 4-(3-{[4-{[(2-fluoro-4-{[(1R,2R)-1-{[(4-fluorophenyl)amino]carbonyl}-2-methylcyclopropyl)carbonyl]amino}phenyl]oxy}-6-(methyloxy)quinolin-7-yl]oxy}propyl)piperazine-1-carboxylate	

Table 2

Entry	Name	Structure
58	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)-1-(phenylmethyl)azetidine-3,3-dicarboxamide	
59	N-(4-{{6,7-bis(methyloxy)quinolin-4-yl}oxy}phenyl)-N'-(4-fluorophenyl)azetidine-3,3-dicarboxamide	
60	(1R,2S)-N-{3-fluoro-4-[[6-(methyloxy)-7-{{3-(4-methylpiperazin-1-yl)propyl}oxy}quinolin-4-yl}oxy]phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
61	(1R,2R)-N-{3-fluoro-4-[[6-(methyloxy)-7-{{3-(4-methylpiperazin-1-yl)propyl}oxy}quinolin-4-yl}oxy]phenyl}-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
62	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
63	N-(3-fluoro-4-{[7-({3-[4-(1-methylethyl)piperazin-1-yl]propyl}oxy)-6-(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
64	N-(4-{[7-({3-(diethylamino)propyl}oxy)-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide	
65	(1R,2R)-N-(4-{[7-({3-(diethylamino)propyl}oxy)-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
66	(1R,2R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
67	(1R,2S)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
68	(1R,2S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
69	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
70	(1R,2S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
71	(1R,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
72	(1R,2R,3S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
73	(1R,2R,3S)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
74	(1R,2R,3S)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
75	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl)oxy]phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
76	(2R,3R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl)oxy]phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
77	(2R,3R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	

Table 2






Entry	Name	Structure
78	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
79	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
80	(1R,2R,3S)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
81	N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
82	(1R,2R,3S)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
83	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
84	N-(4-{[7-([2-(diethylamino)ethyl]oxy)-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
85	N-(4-{[7-([3-(diethylamino)propyl]oxy)-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,2-dimethylcyclopropane-1,1-dicarboxamide	
86	N-(4-{[7-([3-(diethylamino)propyl]oxy)-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
87	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
88	N-[3-fluoro-4-({6-(methyloxy)-7-[(3-piperazin-1-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
89	(2R,3R)-N-[3-fluoro-4-({6-(methyloxy)-7-[(3-morpholin-4-ylpropyl)oxy]quinazolin-4-yl}oxy)phenyl]-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
90	N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	
91	N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinolin-4-yl)oxy]phenyl}-N'-(4-fluorophenyl)cyclobutane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
92	(1R,2R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
93	(1R,2R)-N-{3-fluoro-4-[(6-(methyloxy)-7-{[3-(4-methylpiperazin-1-yl)propyl]oxy}quinazolin-4-yl]oxy}phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	
94	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
95	(2R,3R)-N-(4-{[7-{[3-(diethylamino)propyl]oxy}-6-(methyloxy)quinazolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
96	(1R,2R)-N-[3-fluoro-4-({6-(methyloxy)-7-[3-(3-piperazin-1-ylpropyl)oxy]quinazolin-4-yl]oxy}phenyl]-N'-(4-fluorophenyl)-2-methylcyclopropane-1,1-dicarboxamide	


Table 2

Entry	Name	Structure
97	(2R,3R)-N-(4-{[7-{[2-(diethylamino)ethyl]oxy}-6-(methyloxy)quinolin-4-yl]oxy}-3-fluorophenyl)-N'-(4-fluorophenyl)-2,3-dimethylcyclopropane-1,1-dicarboxamide	
98	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[(4-fluorophenyl)methyl]cyclopropane-1,1-dicarboxamide	
99	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-(2-morpholin-4-ylethyl)cyclopropane-1,1-dicarboxamide	
100	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	
101	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
102	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	
103	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[2-(morpholin-4-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	
104	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-phenylcyclopropane-1,1-dicarboxamide	
105	N-[3-(aminomethyl)phenyl]-N'-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)cyclopropane-1,1-dicarboxamide	
106	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(piperidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	

Table 2

Entry	Name	Structure
107	N-(4-{[6,7-bis(methyloxy)quinolin-4-yl]oxy}phenyl)-N'-[3-(pyrrolidin-1-ylmethyl)phenyl]cyclopropane-1,1-dicarboxamide	

106. A pharmaceutical composition comprising a compound according to any one of claims 1 - 105 and a pharmaceutically acceptable carrier.

107. A metabolite of the compound or the pharmaceutical composition according to any one of claims 1 - 106.

108. A method of modulating the *in vivo* activity of a kinase, the method comprising administering to a subject an effective amount of the compound or the pharmaceutical composition according to any of claims 1 - 105.

109. The method according to claim 108, wherein modulating the *in vivo* activity of the kinase comprises inhibition of said kinase.

110. The method according to claim 108, wherein the kinase is at least one of c-Met, KDR, c-Kit, flt-3, and flt-4.


111. The method according to claim 110, wherein the kinase is c-Met.

112. A method of treating diseases or disorders associated with uncontrolled, abnormal, and/or unwanted cellular activities, the method comprising administering, to a mammal in need thereof, a therapeutically effective amount of the compound or the pharmaceutical composition as described in any one of claims 1 - 106.



113. A method of screening for a modulator of a kinase, said kinase selected from c-Met, KDR, c-Kit, flt-3, and flt-4, the method comprising combining a compound according to any one of claims 1 - 105, and at least one candidate agent and determining the effect of the candidate agent on the activity of said kinase.

114. A method of inhibiting proliferative activity in a cell, the method comprising administering an effective amount of a composition comprising a compound according any one of claims 1 - 105 to a cell or a plurality of cells.

115. A process for preparing a compound of Formula **XXI**,



comprising reaction of a compound of Formula **XXII**, with a compound of Formula **XXIII**



wherein,

each R¹ is independently selected from halogen, -OR³, -NO₂, -NH₂, -NR³R³, -D-R⁵⁰ and optionally substituted C₁₋₆alkyl;

R⁷⁰ is selected from -H, halogen, -OR³, -S(O)₀₋₂R³, -NO₂, -NH₂, -NR³R³, and optionally substituted C₁₋₆alkyl;

J is selected from =N-, =C(H)-, =C(halogen)-, and =C(CN)-;

Z is selected from -S(O)₀₋₂-, -O-, and -NR⁵-;

each R⁵ is independently selected from -H, optionally substituted C₁₋₆alkyl, optionally substituted aryl, and optionally substituted aryl C₁₋₆alkyl;

Ar is either a five- to ten-membered arylene or a five- to ten-membered heteroarylene containing between one and three heteroatoms;

R² is selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, and optionally substituted C₁₋₆alkyl;

each R^3 is independently selected from -H, $-\text{Si}(\text{R}^5)(\text{R}^5)\text{R}^5$, optionally substituted lower alkyl, optionally substituted aryl, optionally substituted arylalkyl, and optionally substituted heteroarylalkyl;

two R^3 , together with the nitrogen to which they are attached, form a four- to seven-membered heteroalicyclic, said four- to seven-membered heteroalicyclic optionally containing one additional heteroatom; when one said additional heteroatom is a nitrogen, then said nitrogen is optionally substituted with a group selected from -H, trihalomethyl, $-\text{SO}_2\text{R}^5$, $-\text{SO}_2\text{NR}^5\text{R}^5$, $-\text{CO}_2\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^5$, $-\text{C}(\text{O})\text{R}^5$, and optionally substituted lower alkyl;

B is selected from absent, $-\text{N}(\text{R}^{13})-$, $-\text{N}(\text{SO}_2\text{R}^{13})-$, $-\text{O}-$, $-\text{S}(\text{O})_{0-2}-$, and $-\text{C}(=\text{O})-$;

L is selected from absent, $-\text{C}(=\text{S})\text{N}(\text{R}^{13})-$, $-\text{C}(=\text{NR}^{14})\text{N}(\text{R}^{13})-$, $-\text{SO}_2\text{N}(\text{R}^{13})-$, $-\text{SO}_2-$, $-\text{C}(=\text{O})\text{N}(\text{R}^{13})-$, $-\text{N}(\text{R}^{13})-$, $-\text{C}(=\text{O})\text{C}_{1-2}\text{alkylN}(\text{R}^{13})-$, $-\text{N}(\text{R}^{13})\text{C}_{1-2}\text{alkylC}(=\text{O})-$, $-\text{C}(=\text{O})\text{C}_{0-1}\text{alkylC}(=\text{O})\text{N}(\text{R}^{13})-$, $-\text{C}(=\text{O})-$, $-\text{C}_{0-4}\text{alkylene}-$, $-\text{C}(=\text{O})\text{C}_{0-1}\text{alkylC}(=\text{O})\text{OR}^3-$, $-\text{C}(=\text{NR}^{14})\text{C}_{0-1}\text{alkylC}(=\text{O})-$, $-\text{C}(=\text{O})\text{C}_{0-1}\text{alkylC}(=\text{O})-$, and an optionally substituted four- to six-membered heterocyclyl containing between one and three annular heteroatoms and comprising at least one nitrogen;

T is selected from -H, $-\text{R}^{13}$, $-\text{C}_{0-4}\text{alkyl}$, $-\text{C}_{0-4}\text{alkylQ}$, $-\text{OC}_{0-4}\text{alkylQ}$, $-\text{C}_{0-4}\text{alkylOQ}$, $-\text{N}(\text{R}^{13})\text{C}_{0-4}\text{alkylQ}$, $-\text{SO}_2\text{C}_{0-4}\text{alkylQ}$, $-\text{C}(=\text{O})\text{C}_{0-4}\text{alkylQ}$, $-\text{C}_{0-4}\text{alkylN}(\text{R}^{13})\text{Q}$, and $-\text{C}(=\text{O})\text{N}(\text{R}^{13})\text{C}_{0-4}\text{alkylQ}$, wherein each of the aforementioned $\text{C}_{0-4}\text{alkyl}$ is optionally substituted;


Q is a five- to ten-membered ring system, optionally substituted with between zero and four of R^{20} ;

each R^{20} is independently selected from -H, halogen, trihalomethyl, -CN, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OR}^3$, $-\text{NR}^3\text{R}^3$, $-\text{S}(\text{O})_{0-2}\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{N}(\text{R}^3)\text{SO}_2\text{R}^3$, $-\text{N}(\text{R}^3)\text{C}(\text{O})\text{R}^3$, $-\text{N}(\text{R}^3)\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{R}^3$, optionally substituted $\text{C}_{1-6}\text{alkyl}$, optionally substituted aryl, optionally substituted aryl $\text{C}_{1-6}\text{alkyl}$, optionally substituted heterocyclyl, and optionally substituted heterocyclyl $\text{C}_{1-6}\text{alkyl}$;

two of R^{20} , together with the atom or atoms to which they are attached, combine to form an optionally substituted three- to seven-membered heteroalicyclic, said optionally substituted three- to seven-membered heteroalicyclic either spiro- to Q or fused to Q;

D is selected from $-\text{O}-$, $-\text{S}(\text{O})_{0-2}-$, and $-\text{NR}^{15}-$;

R⁵⁰ is either R³, or according to formula **XXIV**;



XXIV

wherein X¹, X², and optionally X³, represent the atoms of a saturated bridged ring system, said saturated bridged ring system comprising up to four annular heteroatoms represented by any of X¹, X², and X³; wherein,

each X¹ is independently selected from -C(R⁶)R⁷-, -O-, -S(O)₀₋₂-, and -NR⁸-;

each X² is independently an optionally substituted bridgehead methine or a bridgehead nitrogen;

each X³ is independently selected from -C(R⁶)R⁷-, -O-, -S(O)₀₋₂-, and -NR⁸-;

Y is either:

an optionally substituted C₁₋₆alkylene linker, between D and either 1) any annular atom of the saturated bridged ring system, except X² when X² is a bridgehead nitrogen, or 2) any heteroatom, represented by any of R⁶ or R⁷; provided there are at least two carbon atoms between D and any annular heteroatom of the saturated bridged ring system or any heteroatom represented by any of R⁶ or R⁷;

or Y is absent, when Y is absent, said saturated bridged ring system, is directly attached to D via an annular carbon of said saturated bridged ring system, unless D is -SO₂-, in which case said saturated bridged ring system, is directly attached to D via an any annular atom of said saturated bridged ring system;

m and p are each independently one to four;

n is zero to two, when n is zero, then there is a single bond between the two bridgehead X²'s;

R⁶ and R⁷ are each independently selected from -H, halogen, trihalomethyl, -CN, -NH₂, -NO₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -NCO₂R³, -C(O)R³, optionally substituted C₁₋₆alkyl, optionally substituted

aryl, optionally substituted aryl C₁₋₆alkyl, optionally substituted heterocyclyl, optionally substituted heterocyclyl a C₁₋₆alkyl, and a bond to either Y or D; or

R⁶ and R⁷, when taken together are oxo; or

R⁶ and R⁷, when taken together with a common carbon to which they are attached, form a optionally substituted three- to seven-membered spirocyclyl, said optionally substituted three- to seven-membered spirocyclyl optionally containing at least one additional annular heteroatom selected from N, O, S, and P;

R⁸ is selected from -R³, Y, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -SO₂R³, and -C(O)R³;

R¹³ is selected from -H, -C(=O)R³, -C(=O)OR³, -C(=O)SR³, -SO₂R³, -C(=O)N(R³)R³, and optionally substituted C₁₋₆alkyl;

two R¹³, together with the atom or atoms to which they are attached, can combine to form a heteroalicyclic optionally substituted with between one and four of R⁶⁰, said heteroalicyclic comprising up to four annular heteroatoms, and said heteroalicyclic optionally comprising an aryl or heteroaryl fused thereto, in which case said aryl or heteroaryl is optionally substituted with an additional one to four of R⁶⁰;

R¹⁴ is selected from -H, -NO₂, -NH₂, -N(R³)R³, -CN, -OR³, optionally substituted C₁₋₆alkyl, optionally substituted heteroalicycyl C₁₋₆alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₆alkyl and optionally substituted heteroalicyclic;

R¹⁵ is a group -M¹-M², wherein M¹ is selected from absent, -C(=S)N(R¹³)-, -C(=NR¹⁴)N(R¹³)-, -SO₂N(R¹³)-, -SO₂-, -C(=O)N(R¹³)-, -C(=O)C(=O)N(R¹³)-, -C₀₋₄alkylene-, -C(=O)-, and an optionally substituted four to six-membered heterocyclyl containing between one and three heteroatoms but comprising at least one nitrogen; and M² is selected from -H, -C₀₋₆alkyl, alkoxy, -C(=O)C₀₋₄alkylQ, -C₀₋₄alkylQ, -OC₀₋₄alkylQ-, -N(R¹³)C₀₋₄alkylQ-, and -C(=O)N(R¹³)C₀₋₄alkylQ;

R⁶⁰ is selected from -H, halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted C₁₋₆alkyl, optionally substituted aryl, optionally substituted heteroaryl C₁₋₆alkyl, and optionally substituted aryl C₁₋₆alkyl;

two of R⁶⁰, when attached to a non-aromatic carbon, can be oxo;


P¹ is a suitable leaving group; and


P^2 is selected from -H, a metal, and a group removed *in-situ* when combining **XXII** and **XXIII** to make **XXI**.


116. The process according to claim 115, wherein Ar is *para*-phenylene as defined by the substitution pattern of -Z- and -B-L-T about said phenylene.

117. The process according to claim 116, wherein Z is either -O- or -NR⁵-.

118. The process according to claim 117, wherein -B-L-T is selected from the following:







wherein Q, R²⁰, and R¹³ are as defined above; each E is selected from -O-, -N(R¹³)-, -CH₂-, and -S(O)₀₋₂-; M is selected from -O-, -N(R¹³)-, -CH₂-, and -C(=O)N(R¹³)-; each V is independently either =N- or =C(H)-; each methylene in any of the above formulae is independently optionally substituted with R²⁵; and R²⁵ is selected from halogen, trihalomethyl, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted aryl C₁₋₆alkyl, heteroaryl C₁₋₆alkyl, and optionally substituted C₁₋₆alkyl; two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form an optionally substituted three- to seven-membered alicyclic or heteroalicyclic; two of R²⁵ on a single carbon can be oxo.

119. The process according to claim 118, wherein there is one of R¹ that is -D-R⁵⁰ and another of R¹ that is -OR^{3a}.


120. The process according to claim 119, wherein D is -O-.


121. The process according to claim 120, wherein -O-R⁵⁰ and -OR^{3a} are interchangeably located at the 6-position and 7-position of the quinazoline or quinoline according to Formula XXI.

122. The process according to claim 121, wherein -OR^{3a} is selected from -OH, -OSi(R⁵)(R⁵)R⁵, and optionally substituted -OC₁₋₆alkyl.

123. The process according to claim 122, wherein J is =N- or =C(H)-.


124. The process according to claim 123, wherein -B-L-T is selected from:






wherein Q, R²⁰, R¹³, E, and R⁶⁰ are as defined above; each methylene in any of the above formulae, other than those in a depicted ring, is independently optionally substituted with R²⁵; and R²⁵ is selected from halogen, trihalomethyl, oxo, -CN, -NO₂, -NH₂, -OR³, -NR³R³, -S(O)₀₋₂R³, -SO₂NR³R³, -CO₂R³, -C(O)NR³R³, -N(R³)SO₂R³, -N(R³)C(O)R³, -N(R³)CO₂R³, -C(O)R³, optionally substituted aryl, optionally substituted aryl C₁₋₆alkyl, heteroaryl C₁₋₆alkyl, and optionally substituted C₁₋₆alkyl; two of R²⁵, together with the carbon or carbons to which they are attached, can combine to form a three- to seven-membered optionally substituted alicyclic or heteroalicyclic.

125. The process according to claim 124, wherein Q is selected from the following three formula:




wherein R²⁰ is defined as above, and P is a five- to seven-membered ring, including the two shared carbons of the aromatic ring to which P is fused, P optionally containing between one and three heteroatoms.

126. The process according to claim 125, wherein -B-L-T is either of formula XXV or formula XXVI,




XXV




XXVI

wherein R^{20} is defined as above; G is either an optionally substituted cycloalkyl or an optionally substituted heteroalicyclic; each R^{30} is independently selected from halogen, trihalomethyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{OR}^3$, $-\text{NR}^3\text{R}^3$, $-\text{S}(\text{O})_{0-2}\text{R}^3$, $-\text{SO}_2\text{NR}^3\text{R}^3$, $-\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{NR}^3\text{R}^3$, $-\text{N}(\text{R}^3)\text{SO}_2\text{R}^3$, $-\text{N}(\text{R}^3)\text{C}(\text{O})\text{R}^3$, $-\text{N}(\text{R}^3)\text{CO}_2\text{R}^3$, $-\text{C}(\text{O})\text{R}^3$, and optionally substituted C_{1-6} alkyl; and R^{3a} and R^{3b} are each independently selected from $-\text{H}$ and optionally substituted C_{1-6} alkyl.


127. The process according to claim 126, wherein a compound of formula **XXIIa** is combined with a compound of formula **XXIIIa** to make a compound of formula **XXIa**,



XXIIa



XXIIIa



XXIa

wherein $-\text{B-L-T}$, Z, J, R^{50} , and R^2 are as defined above; R^{70} is selected from $-\text{H}$, $-\text{NO}_2$, $-\text{NH}_2$, and $-\text{NR}^3\text{R}^3$; provided when Z is $-\text{N}(\text{R}^5)-$ that R^5 is selected from $-\text{H}$, C_{1-3} alkyl, and aryl C_{1-3} alkyl; P^1 is selected from halogen, optionally substituted alkyl- $\text{S}(\text{O})_{0-2}-$, optionally substituted arylsulfonate, optionally substituted alkylsulfonate, a group containing boron, an azide, a group containing phosphorus, and a metal; and P^2 is selected from $-\text{H}$ and a metal.

128. The process according to claim 127, wherein P^2 is selected from $-\text{H}$, lithium, sodium, potassium, cesium, copper, palladium, and titanium.

129. The process according to claim 128, wherein Z is $-\text{O}-$.

130. The process according to claim 129, wherein P^1 is selected from chlorine, bromine, a toluene sulfonate, and trifluoromethanesulfonate.
131. The process according to claim 130, wherein R^{70} is -H.
132. The process according to claim 131, wherein J is =C(H)-.
133. The process according to claim 132, wherein R^2 is selected from C_{1-6} alkyl, perfluoro C_{1-6} alkyl, and halogen.
134. The process according to claim 133, wherein **XXIIa** and **XXIIIa** are heated together, optionally with a base, optionally with microwave radiation, to form **XXIa**.
135. The process according to claim 134, wherein the base is selected from an organic base, an inorganic base, and a combination of an organic base and an inorganic base.
136. The process according to claim 135, wherein the base is selected from 2,6-lutidine, 4-N,N-dimethylaminopyridine, and a metal carbonate.
137. The process according to claim 136, wherein **XXIIa** and **XXIIIa** are heated together in a solvent with said base, at between about 40°C and 200°C for between about one hour and twenty-four hours to form **XXIa**.
138. The process according to claim 137, wherein the solvent is an organic solvent.
139. The process according to claim 138, wherein one molar equivalent of **XXIIa** is combined with between about one quarter and four molar equivalents of **XXIIIa**.
140. The process according to claim 139, wherein one molar equivalent of **XXIIa** is combined with more than one but less than two molar equivalents of **XXIIIa**.
141. The process according to claim 140, wherein **XXIIa** is combined with **XXIIIa** and said base in an aromatic solvent to form a mixture, and said mixture is heated to between about 100°C and 200°C for between about one and ten hours to form **Ia**.
142. The process according to claim 141, wherein the aromatic solvent is an optionally substituted benzene.

143. The process according to claim 142, wherein the aromatic solvent is bromobenzene.
144. The process according to claim 143, wherein the base is 4-N,N-dimethylaminopyridine.
145. The process according to claim 144, wherein said mixture is heated to reflux for between about three and seven hours.
146. The process according to claim 145, wherein said mixture is heated to reflux for between about four and six hours.
147. The process according to claim 140, wherein **XXIIa** is combined with **XXIIIa** and said base in a non-aromatic solvent to form a mixture, and said mixture is heated to between about 40°C and 100°C for between about one and twenty hours to form **XXIa**.
148. The process according to claim 147, wherein the non-aromatic solvent comprises a functional group selected from an amide, and ether, a nitrile, a halide, an ester, an amine, and a ketone.
149. The process according to claim 148, wherein the non-aromatic solvent is N,N-dimethylacetamide.
150. The process according to claim 149, wherein the base is potassium carbonate.
151. The process according to claim 150, wherein said mixture is heated to about 50°C between about ten and twenty hours.
152. The process according to claim 151, wherein the aromatic solvent is an optionally substituted pyridine.
153. The process according to claim 152, wherein the aromatic solvent is 2,6-lutidine.
154. The process according to claim 153, wherein the base is 2,6-lutidine.
155. The process according to claim 154, wherein said mixture is heated to reflux for between about three and seven hours.

156. The process according to claim 155, wherein said mixture is heated to reflux for between about four and six hours.

157. The process according to claim 139, wherein one molar equivalent of **XXIIIa** is combined with more than one but less than two molar equivalents of **XXIIa**.

158. The process according to claim 157, wherein **XXIIa** is combined with **XXIIIa** and said base in an aromatic solvent to form a mixture, and said mixture is heated to between about 100°C and 200°C for between about ten and twenty hours to form **XXIa**.




159. The process according to claim 158, wherein the aromatic solvent is an optionally substituted pyridine.



160. The process according to claim 159, wherein the aromatic solvent is 2,6-lutidine.

161. The process according to claim 160, wherein the base is 2,6-lutidine.

162. The process according to claim 161, wherein said mixture is heated to between about 150°C and 200°C for between about fifteen and twenty hours.

163. The process according to any of claims 134 - 162, wherein a compound of formula **XXIIb** is substituted for the compound of formula **XXIIa**, and either a compound of formula **XXIIIb** or a compound of formula **XXIIIc** is substituted for the compound of formula **XXIIIa**, in order to make a compound of formula **XXIb** or a compound of formula **XXIc**, respectively,





wherein J, R⁵⁰, R²⁰ and R² are as defined above.

164. The process according to claim 163, wherein R², if present, is halogen.
165. The process according to claim 164, wherein R², if present, is fluorine.
166. The process according to claim 165, wherein R², if present, is up to two fluorines *ortho* to the oxygen of the phenylene to which R² is attached.
167. The process according to claim 115, used to make a compound listed in either Table 1 or Table 2.
168. The process according to any of claims 115 - 167, further comprising converting said compound to a pharmaceutically acceptable salt, hydrate, or prodrug thereof.